

Completing the Picture: Complexity of Graded Modal Logics with Converse*

BARTOSZ BEDNARCZYK

*Computational Logic Group, TU Dresden, Dresden, Germany and
Institute of Computer Science, University of Wrocław, Wrocław, Poland
(e-mail: bartosz.bednarczyk@cs.uni.wroc.pl)*

EMANUEL KIEROŃSKI and PIOTR WITKOWSKI

*Institute of Computer Science, University of Wrocław, Wrocław, Poland
(e-mails: emanuel.kieronski@cs.uni.wroc.pl, piotr.witkowski@cs.uni.wroc.pl)*

submitted 19 December 2019; revised 5 March 2021; accepted 10 March 2021

Abstract

A complete classification of the complexity of the local and global satisfiability problems for graded modal language over traditional classes of frames has already been established. By “traditional” classes of frames, we mean those characterized by any positive combination of reflexivity, seriality, symmetry, transitivity, and the Euclidean property. In this paper, we fill the gaps remaining in an analogous classification of the graded modal language with graded converse modalities. In particular, we show its NEXPTIME-completeness over the class of Euclidean frames, demonstrating this way that over this class the considered language is harder than the language without graded modalities or without converse modalities. We also consider its variation disallowing graded converse modalities, but still admitting basic converse modalities. Our most important result for this variation is confirming an earlier conjecture that it is decidable over transitive frames. This contrasts with the undecidability of the language with graded converse modalities.

KEYWORDS: modal logic, complexity, graded modalities, satisfiability

1 Introduction

For many years, modal logic has been an active topic in many academic disciplines, including philosophy, mathematics, linguistics, and computer science. Regarding applications in computer science, for example, in knowledge representation or verification, some important variations are those involving graded and converse modalities. In this paper, we investigate their computational complexity.

*We thank Evgeny Zolin for providing us a comprehensive list of gaps in the classification of the complexity of graded modal logics and for sharing with us his tikz files with modal cubes. We thank Emil Jeřábek for his explanations concerning $K5(\diamond, \diamond)$. We also thank Tomasz Gogacz and Filip Murlak for comments concerning Section 4. Finally, we thank the anonymous reviewers for their useful comments and remarks. Bartosz Bednarczyk is supported by Polish Ministry of Science and Higher Education program “Diamentowy Grant” no. DI2017 006447. Emanuel Kieroński and Piotr Witkowski are supported by Polish National Science Centre grant no. 2016/21/B/ST6/01444.




Fig. 1. Complexity of one-way modal logics. All bounds are tight. If local and global satisfiability differ in complexity, then “L:” indicates local and “G:” indicates global satisfiability.

By a modal logic, we will mean a pair $(\mathcal{L}, \mathcal{F})$, represented usually as $\mathcal{F}(\mathcal{L}^*)$, where \mathcal{L} is a modal language, \mathcal{F} is a class of frames, and \mathcal{L}^* is a short symbolic representation of \mathcal{L} (see the next paragraph), characterizing the modalities of \mathcal{L} . For example, $K4(\diamond_{\geq})$ will denote the graded modal logic of transitive frames.

While we are mostly interested in languages with graded and converse modalities, to set the scene we need to mention languages without them. Overall, the following five languages are relevant: the basic one-way modal language ($\mathcal{L}^* = \diamond$) containing only one, forward, modality \diamond ; graded one-way modal language ($\mathcal{L}^* = \diamond_{\geq}$) extending the previous one by graded forward modalities, $\diamond_{\geq n}$, for all $n \in \mathbb{N}$; two-way modal language ($\mathcal{L}^* = \diamond, \diamondleftarrow$) containing basic forward modality and the converse modality \diamondleftarrow ; graded two-way modal language ($\mathcal{L}^* = \diamond_{\geq}, \diamondleftarrow_{\geq}$) containing the forward modality, the converse modality, and their graded versions $\diamond_{\geq n}, \diamondleftarrow_{\geq n}$, for all $n \in \mathbb{N}$; and, additionally, a restriction of the latter without graded converse modalities but with basic converse modality ($\mathcal{L}^* = \diamond_{\geq}, \diamondleftarrow$).

The meaning of graded modalities is natural: $\diamond_{\geq n}\varphi$ means “ φ is true at no fewer than n successors of the current world,” and $\diamondleftarrow_{\geq n}\varphi$ means “ φ is true at no fewer than n predecessors of the current world.” We also recall that $\diamond\varphi$ means “ φ is true at some successor of the current world” and $\diamondleftarrow\varphi$ – “ φ is true at some predecessor of the current world.” Thus, for example, \diamond is equivalent to $\diamond_{\geq 1}$.

Our aim is to classify the complexity of the local (“in a world”) and global (“in all worlds”) satisfiability problems for all the logics obtained by combining any of the above languages with any class of frames from the so-called modal cube, that is, a class of frames characterized by any positive combination of the axioms of reflexivity (T), seriality (D), symmetry (B), transitivity (4), and the Euclidean property (5).

See Figure 1 for a visualization of the modal cube. Nodes of the depicted graph correspond to classes of frames and are labelled by letters denoting the above-mentioned properties, with S used in S4 and S5 for some historical reasons to denote reflexivity, and K denoting the class of all frames. If there is a path from a class X to a class Y , then it means that any class from Y also belongs to X (as all the axioms of X are also present in Y). Note that the modal cube contains only 15 classes, since some different combi-




Fig. 2. Complexities of two-way modal logics. All bounds are tight.

nations of the relevant axioms lead to identical classes, for example, reflexivity implies seriality, symmetry and transitivity imply Euclideaness, and so on.

A lot of work has been already done. The cases of basic one-way language and graded one-way language are completely understood, see Figure 1. The results for the former can be established using some standard techniques, see, for example, Blackburn *et al.* (2001) and the classical paper (Ladner 1977). The local satisfiability of the latter is systematically analyzed in Kazakov and Pratt-Hartmann (2009), with complexities turning out to lie between NP and NEXPTIME. As for its global satisfiability, some of the results follow from Kazakov and Pratt-Hartmann (2009), some are given in Zolin (2017), and the other can be easily obtained using again some standard techniques.

In the case of non-graded two-way modal language, over most relevant classes of frames, tight complexity bounds for local and global satisfiability are also known. The notable exceptions are global satisfiability problems of the logics of transitive frames, $K4(\diamond, \diamond)$, $S4(\diamond, \diamond)$, $D4(\diamond, \diamond)$, which are known to be in EXPTIME (due to a result in Demri and de Nivelle 2005 or due to a translation to description logic (DL) SI , whose satisfiability is in EXPTIME Tobies 2001b). However, according to the survey part of Zolin (2017), the corresponding lower bounds are missing. In the literature, we were also not able find a tight lower bound for the logics of Euclidean frames, $K5(\diamond, \diamond)$, $D5(\diamond, \diamond)$. We provide both missing bounds in Section 5, obtaining them by reductions from the acceptance problem for polynomially space bounded alternating Turing machines.¹ See the left part of Figure 2 for a complete complexity map in this case.

Let us now turn our attention to the most expressive two-way graded modal language with both graded forward and graded converse modalities (the right part of Figure 2). Its local and global satisfiability problems over the class of all frames (K) are known to be, resp., PSPACE-complete and EXPTIME-complete (see the survey part of Zolin 2017 and references therein). In Section 2.2, we explain how to obtain these bounds, as well as the same bounds in all cases involving neither transitivity nor Euclideaness. For the EXPTIME-bound, we employ the so-called *standard translation*. Over $K4$, $D4$, and $S4$,

¹ As explained to the first author by Emil Jeřábek, the latter bound can be alternatively proved by a reduction from TB, whose EXPTIME-hardness follows from Chen and Lin (1994).

the logics turn out to be undecidable (Zolin 2017). We remark that these are the only undecidable members of the whole family of logics considered in this paper. What remains are the classes of frames involving the Euclidean property. We solve them in Section 3. We prove that the logics $K5(\diamond_{\geq}, \diamond_{\geq})$ and $D5(\diamond_{\geq}, \diamond_{\geq})$ are locally and globally NEXPTIME-complete. Interestingly, this is a higher complexity than the EXPTIME-complexity of the language without graded modalities (Demri and de Nivelle 2005) and NP-complexity of the language without converse (Kazakov and Pratt-Hartmann 2009) over the same classes of frames. We also show that, when, additionally, transitivity is required, that is, for the logics $K45(\diamond_{\geq}, \diamond_{\geq})$ and $D45(\diamond_{\geq}, \diamond_{\geq})$, the complexity drops down to NP.

Finally, we consider the above-mentioned intermediate language $(\diamond_{\geq}, \diamond)$ in which we can count the successors, we have the basic converse modality, but we cannot count the predecessors. Our main result here, presented in Section 4, is demonstrating the (local and global) *finite model property* for the logics of transitive frames K4, D4, and S4: whenever a formula is (locally or, resp., globally) satisfiable, it is (locally, resp. globally) satisfiable over a finite frame. This implies the decidability of the (local and global) satisfiability problem (as well as the *finite* satisfiability problem, in which the attention is restricted to finite frames) for these logics and thus solve an open problem posed in Zolin (2017). An analogous problem was formulated also in the richer setting of DLs (Kazakov et al. 2007; Gutiérrez-Basulto et al. 2017), where the corresponding logic is called SIQ^- . That problem only recently was also positively solved (Gogacz et al. 2019). The results from Gogacz et al. (2019) (which we will discuss in more detail in a moment) allow us to derive the precise 2-EXPTIME-complexity bounds for the logics $K4(\diamond_{\geq}, \diamond)$, $D4(\diamond_{\geq}, \diamond)$ and $S4(\diamond_{\geq}, \diamond)$. The logics of the remaining classes of frames retain their complexities from the graded two-way case, so the picture is as in the right part of Figure 2, but the word “Undecidable” should be replaced by “2-EXPTIME.”

Due to a large number of papers in which the complexity bounds from Figures 1 and 2 are scattered, we have not referenced all of them in this introduction. Readers wishing to find an appropriate reference are recommended to use an online tool prepared by the first author ([bartoszjanbednarczyk.github.io/mlnavigator](https://github.com/bartoszanbednarczyk/mlnavigator)).

Related formalisms. Graded modalities are examples of *counting quantifiers* which are present in various formalisms. First of all, counting quantifiers were introduced for first-order logic: $\exists^{\geq n} x \varphi$ means: “at least n elements x satisfy φ ”. The satisfiability problem for some fragments of first-order logic with counting quantifiers was shown to be decidable. In particular, the two-variable fragment is NEXPTIME-complete (Pratt-Hartmann 2005), the two-variable guarded fragment is EXPTIME-complete (Pratt-Hartmann 2007), and the one-variable fragment is NP-complete (Pratt-Hartmann 2008). We will employ the second of those results in our paper.

Counting quantifiers are also present, in the form of the so-called *number restrictions*, in some DLs. As some standard DLs embed modal logics (c.f. a result in Baader et al. 2017, Section 2.6.2), results on DLs with number restrictions may be used to infer upper bounds on the complexity of some graded modal logics.

The DL which is particularly interesting from our point of view is the already mentioned logic SIQ^- . Syntactically, it can be seen as a *multi-modal* logic, that is, a logic whose frames interpret not just one but many accessibility relations, with different modalities associated with these relations. In the case of SIQ^- , each of the accessibility relations

can be independently required to be transitive or not. Recently, the *knowledge base satisfiability* problem for this logic was shown decidable and 2-EXPTIME-complete (Gogacz *et al.* 2019). As we said, from this result the decidability and 2-EXPTIME complexity of both local and global satisfiability of $K4(\diamond_{\geq}, \diamond)$, $S4(\diamond_{\geq}, \diamond)$, and $D4(\diamond_{\geq}, \diamond)$ can be inferred. Nevertheless, our proof of the finite model property for these logics remains valuable as in Gogacz *et al.* (2019) the decidability of the finite model reasoning for SIQ^- is left open (with the exception of the case in which there is only one accessibility relation and this relation is transitive; in this case, however, our finite model construction is used and cross-referred there).

In this context, it is worth noting that the logic $K(\diamond_{\geq}, \diamond)$ (with the accessibility relation not necessarily being transitive) and the logic $K4(\diamond_{\geq}^1, \diamond^1, \diamond_{\geq}^2, \diamond^2)$ (the bi-modal variant of $K4(\diamond_{\geq}, \diamond)$ with two independent transitive accessibility relations) do not have the *global* finite model property. Both these logics are contained in SIQ^- . An example $K(\diamond_{\geq}, \diamond)$ formula which is globally satisfiable (e.g., over an infinite binary tree with reversed edges) but has no finite models is $\diamond p \wedge \diamond \neg p \wedge \diamond_{\leq 1} \top$. This example can be easily adapted to $K4(\diamond_{\geq}^1, \diamond^1, \diamond_{\geq}^2, \diamond^2)$. On the other hand, $K(\diamond_{\geq}, \diamond)$ *does* have the *local* finite model property, as it is a fragment of the DL \mathcal{ALCIQ} , whose local finite model property was shown in Tobies (2001a). The status of the local finite model property for the multi-modal variants of $K4(\diamond_{\geq}, \diamond)$ is open.

Plan of the paper. In Section 2, we formally define the relevant modal languages and their semantics, recall the so-called standard translation, and use it to derive some initial results. In Sections 3 and 4, we investigate the classes of Euclidean frames and, respectively, transitive frames. Finally, in Section 5 we provide two lower bounds filling the gaps in the classification of the complexity of non-graded languages.

This work is an extended version of our conference paper (Bednarczyk *et al.* 2019).

2 Preliminaries

2.1 Languages, Kripke structures, and satisfiability

Let us fix a countably infinite set Π of *propositional variables*. The *language* of graded two-way modal logic is defined inductively as the smallest set of formulas containing Π , closed under Boolean connectives and, for any formula φ , containing $\diamond_{\geq n} \varphi$ and $\diamond_{\leq n} \varphi$, for all $n \in \mathbb{N}$. For a given formula φ , we denote its *length* with $|\varphi|$, and measure it as the number of symbols required to write φ , with numbers in subscripts $\geq n$ encoded in binary (i.e., encoding a number n requires $\log n$ bits rather than n bits).

The basic modality \diamond can be defined in terms of graded modalities: $\diamond \varphi := \diamond_{\geq 1} \varphi$. Analogously, for the converse modality: $\diamondleftarrow := \diamond_{\geq 1}$. Keeping this in mind, we may treat all languages mentioned in the introduction as fragments of the above-defined graded two-way modal language. We remark that we may also introduce other modalities, for example

$$\diamond_{\leq n} \varphi := \neg \diamond_{\geq n+1} \neg \varphi, \quad \diamondleftarrow_{\leq n} \varphi := \neg \diamondleftarrow_{\geq n+1} \neg \varphi, \quad \square \varphi := \neg \diamond \neg \varphi, \quad \text{and} \quad \boxplus \varphi := \neg \diamondleftarrow \neg \varphi.$$

The semantics is defined with respect to *Kripke structures*, that is, structures over the relational signature composed of unary predicates Π and with a binary predicate R . Such structures are represented as triples $\mathfrak{A} = \langle W, R, V \rangle$, where W is the *universe*, R is a binary

accessibility relation on W , and V is a function $V : \Pi \rightarrow \mathcal{P}(W)$ called *valuation*. Elements from the set W are often called *worlds*.

The *satisfaction relation* \models is defined inductively as follows:

- $\mathfrak{A}, w \models p$ iff $w \in V(p)$, for all $p \in \Pi$,
- $\mathfrak{A}, w \models \neg\varphi$ iff $\mathfrak{A}, w \not\models \varphi$ and similarly for the other Boolean connectives,
- $\mathfrak{A}, w \models \Diamond_{\geq n}\varphi$ iff there are $\geq n$ worlds $v \in W$ such that $\langle w, v \rangle \in R$ and $\mathfrak{A}, v \models \varphi$,
- $\mathfrak{A}, w \models \Diamond_{\geq n}\varphi$ iff there are $\geq n$ worlds $v \in W$ such that $\langle v, w \rangle \in R$ and $\mathfrak{A}, v \models \varphi$.

For a given Kripke structure $\mathfrak{A} = \langle W, R, V \rangle$, we call the pair $\langle W, R \rangle$ its *frame*. For a class of frames \mathcal{F} , we define the local (global) satisfiability problem of a modal language \mathcal{L} over \mathcal{F} (or equivalently for a modal logic $\mathcal{F}(\mathcal{L}^*)$) as follows: given a formula φ from a language \mathcal{L} , verify whether φ is satisfied at some world (all worlds) w of some structure \mathfrak{A} whose frame belongs to \mathcal{F} .

We announced in the introduction that we are interested in classes of frames characterized by any positive combination of the axioms of reflexivity (T), seriality (D), symmetry (B), transitivity (4), and the Euclidean property (5), recalled below.

- (D) seriality $\forall x\exists y(xRy)$
- (T) reflexivity $\forall x(xRx)$
- (B) symmetry $\forall xy(xRy \Rightarrow yRx)$
- (4) transitivity $\forall xyz(xRy \wedge yRz \Rightarrow xRz)$
- (5) Euclideaness $\forall xyz(xRy \wedge xRz \Rightarrow yRz)$

We say that a modal logic $\mathcal{F}(\mathcal{L}^*)$ has the *finite local (global) model property* if any formula of \mathcal{L} which is satisfied in some world (all worlds) of some structure from \mathcal{F} is also satisfied in some world (all worlds) of a *finite* structure from \mathcal{F} .

2.2 Standard translations

Modal logic can be seen as a fragment of first-order logic via the so-called *standard translation* (see, e.g., Blackburn et al. 2001). Here we present its variation tailored for graded and converse modalities and discuss how it can be used to establish exact complexity bounds for some of graded two-way modal logics.

In the forthcoming definition, we define a function \mathbf{st}_v for $v \in \{x, y\}$, which takes an input two-way modal logic formula φ and returns an equisatisfiable first-order formula. Definitions of \mathbf{st}_x and \mathbf{st}_y are symmetric; hence, we present the definition of \mathbf{st}_x only.

$$\mathbf{st}_x(p) = p(x) \text{ for all } p \in \Pi \tag{1}$$

$$\mathbf{st}_x(\varphi \wedge \psi) = \mathbf{st}_x(\varphi) \wedge \mathbf{st}_x(\psi) \text{ similarly for } \neg, \vee, \text{ etc.} \tag{2}$$

$$\mathbf{st}_x(\Diamond_{\geq n}\varphi) = \exists_{\geq n}.y(R(x, y) \wedge \mathbf{st}_y(\varphi)) \tag{3}$$

$$\mathbf{st}_x(\Diamond_{\geq n}\varphi) = \exists_{\geq n}.y(R(y, x) \wedge \mathbf{st}_y(\varphi)). \tag{4}$$

Translated formulas lie in the two-variable guarded fragment of first-order logic extended with counting quantifiers GC^2 . Observe that a modal formula $\varphi \in \mathcal{L}$ is (finitely) locally satisfiable iff a formula $\exists x \mathbf{st}_x(\varphi) \in \text{GC}^2$ is (finitely) satisfiable and that φ is (finitely) globally satisfiable iff $\forall x \mathbf{st}_x(\varphi) \in \text{GC}^2$ is (finitely) satisfiable. Since definitions of symmetry, seriality, and reflexivity, as recalled in the previous section, are GC^2 formulas, the standard translation can be used to provide a generic upper bound for the log-

ics $\mathcal{F}(\diamond_{\geq}, \diamond_{\geq})$ over all classes of frames \mathcal{F} involving neither transitivity nor Euclidean-ness. From the fact that the global satisfiability problem is EXPTIME-hard even for the basic modal language $\mathcal{F}(\diamond)$ (Blackburn and van Benthem 2007) and from EXPTIME-completeness of GC^2 (Pratt-Hartmann 2007), we conclude the following theorem.

Theorem 1

The global satisfiability problem for $\mathcal{F}(\diamond_{\geq}, \diamond_{\geq})$, where \mathcal{F} is any class of frames from the modal cube involving neither transitivity nor Euclidean-ness, is EXPTIME-complete.

For the local satisfiability problem, its complexity decreases to PSPACE. For two-way graded language over K, D, and T, we can simply adapt an existing tableaux algorithm by Tobies (Tobies 2001b), which yields a tight PSPACE bound. Moreover, if a class of frames is symmetric, then forward and converse modalities coincide and thus we may simply apply the result on graded one-way languages from Kazakov and Pratt-Hartmann (2009). The PSPACE lower bounds for the above-mentioned logics are inherited from the basic modal logic K (Ladner 1977) and hold even in the case of their propositional-variable-free fragment (Chagrov and Rybakov 2002). Thus, we can conclude the following.

Theorem 2

The local satisfiability problem for $\mathcal{F}(\diamond_{\geq}, \diamond_{\geq})$, where \mathcal{F} is any class of frames from the modal cube involving neither transitivity nor Euclidean-ness, is PSPACE-complete.

3 Euclidean frames: counting successors and predecessors

This section is dedicated to modal languages over the classes of frames satisfying Euclidean-ness. We demonstrate an exponential gap (NEXPTIME versus NP) in the complexities of modal logics over Euclidean frames (K5 and D5) and modal logics over transitive Euclidean frames (K45 and D45).

The two remaining Euclidean logics of our interest, namely KB45 and S5, whose frames are additionally symmetric, may be seen as one-way logics (as \diamond_{\geq} can be always replaced by \diamond_{\geq}). Hence, their NP upper bounds follow from previous works on one-way MLs (Kazakov and Pratt-Hartmann 2009). The lower bound is inherited from the Boolean satisfiability problem (Cook 1971). Thus,

Theorem 3 (Consequence of Kazakov and Pratt-Hartmann 2009.)

The local satisfiability and the global satisfiability problems for modal logics KB45($\diamond_{\geq}, \diamond_{\geq}$) and S5($\diamond_{\geq}, \diamond_{\geq}$) are NP-complete.

3.1 The shape of Euclidean frames

We start by describing the shape of frames under consideration. Let \mathfrak{A} be a *Euclidean structure*, that is, a Kripke structure $\mathfrak{A} = \langle W, R, V \rangle$ whose accessibility relation R satisfies the Euclidean property.

A world $w \in W$ is called a *lantern*, if $\langle w', w \rangle \notin R$ holds for every $w' \in W$. The set of all lanterns in \mathfrak{A} is denoted with $L_{\mathfrak{A}}$. We say that lantern $l \in W$ *illuminates* a world $w \in W$, if $\langle l, w \rangle \in R$ holds. The previous definition is lifted to the sets of worlds in an obvious way: a lantern l *illuminates a set of worlds* $I \subseteq W$ if l illuminates every world w from I .




Fig. 3. A Euclidean structure \mathfrak{A} with lanterns $L_{\mathfrak{A}} = \{l_1, l_2\}$.

We say that two worlds $w_1, w_2 \in W$ are *R-equivalent* (or simply *equivalent* if R is known from the context), if both $\langle w_1, w_2 \rangle \in R$ and $\langle w_2, w_1 \rangle \in R$ hold. The *R-clique* for a world w_1 in a structure \mathfrak{A} is the set $Q_{\mathfrak{A}}(w_1) \subseteq W$ consisting of w_1 together with all of its *R*-equivalent worlds. With $Q_{\mathfrak{A}}$ we denote the set $W \setminus L_{\mathfrak{A}}$ of *inner* (i.e., non-lantern) worlds. See Figure 3 for a drawing of an example Euclidean structure.

It is easy to observe that for any world $w_1 \in W$, all members of the clique $Q_{\mathfrak{A}}(w_1)$ are *R*-equivalent. This justifies why we have chosen the term “clique” to name such sets.

Observation 1

Any distinct worlds w', w'' from the *R*-clique $Q_{\mathfrak{A}}(w)$ of w are *R*-equivalent.

Proof

From the definition of *R*-equivalence, we know that both $\langle w, w' \rangle \in R$ and $\langle w, w'' \rangle \in R$ hold. Since the relation R satisfies the Euclidean property we infer that $\langle w', w'' \rangle \in R$ holds and $\langle w'', w' \rangle \in R$ holds, which implies *R*-equivalence of w and w' . \square

An immediate conclusion from the above observation is that the equality $Q_{\mathfrak{A}}(w) = Q_{\mathfrak{A}}(w_1)$ holds for any world $w \in Q_{\mathfrak{A}}(w_1)$. Thus, we will say that Q is an *R-clique in \mathfrak{A}* if the equality $Q = Q_{\mathfrak{A}}(w_1)$ holds for some (equivalently: for any) world $w_1 \in Q$.

As usual in modal logics, we can restrict our attention to *R-connected* models, that is, those models $\mathfrak{A} = \langle W, R, V \rangle$ for which $\langle W, R \cup R^{-1} \rangle$ is a connected graph. The following lemma describes the shape of Euclidean structures under consideration. It is very similar to Lemma 2 in Kazakov and Pratt-Hartmann (2009).

Lemma 1

If \mathfrak{A} is an *R*-connected structure over a Euclidean frame $\langle W, R \rangle$, then all worlds w in $Q_{\mathfrak{A}}$ are *reflexive* (i.e., $\langle w, w \rangle \in R$ holds) and $Q_{\mathfrak{A}}$ is an *R-clique*.

Proof

In the course of proof, we will refer to the formula (5) from Section 2.1, which defines Euclidean property. We will show that all worlds in $Q_{\mathfrak{A}}$ are reflexive and all worlds in $Q_{\mathfrak{A}}$ are *R*-equivalent. To show reflexivity take any $w \in Q_{\mathfrak{A}}$. By definition of $Q_{\mathfrak{A}}$, there exists $w' \in W$ such that $\langle w', w \rangle \in R$. Since \mathfrak{A} satisfies (5), by taking w' as x and w as both y and z in (5), we infer $\langle w, w \rangle \in R$.

To demonstrate R -equivalence, we will employ some simple observations. First, the relation $R \cap (Q_{\mathfrak{A}} \times Q_{\mathfrak{A}})$ is symmetric. To prove it take any $w_1, w_2 \in Q_{\mathfrak{A}}$ with $\langle w_1, w_2 \rangle \in R$. Then, use reflexivity of w_1 and the Euclidean property (with w_1 taken as x and z simultaneously, and w_2 as y in (5)) to infer $\langle w_2, w_1 \rangle \in R$. Second, $R \cap (Q_{\mathfrak{A}} \times Q_{\mathfrak{A}})$ is transitive. To prove it take any $w_1, w_2, w_3 \in Q_{\mathfrak{A}}$ with $\langle w_1, w_2 \rangle \in R$ and $\langle w_2, w_3 \rangle \in R$. Symmetry of $R \cap (Q_{\mathfrak{A}} \times Q_{\mathfrak{A}})$ gives us $\langle w_2, w_1 \rangle \in R$. Then, by the Euclidean property (with w_2 taken as x , w_3 taken as z , and w_1 as y in (5)) we infer $\langle w_1, w_3 \rangle \in R$. Third, if $\langle l, w_1 \rangle \in R$ and $\langle l, w_2 \rangle \in R$, for some $l \in L$ and $w_1, w_2 \in Q_{\mathfrak{A}}$, then $\langle w_1, w_2 \rangle \in R \cup R^{-1}$. This observation again simply follows from (5).

Now take any $w \in Q_{\mathfrak{A}}$. We will show that $Q_{\mathfrak{A}} = Q_{\mathfrak{A}}(w)$, that is, that $Q_{\mathfrak{A}}$ is the R -clique for w . Take any $w' \in Q_{\mathfrak{A}}$. We will show that both $\langle w, w' \rangle \in R$ and $\langle w', w \rangle \in R$. Since \mathfrak{A} is connected, there exists a $(R \cup R^{-1})$ -path from w to w' in \mathfrak{A} . By inductive application of the third observation above, we may assume that all elements of the path belong to $Q_{\mathfrak{A}}$. Then by the first observation (symmetry) we may assume that this is actually an R -path. Then, by the second observation (transitivity) the path reduces to a single edge $\langle w, w' \rangle \in R$. In the same way, we may show that $\langle w', w \rangle \in R$. Thus, all worlds in $Q_{\mathfrak{A}}$ are R -equivalent with w . Since all other worlds in \mathfrak{A} are lanterns, they cannot be R -equivalent with w . Thus, $Q_{\mathfrak{A}}$ is indeed the R -clique for w . \square

3.2 The universal modality

Before we start proving complexity results for the family of Euclidean logics, we show that global and local satisfiability problems are inter-reducible over any class of frames involving the Euclidean property.

Having restricted our attention to R -connected models, we will show that the *universal modality* \mathbf{U} can be defined in terms of standard (i.e., \diamond and \Box) modalities. Recall that the semantics of $\mathbf{U}\varphi$ is defined as follows: $\mathfrak{A}, w \models \mathbf{U}\varphi$, iff for every world x the condition $\mathfrak{A}, x \models \varphi$ holds. Taking a look at the shape of Euclidean structures (see, e.g., Lemma 1), it is not difficult to see that to propagate satisfaction of a given formula φ through the whole structure, and it is sufficient to first traverse all inner elements and from each of them propagate the satisfaction of φ to their predecessors. This intuition can be formalized by taking $\mathbf{U}\varphi := \varphi \wedge \Box \Box \Box \varphi$.

Lemma 2

Let $\mathfrak{A} = \langle W, R, V \rangle$ be an R -connected Euclidean structure. Then $\mathfrak{A}, w_0 \models \varphi \wedge \Box \Box \Box \varphi$ holds for some world $w_0 \in W$ iff $\mathfrak{A}, v \models \varphi$ holds for all worlds $v \in W$.

Proof

Let $\mathfrak{A} = \langle W, R, V \rangle$ be an R -connected Euclidean structure and let $\mathfrak{A}, w_0 \models \varphi \wedge \Box \Box \Box \varphi$ hold for some world $w_0 \in W$. We will show that it implies that φ is true in every world $w \in W$ (the opposite direction of the Lemma is trivial).

First, if $R = \emptyset$, then \mathfrak{A} is a singleton structure, because it is R -connected. In this case, the implication trivially holds. So, assume that $R \neq \emptyset$. Define $S = R \circ R \circ R^{-1}$. We will show that S is the universal relation $W \times W$. Indeed, take any $a, b \in W$. Then there exists $x \in Q_{\mathfrak{A}}$ such that $R(a, x)$ holds (if $a \in Q_{\mathfrak{A}}$ then, by Lemma 1, a is reflexive, so take $x = a$; if $a \in L_{\mathfrak{A}}$, such an x exists, since \mathfrak{A} is connected). Similarly, there exists $y \in Q_{\mathfrak{A}}$ such that $R(b, y)$. Now we have $R(x, y)$, since R is universal on $Q_{\mathfrak{A}}$ by Lemma 1. Thus, we

have $R(a, x)$, $R(x, y)$, and $R^{-1}(y, b)$, so $S(a, b)$ holds and thus $S = W \times W$. Therefore, $\mathfrak{A}, w_0 \models \square \square \exists \varphi$ implies $\mathfrak{A}, v \models \square \square \exists \varphi$, for any $v \in W$. \square

We now argue that the local and global satisfiability problems coincide for modal logics over Euclidean frames.

Lemma 3

Let $(\mathcal{L}, \mathcal{F})$ be a modal logic whose language contains \diamond and \diamondleftarrow and \mathcal{F} is a class of frames from the modal cube satisfying the Euclidean property. Then the global satisfiability problem for $\mathcal{F}(\mathcal{L}^*)$ is LOGSPACE reducible to the local satisfiability problem for $\mathcal{F}(\mathcal{L}^*)$ and vice versa.

Proof

As usual for modal logics, we may restrict to satisfiability over connected structures. Since \mathcal{F} is Euclidean and we have both \diamond , \diamondleftarrow at our disposal, we know that the universal modality \mathbf{U} is definable in $\mathcal{F}(\mathcal{L}^*)$ (see: Lemma 2). From the semantics of \mathbf{U} , we can immediately conclude that any modal formulas φ_l, φ_g the following equivalences hold: φ_l is locally satisfiable iff $\neg \mathbf{U} \neg \varphi_l$ is globally satisfiable and φ_g is globally satisfiable iff $\mathbf{U} \varphi_g$ is locally satisfiable. \square

3.3 The upper bound for graded two-way K5 and D5

This section is dedicated to the following theorem.

Theorem 4

The local and global satisfiability problems for Euclidean modal logics $\mathbf{K5}(\diamond_{\geq}, \diamondleftarrow_{\geq})$ and $\mathbf{D5}(\diamond_{\geq}, \diamondleftarrow_{\geq})$ are in NEXPTIME.

Proof

Note that here we may again restrict to satisfiability over connected frames. We start with the case of the class of all Euclidean frames $\mathbf{K5}$. We translate a given modal formula φ to the two-variable logic with counting \mathbf{C}^2 , in which both graded modalities and the shape of connected Euclidean structures, as defined in Lemma 1, can be expressed. Since satisfiability of \mathbf{C}^2 is in NEXPTIME (Pratt-Hartmann 2007), we obtain the desired conclusion. Recall the standard translation \mathbf{st} from Section 2.2. Let *lantern*(\cdot) be a new unary predicate and define $\varphi_{\mathbf{tr}}$ as

$$\mathbf{st}_x(\varphi) \wedge \forall x \forall y. (\neg \mathit{lantern}(x) \wedge \neg \mathit{lantern}(y) \rightarrow R(x, y)) \wedge (\mathit{lantern}(y) \rightarrow \neg R(x, y)).$$

Since $\mathbf{st}_x(\varphi)$ belongs to \mathbf{GC}^2 , $\varphi_{\mathbf{tr}}$ belongs to \mathbf{C}^2 (but not to \mathbf{GC}^2) and has one free variable x . Let \mathfrak{B} be a Kripke structure over a Euclidean frame. Expand \mathfrak{B} to a structure \mathfrak{B}^+ by setting $\mathit{lantern}^{\mathfrak{B}^+} = \{w \in \mathfrak{B} \mid w \in L_{\mathfrak{B}}\}$. Taking into account Lemma 1, a structural induction on φ easily establishes the following condition:

$$\mathfrak{B}, w_0 \models \varphi \text{ if and only if } \mathfrak{B}^+ \models \varphi_{\mathbf{tr}}[w_0/x] \text{ for every world } w_0 \in B.$$

Thus, a $\mathbf{K5}(\diamond_{\geq}, \diamondleftarrow_{\geq})$ formula φ is locally satisfiable if and only if the \mathbf{C}^2 formula $\exists_{\geq 1} x. \varphi_{\mathbf{tr}}$ is satisfiable, yielding a NEXPTIME algorithm for $\mathbf{K5}(\diamond_{\geq}, \diamondleftarrow_{\geq})$ local satisfiability. Membership of global satisfiability in NEXPTIME is implied by Lemma 2.

For the case of serial Euclidean frames, D5, it suffices to supplement the C^2 formula defined in the case of K5 with the conjunct $\exists x.(\neg \text{lantern}(x))$ expressing seriality. Correctness follows then from the simple observation that a Euclidean frame is serial iff it contains at least one non-lantern world (recall that all these worlds are reflexive). \square

3.4 Lower bounds for two-way graded K5 and D5

We now show a matching NEXPTIME-lower bound for the logics from the previous section. We concentrate on local satisfiability, but by Lemma 2 the results will hold also for global satisfiability. Actually, we obtain a stronger result, namely we show that the two-way graded modal logics K5 and D5 remain NEXPTIME-hard even if counting in one-way (either backward or forward) is forbidden. Hence, we show hardness of the logics $K5(\diamond_{\geq}, \diamond)$ and $D5(\diamond_{\geq}, \diamond)$. We recall that this gives a higher complexity than the EXPTIME-complexity of the language \diamond, \diamond (Demri and de Nivelles 2005) and NP-complexity of the language \diamond_{\geq} (Kazakov and Pratt-Hartmann 2009) over the same classes of frames.

In order to prove NEXPTIME-hardness of the Euclidean two-way graded modal logics K5 and D5, we employ a variant of the classical tiling problem, namely *exponential torus tiling problem* from Lutz (2002).

Definition 1 (4.15 from Lutz 2002)

A *torus tiling problem* \mathcal{P} is a tuple $(\mathcal{T}, \mathcal{H}, \mathcal{V})$, where \mathcal{T} is a finite set of tile types and $\mathcal{H}, \mathcal{V} \subseteq \mathcal{T} \times \mathcal{T}$ represent the horizontal and vertical matching conditions. Let \mathcal{P} be a tiling problem and $c = t_0, t_1, \dots, t_{n-1} \in \mathcal{T}^n$ an initial condition. A mapping $\tau : \{0, 1, \dots, 2^n - 1\} \times \{0, 1, \dots, 2^n - 1\} \rightarrow \mathcal{T}$ is a *solution* for \mathcal{P} and c if and only if, for all $i, j < 2^n$, the following holds $(\tau(i, j), \tau(i \oplus_{2^n} 1, j)) \in \mathcal{H}$, $(\tau(i, j), \tau(i, j \oplus_{2^n} 1)) \in \mathcal{V}$ and $\tau(0, i) = t_i$ for all $i < n$, where \oplus_i denotes addition modulo i . It is well known that there exists a NEXPTIME-complete torus tiling problem.

3.4.1 Outline of the proof

The proof is based on a polynomial time reduction from a torus tiling problem as in Definition 1. Henceforward, we assume that a NEXPTIME-complete torus tiling problem $\mathcal{P} = (\mathcal{T}, \mathcal{H}, \mathcal{V})$ is fixed. Let $c = t_0, t_1, \dots, t_{n-1} \in \mathcal{T}^n$ be its initial condition. We write a formula which is (locally) satisfiable iff $\langle \mathcal{P}, c \rangle$ has a solution. Each cell of the torus carries a *position* $\langle H, V \rangle \in \{0, 1, \dots, 2^n - 1\} \times \{0, 1, \dots, 2^n - 1\}$, encoded in binary in a natural way by means of propositional letters v_0, v_1, \dots, v_{n-1} and h_0, h_1, \dots, h_{n-1} , with h_0 and v_0 denoting the least significant bits. In the reduction, a single cell of the torus corresponds to a unique *inner*, that is, non-lantern, world. Since there are exactly $2^n \cdot 2^n$ cells, we enforce that also the total number of inner worlds is equal to $2^n \cdot 2^n$. We make use of graded modalities to specify that every inner world has exactly $2^n \cdot 2^n$ successors. We stress here that this is the only place where we employ counting. Thus, the proof works in the case where graded converse modalities are disallowed (but the basic converse modality will be necessary). Alternatively we could equivalently write that every inner world has exactly $2^n \cdot 2^n$ inner predecessors and obtain hardness of the language with graded converse modalities but without graded forward modalities.

Once we enforced a proper size of our torus, we must be sure that two distinct inner worlds carry different positions. We do this in two steps. We first write that a world with position $(0, 0)$ occurs in a model. For the second step, we assume that the grid is chessboard-like, that is, all elements are colored black or white in the same way as a chessboard is. Then, we say that every world is illuminated by four lanterns, where each of them propagates $\oplus_{2^n} 1$ relation on the proper axis (from a black node to a white one and vice versa). Finally, having the torus prepared we encode a solution for the given tiling by simply labelling each inner world with some tile letter t and ensure (from the vantage point of the lanterns) that any two horizontal or vertical neighbors do not violate the tiling constraints.

3.4.2 Encoding the exponential torus

Our goal is now to define a formula describing the exponential torus. The shape of the formula is as follows:

$$\varphi_{\text{torus}} \stackrel{\text{def}}{=} \varphi_{\text{firstCell}} \wedge \mathbf{U}(\varphi_{\text{partition}} \wedge \varphi_{\text{chessboard}} \wedge \varphi_{\text{torusSize}} \wedge \varphi_{\text{succ}})$$

where \mathbf{U} is the universal modality as in Lemma 2. The formula is going to say that: (i) the current world has position $(0, 0)$; (ii) every world is either a lantern or an inner world; (iii) the torus is chessboard-like, that is, its cells are colored with *blk* (black) and with *wht* (white) exactly as a real chessboard is; (iv) the overall size of the torus is equal to $2^n \cdot 2^n$; (v) each world of the torus has a proper vertical and a proper horizontal successor. The first four properties are straightforward to define:

$$\begin{aligned} \varphi_{\text{firstCell}} &\stackrel{\text{def}}{=} \text{inner} \wedge \text{wht} \wedge \bigwedge_{i=0}^{n-1} (\neg v_i \wedge \neg h_i) \\ \varphi_{\text{partition}} &\stackrel{\text{def}}{=} (\text{lantern} \leftrightarrow \neg \text{inner}) \wedge (\text{lantern} \leftrightarrow \neg \diamond \top) \\ \varphi_{\text{chessboard}} &\stackrel{\text{def}}{=} (\text{wht} \leftrightarrow \neg \text{blk}) \wedge (\text{wht} \leftrightarrow (v_0 \leftrightarrow h_0)) \\ \varphi_{\text{torusSize}} &\stackrel{\text{def}}{=} \text{inner} \rightarrow \diamond_{=2^n \cdot 2^n} \top \end{aligned}$$

Note that the formula $\varphi_{\text{torusSize}}$ indeed expresses (iv), as the set of all inner worlds forms a clique. The obtained formulas are of polynomial length since the number $2^n \cdot 2^n$ is encoded in binary.

What remains is to define φ_{succ} . For this, for every inner world we ensure that there exists a proper lantern responsible for establishing the appropriate successor relation. There will be four different types of such lanterns, denoted by symbols: *vbw*, *hbw*, *vwb*, *hwb*. The intuition is the following: the first letter *h* or *v* indicates whether a lantern is responsible for an *H*- or *V*-relation. The last two letters say whether a successor relation will be established between black and white worlds, or in the opposite way.

$$\begin{aligned} \varphi_{\text{succ}} &\stackrel{\text{def}}{=} (\text{lantern} \rightarrow \bigvee_{\heartsuit \in \{vbw, hbw, vwb, hwb\}} (\heartsuit \wedge \varphi_{\heartsuit})) \wedge \\ &\quad (\text{inner} \rightarrow \bigwedge_{\heartsuit \in \{vbw, hbw, vwb, hwb\}} \diamond (\text{lantern} \wedge \varphi_{\heartsuit})). \end{aligned}$$

It suffices to define formulas φ_{vbw} , φ_{hbw} , φ_{vwb} , and φ_{hwb} . Let us first define φ_{vbw} . The formula below, intended to be interpreted at a lantern, consists of three parts: (i)

the black and the white worlds illuminated by the lantern are pseudo-unique, that is, all white (respectively, black) worlds illuminated by the same lantern carry the same position; uniqueness will follow later from $\varphi_{\text{torusSize}}$; (ii) all black worlds illuminated by the lantern have the same H -position as all white worlds illuminated by this lantern; (iii) if V_w (respectively, V_b) encodes a V -position of the white (respectively, black) worlds illuminated by the lantern, then $V_w = V_b \oplus_{2^n} 1$. Let us define φ_{vbw} as:

$$\varphi_{vbw} \stackrel{\text{def}}{=} \varphi_{\text{pseudoUniqueness}} \wedge \varphi_{\text{equalH}} \wedge \varphi_{V_w=V_b \oplus_{2^n} 1}.$$

The definitions of the first and the second part of φ_{vbw} are simple:

$$\varphi_{\text{pseudoUniqueness}} \stackrel{\text{def}}{=} \bigwedge_{c \in \{wht, blk\}} \bigwedge_{p \in \{v, h\}} \bigwedge_{i=0}^{n-1} \diamond(c \wedge p_i) \rightarrow \square(c \wedge p_i)$$

$$\varphi_{\text{equalH}} \stackrel{\text{def}}{=} \bigwedge_{i=0}^{n-1} \diamond(blk \wedge h_i) \leftrightarrow \diamond(wht \wedge h_i)$$

Finally, we encode the \oplus_{2^n} -operation as the formula $\varphi_{V_w=V_b \oplus_{2^n} 1}$ by, a rather standard, implementation of binary addition. Below we distinguish two cases: when V_b is equal to $2^n - 1$ and when V_b is smaller than $2^n - 1$.

$$\begin{aligned} \varphi_{V_w=V_b \oplus_{2^n} 1} &\stackrel{\text{def}}{=} (\diamond(blk \wedge \bigwedge_{i=0}^{n-1} v_i) \rightarrow \diamond(wht \wedge \bigwedge_{i=0}^{n-1} \neg v_i)) \wedge \\ &\bigvee_{i=0}^{n-1} (\diamond(blk \wedge \neg v_i \wedge \bigwedge_{j=0}^{i-1} v_j) \wedge \diamond(wht \wedge v_i \wedge \bigwedge_{j=0}^{i-1} \neg v_j) \wedge \bigwedge_{j=i+1}^{n-1} \diamond(blk \wedge v_j) \leftrightarrow \diamond(wht \wedge v_j)). \end{aligned}$$

This completes the definition of φ_{vbw} . The following three definitions are analogous:

$$\varphi_{hbw} \stackrel{\text{def}}{=} \varphi_{\text{pseudoUniqueness}} \wedge \varphi_{\text{equalV}} \wedge \varphi_{H_w=H_b \oplus_{2^n} 1}$$

$$\varphi_{vwb} \stackrel{\text{def}}{=} \varphi_{\text{pseudoUniqueness}} \wedge \varphi_{\text{equalH}} \wedge \varphi_{V_b=V_w \oplus_{2^n} 1}$$

$$\varphi_{hwb} \stackrel{\text{def}}{=} \varphi_{\text{pseudoUniqueness}} \wedge \varphi_{\text{equalV}} \wedge \varphi_{H_b=H_w \oplus_{2^n} 1}.$$

The formula φ_{equalV} can be obtained from φ_{equalH} by replacing, for every i , the letter h_i with the letter v_i , and defining the formulas $\varphi_{H_w=H_b \oplus_{2^n} 1}$, $\varphi_{V_b=V_w \oplus_{2^n} 1}$, and $\varphi_{H_b=H_w \oplus_{2^n} 1}$ as simple modifications of $\varphi_{V_w=V_b \oplus_{2^n} 1}$. While modifying the mentioned formula, one should only switch blk and wht propositional symbols and possibly change v to h (when we consider adding $\oplus_{2^n} 1$ on the H axis).

The following lemma simply states that the formula φ_{torus} indeed defines a valid torus. Its proof is routine and follows directly from correctness of all presented formulas.

Lemma 4

Assume that the formula φ_{torus} is locally satisfied at a world w of a Euclidean structure $\mathfrak{A} = \langle W, R, V \rangle$. Then, set $Q_{\mathfrak{A}}(w)$, that is, the R -clique for w , contains exactly $2^n \cdot 2^n$ elements and each of them carries a different position $\langle H, V \rangle$, that is, there are no two worlds v, v' satisfying exactly the same h_i - and v_i -predicates.

Having defined a proper torus, it is quite easy to encode a solution to the torus tiling problem \mathcal{P} with the initial condition c . Each inner node will be labelled with a single tile

from \mathcal{T} and using appropriate lanterns we enforce that any two neighboring worlds do not violate the tiling rules \mathcal{H} and \mathcal{V} . This is the purpose of the formula φ_{tiling} defined below:

$$\varphi_{\text{tiling}} \stackrel{\text{def}}{=} \mathbf{U}(\varphi_{\text{tile}} \wedge \varphi_{\text{initCond}} \wedge \varphi_{\text{tilingRules}}).$$

The first conjunct specifies that each inner world is labelled with exactly one tile.

$$\varphi_{\text{tile}} \stackrel{\text{def}}{=} \text{inner} \rightarrow \left(\bigvee_{t \in \mathcal{T}} t \right) \wedge \bigwedge_{t, t' \in \mathcal{T}, t \neq t'} (\neg t \vee \neg t').$$

The second conjunct distributes the initial tiling among torus cells. To define it we use handy macros $V=k$ and $H=k$, with their intuitive meaning that the binary representation of the number k is written on atomic letters v_0, v_1, \dots, v_{n-1} and h_0, h_1, \dots, h_{n-1} , respectively. Thus,

$$\varphi_{\text{initCond}} \stackrel{\text{def}}{=} \bigwedge_{i=0}^{n-1} (\text{inner} \wedge H=0 \wedge V=i) \rightarrow t_i.$$

The last formula says that any two successive worlds do not violate tiling rules. Since any two neighbors are connected via a lantern, we describe the formula from the point of view of such lantern.

$$\begin{aligned} \varphi_{\text{tilingRules}} \stackrel{\text{def}}{=} & (\text{lantern} \wedge vbw \rightarrow \bigvee_{(t,t') \in \mathcal{V}} (\diamond(\text{blk} \wedge t) \wedge \diamond(\text{wht} \wedge t'))) \wedge \\ & (\text{lantern} \wedge vwb \rightarrow \bigvee_{(t',t) \in \mathcal{V}} (\diamond(\text{wht} \wedge t) \wedge \diamond(\text{blk} \wedge t'))) \wedge \\ & (\text{lantern} \wedge hbw \rightarrow \bigvee_{(t,t') \in \mathcal{H}} (\diamond(\text{blk} \wedge t) \wedge \diamond(\text{wht} \wedge t'))) \wedge \\ & (\text{lantern} \wedge hwb \rightarrow \bigvee_{(t',t) \in \mathcal{H}} (\diamond(\text{wht} \wedge t) \wedge \diamond(\text{blk} \wedge t))). \end{aligned}$$

In the following lemma, we claim that the presented reduction is correct. Its proof is once again routine and follows directly from correctness of all presented formulas.

Lemma 5

Let $\varphi_{\text{reduction}} \stackrel{\text{def}}{=} \varphi_{\text{torus}} \wedge \varphi_{\text{tiling}}$. The torus tiling problem instance $\langle \mathcal{P}, c \rangle$ has a solution if and only if the formula is $\varphi_{\text{reduction}}$ locally satisfiable.

Note that our intended models are serial. Thus, the result holds also for the logic D5. This gives the following theorem.

Theorem 5

The local and global satisfiability problems for the logics $\text{K5}(\diamond_{\geq}, \diamond)$ and $\text{D5}(\diamond_{\geq}, \diamond)$ are NEXPTIME-hard.

Together with Theorem 4, this gives:

Theorem 6

The local and global satisfiability problems for the logics $\text{K5}(\diamond_{\geq}, \diamond)$, $\text{K5}(\diamond_{\geq}, \diamond_{\geq})$ and for logics $\text{D5}(\diamond_{\geq}, \diamond)$, $\text{D5}(\diamond_{\geq}, \diamond_{\geq})$ are NEXPTIME-complete.

3.5 Transitive Euclidean frames

It turns out that the logics of transitive Euclidean frames have lower computational complexity. This is due to the following lemma.

Lemma 6

Let \mathfrak{A} be an R -connected structure over a transitive Euclidean frame $\langle W, R \rangle$. Then, every world $l \in L_{\mathfrak{A}}$ illuminates $Q_{\mathfrak{A}}$.

Proof

Take any world $q \in Q_{\mathfrak{A}}$. We will show that a lantern l illuminates q . Since l has no R -predecessor and \mathfrak{A} is R -connected, there exists a world $q' \in Q_{\mathfrak{A}}$ such that $\langle l, q' \rangle \in R$. By Lemma 1 set $Q_{\mathfrak{A}}$ is an R -clique, and thus we have $\langle q', q \rangle \in R$. By transitivity, we conclude that $\langle l, q \rangle \in R$. Thus, a lantern l illuminates $Q_{\mathfrak{A}}$. \square

A first-order formula stating that all non-lanterns are R -successors of all lanterns requires only two variables. Thus, as an immediate conclusion from Lemma 6, we can extend the translation developed in the previous section to handle the logic $K45(\diamond_{\geq}, \diamond_{\geq})$, and obtain a NEXPTIME-upper bound for the satisfiability problem. In fact, the shape of transitive Euclidean structures is so simple that two-variable logic is no longer necessary. Below we translate $K45(\diamond_{\geq}, \diamond_{\geq})$ and $D45(\diamond_{\geq}, \diamond_{\geq})$ to one-variable logic with counting C^1 , which is NP-complete (Pratt-Hartmann 2008).

Theorem 7

The local and the global satisfiability problems for transitive Euclidean modal logics $K45(\diamond_{\geq}, \diamond_{\geq})$ and $D45(\diamond_{\geq}, \diamond_{\geq})$ are in NP.

Proof

The proof is similar in spirit to the proof of Lemma 3 in Kazakov and Pratt-Hartmann (2009). Let $lantern(\cdot)$ be a new unary predicate. We first define translation function \mathbf{tr} that, given a $K45(\diamond_{\geq}, \diamond_{\geq})$ formula φ , produces an equisatisfiable C^1 formula $\mathbf{tr}(\varphi)$. We assume that all counting subscripts φ are non-zero.

$$\mathbf{tr}(p) = p(x) \text{ for all } p \in \Pi \tag{5}$$

$$\mathbf{tr}(\varphi \wedge \psi) = \mathbf{tr}(\varphi) \wedge \mathbf{tr}(\psi) \text{ similarly for } \neg, \vee, \text{ etc.} \tag{6}$$

$$\mathbf{tr}(\diamond_{\geq n} \varphi) = \exists_{\geq n}.x(\neg lantern(x) \wedge \mathbf{tr}(\varphi)) \tag{7}$$

$$\mathbf{tr}(\diamond_{\leq n} \varphi) = \exists_{\leq n}.x(\neg lantern(x) \wedge \mathbf{tr}(\varphi)) \tag{8}$$

$$\mathbf{tr}(\diamond_{\geq n} \varphi) = \neg lantern(x) \wedge \exists_{\geq n}.x(\mathbf{tr}(\varphi)) \tag{9}$$

$$\mathbf{tr}(\diamond_{\leq n} \varphi) = lantern(x) \vee \exists_{\leq n}.x(\mathbf{tr}(\varphi)). \tag{10}$$

Observe that $\mathbf{tr}(\varphi)$ is linear in the size of φ . Let \mathfrak{B} be a Kripke structure over a transitive Euclidean frame. Expand \mathfrak{B} to a structure \mathfrak{B}^+ by setting an interpretation of a symbol $lantern$ to be $lantern^{\mathfrak{B}^+} = \{w \in \mathfrak{B} \mid w \in L_{\mathfrak{B}}\}$. Taking into account Lemmas 1 and 6, a structural induction on φ easily establishes the following condition:

$$\mathfrak{B}, w_0 \models \varphi \text{ if and only if } \mathfrak{B}^+ \models \mathbf{tr}(\varphi)[w_0/x] \text{ for every world } w_0.$$

Thus, a $K45(\diamond_{\geq}, \diamond_{\geq})$ formula φ is locally satisfiable if and only if C^1 formula $\exists_{\geq 1}.x(\mathbf{tr}(x))$ is satisfiable, yielding an NP algorithm for $K45(\diamond_{\geq}, \diamond_{\geq})$ satisfiability. The algorithm

for $D45(\diamond_{\geq}, \diamond_{\geq})$ is obtained by just a slight update to the one given above. It suffices to supplement the C^1 formula defined in the case of $K45$ with the conjunct $\exists x.(\neg lantern(x))$ expressing seriality (cf. the proof of Theorem 4). \square

4 Transitive frames: counting successors, accessing predecessors

In this section, we consider the language $\diamond_{\geq}, \diamond$, that is, the modal language in which we can count the successors, but cannot count the predecessors, having at our disposal only the basic converse modality. Over all classes of frames involving neither transitivity nor Euclideaness, local satisfiability is PSPACE-complete and global satisfiability is EXPTIME-complete, as the tight lower and upper bounds can be transferred from, resp., the one-way non-graded language \diamond and the full two-way graded language. Over the classes of Euclidean frames $K5$ and $D5$, both problems are NEXPTIME-complete, as proved in Theorem 6. Over the classes of transitive Euclidean frames $KB45$, $K45$, $D45$, and $S5$, the problems are NP-complete, as the lower bound transfers from the language \diamond and the upper bound from the full two-way graded language (Theorem 7). So, over all the above-discussed classes of frames the complexities of $\diamond_{\geq}, \diamond$ and $\diamond_{\geq}, \diamond_{\geq}$ coincide.

What is left are the classes of transitive frames $K4$, $D4$, and $S4$. Recall that, in contrast to their one-way counterparts, the two-way graded logics of transitive frames $K4(\diamond_{\geq}, \diamond_{\geq})$, $D4(\diamond_{\geq}, \diamond_{\geq})$, and $S4(\diamond_{\geq}, \diamond_{\geq})$ are undecidable (Zolin 2017). In Zolin (2017), the question is asked if the decidability is regained when the language is restricted to $\diamond_{\geq}, \diamond$. Here we answer this question, demonstrating the local and global finite model property for the obtained logics; this implies that their satisfiability problems are indeed decidable.

In Lemma 5.5 from Zolin (2017), it is shown that over the class of transitive frames the global satisfiability and local satisfiability problems for the considered language are polynomially equivalent. Moreover, they are polynomially equivalent to the *combined* satisfiability problem, asking if for a given pair of formulas ϕ, ϕ' there exists a structure in which ϕ is true at every world and ϕ' is true at some world. The remark following the proof of that lemma says that it holds also for reflexive transitive frames. The same can be easily shown also for serial transitive frames. We thus have:

Lemma 7

For each of the logics $K4(\diamond_{\geq}, \diamond)$, $D4(\diamond_{\geq}, \diamond)$, and $S4(\diamond_{\geq}, \diamond)$, their global, local, and combined satisfiability problems are polynomially equivalent.

Below we explicitly deal with global satisfiability. The above lemma implies, however, that our results apply also to local satisfiability.

Let us concentrate on the class $K4$ of all transitive frames. The finite model construction we are going to present is the most complicated part of this paper. It begins similarly to the exponential model construction in the case of local satisfiability of $K4(\diamond_{\geq})$ from Kazakov and Pratt-Hartmann (2009): we introduce a Scott-type normal form (Lemma 8) and then generalize two pieces of model surgery used there (Lemma 9) to our setting: starting from any model, we first obtain a model with short *paths of cliques* and then we decrease the size of the cliques. Some modifications of the constructions from Kazakov and Pratt-Hartmann (2009) are necessary to properly deal with the converse modality; they are, however, rather straightforward. Having a model with short paths of cliques and

small cliques, we develop some new machinery of *clique profiles* and *clique types* allowing us to decrease the overall size of the structure; this fragment is our main contribution.

Lemma 8

Given a formula φ of the language $(\diamond_{\geq}, \diamond)$, we can compute in polynomial time a formula ψ of the form

$$\eta \wedge \bigwedge_{1 \leq i \leq l} (p_i \rightarrow \diamond_{\geq C_i} \pi_i) \wedge \bigwedge_{1 \leq i \leq m} (q_i \rightarrow \diamond_{\leq D_i} \chi_i) \wedge \bigwedge_{1 \leq i \leq l'} (p'_i \rightarrow \diamond \pi'_i) \wedge \bigwedge_{1 \leq i \leq m'} (q'_i \rightarrow \Box \neg \chi'_i), \quad (11)$$

where the p_i, q_i, p'_i, q'_i are propositional variables, the C_i, D_i are natural numbers, and η and the $\pi_i, \chi_i, \pi'_i, \chi'_i$ are propositional formulas, such that φ and ψ are globally satisfiable over exactly the same transitive frames.

Proof

Follows by a routine renaming process, which is similar to the proof of Lemma 4 from Kazakov and Pratt-Hartmann (2009). \square

Next, let us introduce some helpful terminology, copying it mostly from the above-mentioned paper (Kazakov and Pratt-Hartmann 2009). Let $\mathfrak{A} = \langle W, R, V \rangle$ be a transitive structure, and $w_1, w_2 \in W$. We say that w_2 is an *R-successor* of w_1 if $\langle w_1, w_2 \rangle \in R$; w_2 is a *strict R-successor* of w_1 if $\langle w_1, w_2 \rangle \in R$, but $\langle w_2, w_1 \rangle \notin R$; w_2 is a *direct R-successor* of w_1 if w_2 is a strict *R-successor* of w_1 and, for every $w \in W$ such that $\langle w_1, w \rangle \in R$ and $\langle w, w_2 \rangle \in R$ we have either $w \in Q_{\mathfrak{A}}(w_1)$ or $w \in Q_{\mathfrak{A}}(w_2)$. Recall that $Q_{\mathfrak{A}}(w)$ denotes the *R-clique* for w in \mathfrak{A} .

The *depth* of a structure \mathfrak{A} is the maximum over all $k \geq 0$ for which there exist worlds $w_0, \dots, w_k \in W$ such that w_i is a strict *R-successor* of w_{i-1} for every $1 \leq i \leq k$, or ∞ if no such a maximum exists. The *breadth* of \mathfrak{A} is the maximum over all $k \geq 0$ for which there exist worlds w, w_1, \dots, w_k such that w_i is a direct *R-successor* of w for every $1 \leq i \leq k$, and the sets $Q_{\mathfrak{A}}(w_1), \dots, Q_{\mathfrak{A}}(w_k)$ are disjoint, or ∞ if no such a maximum exists. The *width* of \mathfrak{A} is the smallest k such that $k \geq |Q_{\mathfrak{A}}(w)|$ for all $w \in W$, or ∞ if no such k exists.

Lemma 9

Let φ be a normal form formula as in equation (11). If φ is globally satisfied in a transitive model \mathfrak{A} , then it is globally satisfied in a transitive model \mathfrak{A}' with depth $d' \leq (\sum_{i=1}^m D_i) + m + m' + 1$ and width $c' \leq (\sum_{i=1}^l C_i) + l' + 1$.

Proof

The proof is a construction being a minor modification of Stages 1 and 4 of the construction from the proof of Lemma 6 in Kazakov and Pratt-Hartmann (2009), where the language without backward modalities is considered. We closely follow the lines of Kazakov and Pratt-Hartmann's construction, just taking additional care of backward witnesses. We remark here that also Stage 2 of the above-mentioned construction could be adapted, giving a better bound on the depth of \mathfrak{A}' . We omit it here since such an improvement would not be crucial for our purposes. Stage 3 cannot be directly adapted.

Let us turn to the detailed proof.

Stage 1. Small depth. Let $\mathfrak{A} = \langle W, R, V \rangle$. For $w \in W$ define $d_{\mathfrak{A}}^i(w) := \min(D_i + 1, |\{w' : \mathfrak{A}, w' \models \chi_i, \langle w, w' \rangle \in R^*\}|)$ where D_i and χ_i , $1 \leq i \leq m$, are as in Equation 11 and R^* is the reflexive closure of R . We also define $S_{\mathfrak{A}}(w) := \{\chi'_i : \text{there is } w' \text{ such that } \mathfrak{A}, w' \models \chi'_i \text{ and } \langle w', w \rangle \in R^*\}$, where χ'_i , $1 \leq i \leq m'$ are also as in equation (11).

Let $R_{\sim} := \{\langle w_1, w_2 \rangle \in R : d_{\mathfrak{A}}^i(w_1) = d_{\mathfrak{A}}^i(w_2) \text{ for all } 1 \leq i \leq m \text{ and } S_{\mathfrak{A}}(w_1) = S_{\mathfrak{A}}(w_2)\}$ be the restriction of R to pairs of worlds that have the same values of the $d_{\mathfrak{A}}^i$ and $S_{\mathfrak{A}}$. Let R_{\sim}^{-} be the inverse of R_{\sim} . Let $\mathfrak{A}' = \langle W, R', V \rangle$ be obtained from $\mathfrak{A} = \langle W, R, V \rangle$ by setting $R' := (R \cup R_{\sim}^{-})^+$, where the superscript $+$ is the transitive closure operator. Intuitively, if w_1 is R -reachable from w_2 and w_1 and w_2 agree on the number (up to the limit of D_i) of the worlds satisfying χ_i reachable from them, for all $1 \leq i \leq m$, and, for all i , w_1 is an R -successor of a world satisfying χ'_i iff w_2 is, then we make w_1 and w_2 R' -equivalent. The effect is that some R -cliques of \mathfrak{A} are joined into bigger R -cliques in \mathfrak{A}' . We show that \mathfrak{A}' satisfies φ and has appropriately bounded depth.

For every $w_1, w_2 \in W$ such that w_2 is a strict R' -successor of w_1 , we have $d_{\mathfrak{A}}^i(w_1) \geq d_{\mathfrak{A}}^i(w_2)$ for all $1 \leq i \leq m$, $S_{\mathfrak{A}}(w_1) \subseteq S_{\mathfrak{A}}(w_2)$ and either $d_{\mathfrak{A}}^i(w_1) > d_{\mathfrak{A}}^i(w_2)$ for some i , and thus $\sum_{i=1}^m d_{\mathfrak{A}}^i(w_1) > \sum_{i=1}^m d_{\mathfrak{A}}^i(w_2)$ or the inclusion $S_{\mathfrak{A}}(w_1) \subseteq S_{\mathfrak{A}}(w_2)$ is strict. Since $d_{\mathfrak{A}}^i(w) \leq D_i + 1$ for every $w \in W$ and every $1 \leq i \leq m$, and the size of $S_{\mathfrak{A}}(w)$ is bounded by m' , the length of every chain w_0, \dots, w_k such that w_i is a strict R' -successor of w_{i-1} is bounded by $(\sum_{i=1}^m D_i) + m + m' + 1$.

In order to prove that $\mathfrak{A}' \models \varphi$, we first prove that $d_{\mathfrak{A}}^i(w) = d_{\mathfrak{A}'}^i(w)$ for every $w \in W$ and $1 \leq i \leq m$. Assume to the contrary that $d_{\mathfrak{A}}^i(w) \neq d_{\mathfrak{A}'}^i(w)$ for some $w \in W$ and some i . Since $R \subseteq R'$, we have $d_{\mathfrak{A}}^i(w) < d_{\mathfrak{A}'}^i(w) \leq D_i + 1$, which means, in particular, that there exists an element $w' \in W$ with $\mathfrak{A}, w' \models \chi_i$, such that $\langle w, w' \rangle \in R'$ but $\langle w, w' \rangle \notin R$.

Since $\langle w, w' \rangle \in R'$, by definition of R' , there exists a sequence w_0, \dots, w_k of different worlds in W such that $w_0 = w$, $w_k = w'$ and $\langle w_{j-1}, w_j \rangle \in R \cup R_{\sim}^{-}$ for every $1 \leq j \leq k$. Note that $d_{\mathfrak{A}}^i(w_{j-1}) \geq d_{\mathfrak{A}}^i(w_j)$ for every $1 \leq j \leq k$ and every $1 \leq i \leq m$. Take the maximal j such that $\langle w_{j-1}, w' \rangle \notin R$. Since $\langle w_0, w' \rangle = \langle w, w' \rangle \notin R$, such a maximal j always exists. Then $\langle w_j, w' \rangle \in R^*$, and $\langle w_{j-1}, w_j \rangle \notin R$. Since $\langle w_{j-1}, w_j \rangle \in R \cup R_{\sim}^{-}$, we have $\langle w_{j-1}, w_j \rangle \in R_{\sim}^{-}$, and so $d_{\mathfrak{A}}^i(w_{j-1}) = d_{\mathfrak{A}}^i(w_j)$ by definition of R_{\sim} . Since $d_{\mathfrak{A}}^i(w_j) \leq d_{\mathfrak{A}}^i(w_0) = d_{\mathfrak{A}}^i(w) < D_i + 1$, we obtain a contradiction, due to the fact that $d_{\mathfrak{A}}^i(w_{j-1}) = d_{\mathfrak{A}}^i(w_j) \leq D_i$, $\langle w_{j-1}, w \rangle \notin R^*$, $\langle w_j, w' \rangle \in R^*$, $\langle w_j, w_{j-1} \rangle \in R$, and $\mathfrak{A}, w' \models \chi_i$.

The observation that $S_{\mathfrak{A}}(w) = S_{\mathfrak{A}'}(w)$ for all $w \in W$ is even simpler. Assume to the contrary that this equality does not hold for some $w \in W$. This means that $\chi'_i \in S_{\mathfrak{A}'}(w)$ and $\chi'_i \notin S_{\mathfrak{A}}(w)$ for some $1 \leq i \leq m'$. In particular, there exists an element $w' \in W$ with $\mathfrak{A}, w' \models \chi'_i$, such that $\langle w', w \rangle \in R'$ but $\langle w', w \rangle \notin R$. Thus, there is a sequence of different worlds $w' = w_0, \dots, w_k = w$ such that $\langle w_{j-1}, w_j \rangle \in R \cup R_{\sim}^{-}$ for every $1 \leq j \leq k$. Note that $S_{\mathfrak{A}}(w_{j-1}) \subseteq S_{\mathfrak{A}}(w_j)$ for every $1 \leq j \leq k$. Since $\chi'_i \in S_{\mathfrak{A}}(w_0)$ it follows that $\chi'_i \in S_{\mathfrak{A}}(w_k)$. Contradiction.

To complete the proof that $\mathfrak{A}' \models \varphi$ we demonstrate that if ψ is any conjunct of φ and $w \in W$, then $\mathfrak{A}, w \models \psi$ implies $\mathfrak{A}', w \models \psi$. Indeed, for the propositional formula η it is immediate. For subformulas $(p_i \rightarrow \diamond_{\geq C_i} \pi_i)$ and $(p'_i \rightarrow \diamond \pi'_i)$, this holds since $R \subseteq R'$. For subformulas $(q_i \rightarrow \diamond_{\leq D_i} \chi_i)$, this follows from the property $d_{\mathfrak{A}}^i(w) = d_{\mathfrak{A}'}^i(w)$. Finally, for subformulas $(q'_i \rightarrow \boxminus \neg \chi'_i)$ this follows from the property $S_{\mathfrak{A}}(w) = S_{\mathfrak{A}'}(w)$.

Stage 2. Small width. By Stage 1, we may assume that \mathfrak{A} has depth bounded by $(\sum_{i=1}^m D_i) + m + m' + 1$. For every element $w \in W$, we define $Q_{\pi_i}(w)$ to be the set of elements of $Q_{\mathfrak{A}}(w)$ for which π_i holds ($1 \leq i \leq l$) and $Q_{\pi'_i}(w)$ to be the set of elements of $Q_{\mathfrak{A}}(w)$ for which π'_i holds ($1 \leq i \leq l'$). We call the elements of each $Q_{\pi}(w)$ the *equivalent π -witnesses for w* . Note that for each relevant π we have $Q_{\pi}(w_1) = Q_{\pi}(w_2)$ when w_1 and w_2 are R -equivalent. For $1 \leq i \leq l$, let $Q'_{\pi_i}(w)$ be $Q_{\pi_i}(w)$ if $|Q_{\pi_i}(w)| \leq C_i$, or, otherwise, a subset of $Q_{\pi_i}(w)$ which contains exactly C_i elements. We call $Q'_{\pi_i}(w)$ the *selected equivalent π_i -witnesses for w* . For $1 \leq i \leq l'$, let $Q'_{\pi'_i}(w)$ be $Q_{\pi'_i}(w)$ if $|Q_{\pi'_i}(w)| \leq 1$, or, otherwise, a singleton subset of $Q_{\pi'_i}(w)$. We call $Q'_{\pi'_i}(w)$ the *selected equivalent π'_i -witness for w* . Additionally, define $Q'_*(w)$ to be any singleton subset of $Q_{\mathfrak{A}}(w)$. We assume that if w_1 and w_2 are R -equivalent then $Q'_{\pi_i}(w_1) = Q'_{\pi_i}(w_2)$ for all $1 \leq i \leq l$, $Q'_{\pi'_i}(w_1) = Q'_{\pi'_i}(w_2)$ for $1 \leq i \leq l'$, and $Q'_*(w_1) = Q'_*(w_2)$. Define the structure $\mathfrak{A}' = \langle W', R', V' \rangle$ by setting $W' = \bigcup_{w \in W, 1 \leq i \leq l} Q'_{\pi_i}(w) \cup \bigcup_{w \in W, 1 \leq i \leq l'} Q'_{\pi'_i}(w) \cup Q'_*(w)$, $R' := R \upharpoonright W'$, and $V' = V \upharpoonright W'$. Intuitively \mathfrak{A}' is obtained from \mathfrak{A} by removing elements in every R -clique, except for those that are selected witnesses for other elements or are members of the singleton set Q_* , guaranteeing that the clique will remain non-empty. It is not difficult to see that \mathfrak{A}' has the required properties. In particular, our selection process selects at most $(\sum_{i=1}^l C_i) + l' + 1$ elements in every R -clique. \square

To describe our next step, we need a few more definitions. Given a world w of a structure \mathfrak{A} , we define its *depth* as the maximum over all $k \geq 0$ for which there exist worlds $w = w_0, \dots, w_k \in W$ such that w_i is a strict R -successor of w_{i-1} for every $1 \leq i \leq k$, or as ∞ if no such a maximum exists. For an R -clique Q , we define its *depth* as the depth of w for any $w \in Q$; this definition is sound since for all $w_1 \in Q_{\mathfrak{A}}(w)$ the depth of w is equal to the depth of w_1 .

From this point, we will mostly work on the level of cliques rather than individual worlds. We may view any structure \mathfrak{A} as a partially ordered set of cliques. We write $\langle Q_1, Q_2 \rangle \in R$, and say that a clique Q_1 *sends* an edge to a clique Q_2 (or that Q_2 *receives* an edge from Q_1) if $\langle w_1, w_2 \rangle \in R$ for any (equivalently: for all) $w_1 \in Q_1, w_2 \in Q_2$.

A 1-type of a world w in \mathfrak{A} is the set of all propositional variables p such that $\mathfrak{A} \models p$. We sometimes identify a 1-type with the conjunction of all its elements and negations of variables it does not contain. Given a natural number k , a structure \mathfrak{A} , and a clique Q in this structure \mathfrak{A} , we define a k -*profile* of Q (called just a *profile* if k is clear from the context) in \mathfrak{A} as the tuple $prof_{\mathfrak{A}}^k(Q) = (\mathcal{H}, \mathcal{A}, \mathcal{B}, irref)$, where \mathcal{H} is the multiset of 1-types in which the number of copies of each 1-type α equals $\min(k + 1, |\{w \in Q : \mathfrak{A}, w \models \alpha\}|)$, \mathcal{A} is the multiset of 1-types in which the number of copies of each 1-type α equals $\min(k, |\{w : \mathfrak{A}, w \models \alpha \text{ and } w \text{ is a strict } R\text{-successor of a world from } Q\}|)$, \mathcal{B} is the set of 1-types of worlds for which a world from Q is its strict R -successor, and *irref* is a Boolean variable set to 1 iff the clique consists of a single irreflexive element (note that if the clique contains at least two elements then they all must be reflexive). Intuitively, \mathcal{H} counts (up to $k + 1$) realizations of 1-types (*H*)ere in Q , \mathcal{A} counts (up to k) realizations 1-types (*A*)bove Q , and \mathcal{B} says which 1-types appear (*B*)elow Q . Usually, given a normal form φ as in equation (11), we will be interested in M_{φ} -profiles of cliques, where $M_{\varphi} = \max(\{C_i\}_{i=1}^l \cup \{D_i + 1\}_{i=1}^m)$. Note that, given the M_{φ} -profiles of all cliques in a structure we are able to determine whether this structure is a global model of φ . Indeed, given the M_{φ} -profile of a clique we know

the 1-types of elements it contains, for each such element we can count, at least up to M_φ , how many successors of each 1-type it has (for this we use the values of \mathcal{H} , \mathcal{A} , and *irref*), and for each element we know the set of 1-types of its predecessors (for this we use the values of \mathcal{H} , \mathcal{B} and *irref*). Clearly, this information is sufficient to check if every conjunct of φ is satisfied. The following observation is also straightforward.

Lemma 10

If $\mathfrak{A} \models \varphi$ for a normal form φ , and if in a structure \mathfrak{A}' the M_φ -profile of every clique is equal to the M_φ -profile of some clique from \mathfrak{A} , then $\mathfrak{A}' \models \varphi$.

We now prove the finite model property.

Lemma 11

Let φ be a normal form formula. If φ is globally satisfied in a transitive model \mathfrak{A} , then it is globally satisfied in a finite transitive model \mathfrak{A}' .

Proof

Construction of \mathfrak{A}' . We assume that φ is as in equation (11). By Lemma 9, we may assume that $\mathfrak{A} = \langle W, R, V \rangle$ has depth $d \leq (\sum_{i=1}^m D_i) + m + m' + 1$ and width $c \leq (\sum_{i=1}^l C_i) + l' + 1$. Note that \mathfrak{A} may be infinite due to possibly infinite breadth.

Let us split W into sets U_0, \dots, U_d with U_i consisting of all elements of W of depth i in \mathfrak{A} (equivalently speaking: being the union of all cliques of depth i in \mathfrak{A}). They are called *layers*. Note that cliques from U_i may send R -edges only to cliques from U_j with $j < i$.

We now inductively define a sequence of models $\mathfrak{A} = \mathfrak{A}_{-1}, \mathfrak{A}_0, \dots, \mathfrak{A}_d = \mathfrak{A}'$, with $\mathfrak{A}_i = \langle W_i, R_i, V_i \rangle$ such that

- $W_i = U'_0 \cup \dots \cup U'_i \cup U_{i+1} \cup \dots \cup U_d$, where each U'_i is a finite union of some cliques from U_i ,
- $V_i = V \upharpoonright W_i$
- $\mathfrak{A}_i \upharpoonright (U'_0 \cup \dots \cup U'_i) = \mathfrak{A}_{i-1} \upharpoonright (U'_0 \cup \dots \cup U'_i)$,
- $\mathfrak{A}_i \upharpoonright (U'_0 \cup \dots \cup U'_{i-1} \cup U_{i+1} \cup \dots \cup U_d) = \mathfrak{A}_{i-1} \upharpoonright (U'_0 \cup \dots \cup U'_{i-1} \cup U_{i+1} \cup \dots \cup U_d)$
- in particular: $\mathfrak{A}_i \upharpoonright (U_{i+1} \cup \dots \cup U_d) = \mathfrak{A} \upharpoonright (U_{i+1} \cup \dots \cup U_d)$.

We obtain \mathfrak{A}_i from \mathfrak{A}_{i-1} by distinguishing a fragment U'_i of U_i , removing $U_i \setminus U'_i$ and adding some edges from $U_{i+1} \cup \dots \cup U_d$ to U'_i ; all the other edges remain untouched. We do it carefully, to avoid modifications of the profiles of the surviving cliques. Let us describe the process of constructing \mathfrak{A}_i in detail.

Assume $i \geq 0$. We first distinguish a finite subset U'_i of U_i . We define a *clique type* of every clique Q from U_i in \mathfrak{A}_{i-1} as a triple $(\mathcal{H}, \mathcal{B}, S)$, where \mathcal{H} and \mathcal{B} are as in $prof_{\mathfrak{A}_{i-1}}^{M_\varphi}(Q)$ and S is the subset of cliques from $U'_0 \cup \dots \cup U'_{i-1}$, consisting of those cliques to which Q sends an R_{i-1} -edge. We stress that during the construction of \mathfrak{A}_i , the clique types of cliques are always computed in \mathfrak{A}_{i-1} . In particular, S is empty for $i = 0$ and, as we will always have that $U'_0 \cup \dots \cup U'_{i-1}$ is finite, S is finite for any $i > 0$. Thus, for each i there will be only finitely many clique types.

For every clique type β realized in U_i , we mark M_φ cliques of this type, or all such cliques if there are less than M_φ of them. Let U'_i be the union of the marked cliques. We fix some arbitrary numbering of the marked cliques.

Now we define the relation R_i . As said before, for any pair of cliques Q_1, Q_2 both of which are contained in $U'_0 \cup \dots \cup U'_{i-1} \cup U_{i+1} \cup \dots \cup U_d$ or in $U'_0 \cup \dots \cup U'_i$, we set $\langle Q_1, Q_2 \rangle \in R_i$ iff $\langle Q_1, Q_2 \rangle \in R_{i-1}$. It remains to define the R_i -edges from $U_{i+1} \cup \dots \cup U_d$ to U'_i . For every clique Q from $U_{i+1} \cup \dots \cup U_d$ and every clique type β realized in U'_i , let $f(\beta)$ be the number of R_{i-1} -edges sent by Q to cliques of type β in U_i , if this number is not greater than M_φ , or, otherwise, let $f(\beta) = M_\varphi$. Let $f'(\beta)$ be the number of R_{i-1} -edges sent by Q to cliques of type β in U'_i (recall that this number is not greater than M_φ). We let all the R_{i-1} -edges sent by Q to the cliques of type β in U'_i to be also members of R_i , that is, to be edges in \mathfrak{A}_i . Additionally, we link Q by R_i to the first (with respect to the numbering we have fixed) $f(\beta) - f'(\beta)$ cliques of type β in U'_i to which Q is not linked by R_{i-1} . By the choice of U'_i , we have enough such cliques in U'_i . We finish the construction of \mathfrak{A}_i by removing all the cliques from $U_i \setminus U'_i$.

That \mathfrak{A}_i has the desired properties is shown in the following two claims.

Claim 1: Each of the \mathfrak{A}_i is a transitive structure.

We show this by induction by $i = -1, 0, \dots, d$. Obviously $\mathfrak{A}_{-1} = \mathfrak{A}$ is transitive. Assume that \mathfrak{A}_{i-1} is transitive, and assume to the contrary that \mathfrak{A}_i is not. This means there are cliques Q_1, Q_2, Q_3 in \mathfrak{A}_i such that $\langle Q_1, Q_2 \rangle \in R_i$, $\langle Q_2, Q_3 \rangle \in R_i$ but $\langle Q_1, Q_3 \rangle \notin R_i$. It is easy to see that the cliques Q_1, Q_2, Q_3 must belong to three different layers, and that precisely one of the two cases holds: either $Q_2 \subseteq U'_i$, $\langle Q_1, Q_2 \rangle \notin R_{i-1}$, $\langle Q_2, Q_3 \rangle \in R_{i-1}$ or $Q_3 \subseteq U'_i$, $\langle Q_1, Q_2 \rangle \in R_{i-1}$, $\langle Q_2, Q_3 \rangle \notin R_{i-1}$. In the first case, our construction implies that there is a clique $Q' \subseteq U_i \setminus U'_i$ such that $\langle Q_1, Q' \rangle \in R_{i-1}$, and the clique types of Q_2 and Q' are identical. But from the latter it follows that $\langle Q', Q_3 \rangle \in R_{i-1}$ and from transitivity of R_{i-1} we have $\langle Q_1, Q_3 \rangle \in R_{i-1}$. Since none of Q_1, Q_3 is contained in U_i , by our construction we have that $\langle Q_1, Q_3 \rangle \in R_i$. Contradiction. In the second case, let β be the clique type of Q_3 and let Q'_1, \dots, Q'_{k_1} be the cliques of type β from U'_i to which Q_2 sends R_{i-1} -edges, Q''_1, \dots, Q''_{k_2} be the cliques of type β from $U_i \setminus U'_i$ to which Q_2 sends R_{i-1} -edges, and let $Q'''_1, \dots, Q'''_{k_3}$ be the cliques of type β from U'_i to which Q_1 sends R_{i-1} -edges, but Q_2 does not. Note that, by transitivity of R_{i-1} , Q_1 sends R_{i-1} -edges to all of the Q'_i and all of the Q''_i . If $k_1 + k_2 + k_3 \geq M_\varphi$, then Q_1 must send, by our construction, an R_i -edge to every clique of type β from U'_i , in particular to Q_3 ; contradiction. Thus, $k_1 + k_2 + k_3 < M_\varphi$ and Q_1 sends at least k_2 R_i -edges to cliques of type β from U'_i to which it does not send R_{i-1} -edges. Q_2 sends precisely k_2 such edges. Thus, since our strategy of choosing always cliques of type β with minimal possible numbers in the numbering we have fixed requires Q_2 to send an R_i -edge to Q_3 , the same strategy requires Q_1 also to send an R_i -edge to Q_3 . Contradiction.

Claim 2: The M_φ -profile of every clique in \mathfrak{A}_i is the same as its M_φ -profile in \mathfrak{A} . Again we work by induction. Assume that the M_φ -profiles of the surviving cliques in \mathfrak{A}_{i-1} are the same as in \mathfrak{A} . We show that the M_φ -profiles of cliques surviving in \mathfrak{A}_i are the same as in \mathfrak{A}_{i-1} . It is obvious for the \mathcal{H} -components and the values of *irref*, as we do not change the cliques. The \mathcal{A} -components for the cliques from $U'_0 \dots U'_i$ cannot change since they send R_i -edges to precisely the same cliques they send R_{i-1} -edges. Similarly, the \mathcal{B} -components for the cliques from $U_{i+1} \dots U_d$ cannot change since they receive R_i -edges precisely from the same cliques they receive R_{i-1} -edges.

Consider a clique Q from $U'_0 \dots U'_{i-1}$. Note that $prof_{\mathfrak{A}_i}^{M_\varphi}(Q) \cdot \mathcal{B} \subseteq prof_{\mathfrak{A}_{i-1}}^{M_\varphi}(Q) \cdot \mathcal{B}$ since any R_i -edge received by Q is also an R_{i-1} -edge. To see that \supseteq also holds take any

1-type $\alpha \in \text{prof}_{\mathfrak{A}_{i-1}}^{M_\varphi}(Q).\mathcal{B}$. Then there exists a clique Q' containing a realization of α such that Q' sends an R_{i-1} -edge to Q . If Q' survives in \mathfrak{A}_i then it sends an R_i -edge to Q . Otherwise $Q' \setminus U_i \setminus U'_i$ and there is a clique Q'' of the same clique type as Q' in U'_i . This equality of the clique types implies that α is realized in Q'' and Q'' sends an R_i -edge to Q . It follows that $\alpha \in \text{prof}_{\mathfrak{A}_i}^{M_\varphi}(Q)$. Thus, $\text{prof}_{\mathfrak{A}_{i-1}}^{M_\varphi}(Q) = \text{prof}_{\mathfrak{A}_i}^{M_\varphi}(Q)$.

Consider a clique Q from U'_i . Obviously $\text{prof}_{\mathfrak{A}_{i-1}}^{M_\varphi}(Q).\mathcal{B} \subseteq \text{prof}_{\mathfrak{A}_i}^{M_\varphi}(Q).\mathcal{B}$ since all the R_{i-1} -edges received by Q remain R_i -edges. To see \supseteq assume $\alpha \in \text{prof}_{\mathfrak{A}_i}^{M_\varphi}(Q).\mathcal{B}$ for some 1-type α . Then there exists a clique Q' containing a realization of α such that Q' sends an R_i -edge to Q . If Q' sends also an R_{i-1} -edge to Q , then $\alpha \in \text{prof}_{\mathfrak{A}_{i-1}}^{M_\varphi}(Q).\mathcal{B}$. Otherwise, by our construction, Q' sends an R_{i-1} -edge to a clique $Q'' \subseteq U_i \setminus U'_i$ such that the clique types of Q and Q'' are equal. But then α belongs to the \mathcal{B} -component of the clique type of Q'' and also of Q . So, $\alpha \in \text{prof}_{\mathfrak{A}_{i-1}}^{M_\varphi}(Q).\mathcal{B}$. It follows that $\text{prof}_{\mathfrak{A}_{i-1}}^{M_\varphi}(Q).\mathcal{B} = \text{prof}_{\mathfrak{A}_i}^{M_\varphi}(Q).\mathcal{B}$.

Finally, consider a clique Q from $U_{i+1} \cup \dots \cup U_d$. It remains to show that $\text{prof}_{\mathfrak{A}_{i-1}}^{M_\varphi}(Q).\mathcal{A} = \text{prof}_{\mathfrak{A}_i}^{M_\varphi}(Q).\mathcal{A}$. By our construction, the R_i -edges sent by Q to $U'_0 \cup \dots \cup U'_{i-1} \cup U_{i+1} \cup \dots \cup U_d$ are the same as R_{i-1} -edges sent by Q to this set. The desired equality of the \mathcal{A} -components (as multisets) follows now easily from the fact that, for any clique-type β , whenever Q sends precisely k R_{i-1} -edges to cliques of U_i of type β then it sends precisely k' -edges to cliques of U'_i of type β , where $k' = \min(k, M_\varphi)$. This finishes the proof of Claim 2.

The two above claims and Lemma 10 imply that $\mathfrak{A}' = \mathfrak{A}_d$ is indeed a model of φ . As each of the U'_i contains a finite number of cliques and each of the cliques is finite, we get that \mathfrak{A}' is finite. This finishes the proof of Lemma 11. \square

Let us estimate the size of the constructed finite model \mathfrak{A}' . For U'_0 we take at most M_φ realizations of every clique type from U_0 . M_φ is bounded exponentially, and the number of possible clique types in U_0 is bounded doubly exponentially in $|\varphi|$ (note that such cliques do not send any edges). Then, to construct U'_i we consider clique types distinguished, in particular, by the sets of cliques from $U'_0 \cup \dots \cup U'_{i-1}$ to which a given clique sends edges. Thus, the number of cliques in U'_i may become exponentially larger than the number of cliques in U'_{i-1} . Thus, we can only estimate the number of cliques in our eventual finite model by a tower of exponents of height d (recall that our bound on d is exponential in $|\varphi|$, though a polynomial bound would not be difficult to obtain). We leave open the question if a construction building smaller (e.g., doubly exponential in $|\varphi|$) models exist.

A careful inspection shows that all our constructions respect reflexivity and seriality, that is, if we replace the word *transitive* in the statements of Lemmas 9 and 11 with the phrases *reflexive transitive* or *serial transitive* then they remain correct.

Theorem 8

The logics $K4(\diamond_{\geq}, \diamond)$, $D4(\diamond_{\geq}, \diamond)$, $S4(\diamond_{\geq}, \diamond)$ have the finite model property. Their local and global satisfiability problems are decidable.

A natural decision procedure arising from our work is as follows: guess a finite model of the given formula and check that it indeed is a model. However, this procedure does not give a good upper complexity bound, since it needs to take into account very large finite models. The precise complexity can be established using the above-mentioned results from Gogacz et al. (2019) concerning the DL \mathcal{SIQ}^- .

Theorem 9

The local and global satisfiability problems for the logics $K4(\diamond_{\geq}, \diamond)$, $D4(\diamond_{\geq}, \diamond)$, $S4(\diamond_{\geq}, \diamond)$ are 2-EXPTIME-complete.

Proof

In [Gogacz et al. \(2019\)](#), it is shown that the knowledge base satisfiability problem for the logic SIQ^- , restricted to a single transitive role, is 2-EXPTIME-complete. With this single role restriction, the language of SIQ^- becomes a syntactic variant of $K4(\diamond_{\geq}, \diamond)$. The knowledge base satisfiability in SIQ^- is the question if for a given pair $(\mathcal{T}, \mathcal{A})$, where \mathcal{T} is a *TBox* and \mathcal{A} is an *ABox*, there exists a structure containing \mathcal{A} and respecting \mathcal{T} at every element. We do not want to define these notions formally here and refer the interested reader to [Gogacz et al. \(2019\)](#) or some other articles on DLs. For our purposes, it is sufficient to say that \mathcal{T} consists of implications of the form $\phi \rightarrow \psi$ and \mathcal{A} is a collection of assertions of the form $\phi(a)$ or $T(a, b)$ where a, b are names for domain elements (which can be used only in \mathcal{A}), $\phi(a)$ means that ϕ is satisfied at a , and $T(a, b)$ means that there is an edge from a to b .

To solve global satisfiability for $K4(\diamond_{\geq}, \diamond)$, we just translate the input formula ϕ to the knowledge base $(\{\top \rightarrow \phi\}, \emptyset)$ and ask for its satisfiability. Regarding the lower bound, we can easily adapt the lower bound proof from [Gogacz et al. \(2019\)](#) (Theorem 4) to our scenario. The proof there goes by a reduction from the acceptance problem for alternating Turing machines with exponentially bounded space and uses both TBoxes and ABoxes. However, ABoxes are always of a simple form $\phi'(a)$. What we can do is to take the conjunction ϕ of the $K4$ -counterparts of the implications from the given TBox and ask for combined satisfiability of ϕ and ϕ' . This gives the 2-EXPTIME-lower bound for the combined complexity of $K4(\diamond_{\geq}, \diamond)$. Due to Lemma 7 we infer 2-EXPTIME-completeness of local and global satisfiability in $K4(\diamond_{\geq}, \diamond)$.

The upper and lower complexity bounds for $K4(\diamond_{\geq}, \diamond)$ and $S4(\diamond_{\geq}, \diamond)$ can be obtained by an inspection of the proofs from [Gogacz et al. \(2019\)](#) and observing that they work for structures with a reflexive or serial transitive relation. \square

5 Missing lower bounds for logics with converse and without graded modalities

To complete the picture, we consider in this section the modal language with converse but without graded modalities. Over most relevant classes of frames tight complexity bounds for local and global satisfiability of this language are known. However, according to Zolin's survey ([Zolin 2017](#)), the three logics of transitive frames $K4(\diamond, \diamond)$, $S4(\diamond, \diamond)$, and $D4(\diamond, \diamond)$ whose global satisfiability is known to be in EXPTIME lack the corresponding lower bound. We provide it here. We were also not able to find a tight lower bound in the literature for the logics of Euclidean frames, $K5(\diamond, \diamond)$, $D5(\diamond, \diamond)$. We also show it here. Interestingly, the two reductions are identical, that is, in both cases we produce the same formulas (but the shapes of the intended models differ).

In the conference version of this paper, we used a rather heavy reductions from the halting problem for alternating Turing machines working in polynomial space. Following the suggestion of one of the referees, we looked for an alternative proof by a reduction

from global satisfiability of the logic $K(\diamond)$. The general idea is essentially the same as in our previous proof, but the reduction is arguably simpler.

Theorem 10

The global satisfiability problem for $K4(\diamond, \boxplus)$, $D4(\diamond, \boxplus)$, and $S4(\diamond, \boxplus)$ is EXPTIME-hard.

Proof

We recall that global satisfiability problem for $K(\diamond)$ is EXPTIME-hard. We reduce this problem simultaneously to global satisfiability of the three logics we consider.

Take any modal formula φ of $K(\diamond)$. Without loss of generality, we assume that φ contains no nested occurrences of \diamond and \square . (Indeed, if φ contains a nested occurrence of a modal operator, that is, it contains a subformula $\diamond\psi$ or $\square\psi$ in the scope of another \diamond or \square , then we replace that subformula by a fresh variable p and append the conjunct $p \leftrightarrow \diamond\psi$, resp., $p \leftrightarrow \square\psi$. Successively treating in this way all occurrences of modal operators we eventually end up with a formula equisatisfiable to φ in which they are not nested.)

Assuming that c_0, c_1, c_2 , and c_3 are fresh propositional variables not occurring in φ we define the translation $\mathbf{tr}(\varphi)$ as follows:

- $\mathbf{tr}(p) = p$ for all propositional variables p ,
- $\mathbf{tr}(\varphi' \vee \varphi'') = \mathbf{tr}(\varphi') \vee \mathbf{tr}(\varphi'')$ and analogously for $\vee, \rightarrow, \leftrightarrow$,
- $\mathbf{tr}(\neg\varphi') = \neg\mathbf{tr}(\varphi')$,
- $\mathbf{tr}(\diamond\varphi') = [c_0 \rightarrow \diamond(c_1 \wedge \mathbf{tr}(\varphi'))] \wedge [c_1 \rightarrow \boxplus(c_2 \wedge \mathbf{tr}(\varphi'))] \wedge [c_2 \rightarrow \diamond(c_3 \wedge \mathbf{tr}(\varphi'))] \wedge [c_3 \rightarrow \boxplus(c_0 \wedge \mathbf{tr}(\varphi'))]$ and analogously for $\square\varphi'$

Let $\varphi^* = \mathbf{tr}(\varphi) \wedge (\bigvee_{0 \leq i \leq 3} c_i) \wedge (\bigwedge_{0 \leq i < j \leq 3} (\neg c_i \vee \neg c_j))$. Note that φ^* is composed of the translated φ and a formula stipulating that for each node exactly one of c_i holds true. The size of φ^* is clearly polynomial in $|\varphi|$ since \diamond and \square have no nested occurrences in φ .

Claim 1

If φ is globally satisfiable in $K(\diamond)$, then φ^* is globally satisfiable in $K4(\diamond, \boxplus)$, $D4(\diamond, \boxplus)$, and $S4(\diamond, \boxplus)$.

Proof

Let $\mathfrak{A} = \langle W, R, V \rangle$ be a model of φ . We assume that \mathfrak{A} is tree-shaped (this is done without loss of generality since $K(\diamond)$ has the tree-shaped model property). Let w_r denote the root of \mathfrak{A} .

For any world w define its distance from the root, denoted with $d(w)$, as the length of the R -path from w_r , that is, $d(w_r) = 0$, $d(w) = 1$ iff $R(w_r, w)$ holds, $d(w) = 2$ iff there is a world v such that $R(w_r, v), R(v, w)$, etc. We define the Kripke structure $\mathfrak{A}' = \langle W', R', V' \rangle$ by inverting every second R -edge of \mathfrak{A} and labelling the worlds on every path, leading from the root, repetitively $c_0, c_1, c_2, c_3, c_0, \dots$. Formally:

- $W = W'$,
- For every propositional variable $p \notin \{c_0, c_1, c_2, c_3\}$, we set $V'(p) = V(p)$ while for the variables c_i we set $V'(c_i) = \{w : d(w) \bmod 4 = i\}$ for $i \in \{0, 1, 2, 3\}$,
- R' is the reflexive closure of $R_{(0,1)} \cup R_{(1,2)}^{-1} \cup R_{(2,3)} \cup R_{(3,0)}^{-1}$, with $R_{(i,j)} = R \cap V'(c_i) \times V'(c_j)$.




Fig. 4. Shape of intended models in the proof of Theorem 10. All worlds are reflexive.

The shape of the obtained model is illustrated in Figure 4.

Now we show that φ^* is globally satisfiable in $K4(\diamond, \diamond)$, $D4(\diamond, \diamond)$, and $S4(\diamond, \diamond)$. First, note that due to the construction R' is transitive and reflexive (and thus also serial). Next, note that the second part of the formula φ^* is globally satisfied in \mathfrak{A}' , since every world belongs to exactly one of $V'(c_i)$ (due to the fact that the satisfaction of c_i depends on a distance from the root, which is unique since \mathfrak{A} is assumed to be tree-shaped). Finally, we show that $\mathfrak{A}' \models \mathbf{tr}(\varphi)$. The proof is by induction, where the inductive hypothesis states that for any subformula ψ of φ and every world w we have $\mathfrak{A}, w \models \psi$ if and only if $\mathfrak{A}', w \models \mathbf{tr}(\psi)$. The case of ψ being a propositional variable follows from the second item of definition of \mathfrak{A}' . The case when ψ is a Boolean combination of formulas is immediate from the inductive hypothesis and the semantics of \models . Hence, the only interesting case is when ψ is of the form $\diamond(\psi')$. We prove only one implication; the second one is analogous. Assume that $\mathfrak{A}, w \models \diamond(\psi')$. Thus, there is a world v such that $R(w, v)$ and $\mathfrak{A}, v \models \psi'$. By induction hypothesis, we deduce that $\mathfrak{A}', v \models \mathbf{tr}(\psi')$. Moreover, for $i = d(w) \bmod 4$ and $j = d(v) \bmod 4 = (i + 1) \bmod 4$, we have $\mathfrak{A}', w \models c_i$ and $\mathfrak{A}', v \models c_j$. Moreover, if i is even, then $(w, v) \in R'$ and $(v, w) \in R'$ otherwise. In each of the cases $i \in \{0, 1, 2, 3\}$ these all imply that $\mathfrak{A}', w \models \mathbf{tr}(\psi)$, which finishes the proof. \square

Claim 2

If φ^* is globally satisfiable in $K4(\diamond, \diamond)$, $D4(\diamond, \diamond)$, or $S4(\diamond, \diamond)$ then φ is globally satisfiable in $K(\diamond)$.

Proof

Let $\mathfrak{A} = \langle W, R, V \rangle$ be a model of φ^* . We will define an increasing chain of structures $\mathfrak{A}'_0, \mathfrak{A}'_1, \dots$, in which $\mathfrak{A}'_i = \langle W'_i, R'_i, V'_i \rangle$, together with a pattern function $f : A_0 \cup A_1 \cup \dots \rightarrow W$. The Kripke structure $\mathfrak{A}' = \langle W', R', V' \rangle$ defined as the union of the chain will turn out to be a model of φ . Our chain of structures is defined as follows.

We fix a world $w \in W$, set $\mathfrak{A}'_0 = \langle \{w'\}, \emptyset, V'_0 \rangle$ with $V'_0(w') = V(w)$ and set $f(w') = w$. For simplicity, let us assume that $\mathfrak{A}, w \models c_0$. In our construction, for every element w' freshly added to \mathfrak{A}'_i , we will have that $f(w')$ satisfy $c_{i \bmod 4}$.

Assume now that \mathfrak{A}'_i is defined. To construct \mathfrak{A}'_{i+1} we repeat for every element w' freshly added to \mathfrak{A}'_i : if i is even (odd) then for every R -successor (R -predecessor) v of $f(w')$ in \mathfrak{A} such that $\mathfrak{A}, v \models c_{i+1 \bmod 4}$ add to W'_{i+1} a fresh R -successor v' of w' and let $V'_i(v') = V(v)$ and $f(v') = v$.

We prove inductively over the shape of ψ that $\mathfrak{A}', w' \models \psi$ iff $\mathfrak{A}, f(w') \models \mathbf{tr}(\psi)$. The case of atomic propositions and Boolean combinations follows immediately from the definition. The only interesting case is of $\psi = \diamond(\psi')$. Here we show only one case of one implication; the other cases are analogous. Assume that $\mathfrak{A}, f(w') \models \mathbf{tr}(\psi)$ holds as well as $f(w') \models c_0$. Then there is an R -successor v of $f(w')$ satisfying $\mathbf{tr}(\psi') \wedge c_1$. Note that the R' -successors of w' are copies of R -successors of $f(w')$ satisfying c_1 ; thus, there is a world v' being an R' -successor of $f(w')$ and satisfying $f(v') = v$. Hence, from the inductive assumption, we infer $\mathfrak{A}', v' \models \psi'$, which implies $\mathfrak{A}', w' \models \psi$. Analyzing analogously the other cases we finish the inductive proof of the claim. \square

The two preceding claims show the correctness of the translation, allowing us to conclude Theorem 10. \square

We next handle the case of Euclidean frames.

Theorem 11

The global and local satisfiability problem for $K5(\diamond, \diamond)$ and $D5(\diamond, \diamond)$ is EXPTIME-hard.

Proof

We explicitly consider the global satisfiability problem, but due to Lemma 3 our proof applies also to local satisfiability. The proof goes as the proof of Theorem 10. Our current intended models are similar to the intended models there (as on Figure 4). The difference is that all the worlds satisfying c_1 or c_3 are made equivalent to each other, and the worlds satisfying c_0 or c_2 are irreflexive. Note that this does not violate the property that each world can identify its children in the tree. Observe also that such intended models are indeed Euclidean and serial (however, they are neither transitive nor reflexive); in particular all worlds satisfying c_0 or c_2 are lanterns. Now, for a given K formula we can construct precisely the same formula as in the previous proof. We leave the routine details to the reader. The correctness proof is essentially identical to the correctness proof of Theorem 10 so we omit it here. \square

6 Conclusions

We have filled the gaps remaining in the classification of the complexity of the local and global satisfiability problems for natural modal languages with graded and converse modalities over traditional classes of frames. What we have not systematically studied are the problem of combined satisfiability (given two formulas check if there exists a model in which the first is satisfied locally and the second is satisfied globally) and the problem of finite (local, global, combined) satisfiability (asking about the existence of *finite* models). We suspect that the classification could be extended to cover these problems using results/techniques from our paper and the referenced articles without major obstacles.

Two other questions we leave open are if the NEXPTIME-lower bound in Theorem 5 remains valid, if the numbers in graded modalities are encoded in unary rather than in binary, and if our finite model construction from Section 4 can be replaced by a one producing smaller models.

Conflicts of interest

The author(s) declare none.

References

- BAADER, F., HORROCKS, I., LUTZ, C. AND SATTTLER, U. 2017. *An Introduction to Description Logic*. Cambridge University Press.
- BEDNARCZYK, B., KIERONSKI, E. AND WITKOWSKI, P. 2019. On the complexity of graded modal logics with converse. In *Logics in Artificial Intelligence - 16th European Conference, JELIA 2019, Rende, Italy, May 7–11, 2019, Proceedings*, F. Calimeri, N. Leone and M. Manna, Eds. Lecture Notes in Computer Science, vol. 11468. Springer, 642–658.
- BLACKBURN, P., DE RIJKE, M. AND VENEMA, Y. 2001. *Modal Logic*. Cambridge University Press, New York, NY, USA.
- BLACKBURN, P. AND VAN BENTHEM, J. 2007. Modal logic: A semantic perspective. In *Handbook of Modal Logic*, P. Blackburn, J. F. A. K. van Benthem and F. Wolter, Eds. Studies in Logic and Practical Reasoning, vol. 3. North-Holland, 1–84.
- CHAGROV, A. V. AND RYBAKOV, M. N. 2002. How many variables does one need to prove PSPACE-hardness of modal logics. In *Advances in Modal Logic 4, Papers from the Fourth Conference on “Advances in Modal Logic,” Held in Toulouse, France, 30 September–2 October 2002*, P. Balbiani, N. Suzuki, F. Wolter and M. Zakharyashev, Eds. King’s College Publications, 71–82.
- CHEN, C. AND LIN, I. 1994. The complexity of propositional modal theories and the complexity of consistency of propositional modal theories. In *Logical Foundations of Computer Science, Third International Symposium, LFCS’94, St. Petersburg, Russia, July 11–14, 1994, Proceedings*, 69–80.
- COOK, S. A. 1971. The complexity of theorem-proving procedures. In *Proceedings of the 3rd Annual ACM Symposium on Theory of Computing, May 3–5, 1971, Shaker Heights, Ohio, USA*, M. A. Harrison, R. B. Banerji and J. D. Ullman, Eds. ACM, 151–158.
- DEMRI, S. AND DE NIVELLE, H. 2005. Deciding regular grammar logics with converse through first-order logic. *Journal of Logic, Language and Information* 14, 3, 289–329.
- GOGACZ, T., GUTIÉRREZ-BASULTO, V., IBÁÑEZ-GARCÍA, Y., JUNG, J. C. AND MURLAK, F. 2019. On finite and unrestricted query entailment beyond SQ with number restrictions on transitive roles. In *Proceedings of the Twenty-Eighth International Joint Conference on Artificial Intelligence, IJCAI 2019, Macao, China, August 10–16, 2019*. ijcai.org, 1719–1725.
- GUTIÉRREZ-BASULTO, V., IBÁÑEZ-GARCÍA, Y. A. AND JUNG, J. C. 2017. Number restrictions on transitive roles in description logics with nominals. In *Proceedings of the Thirty-First AAAI Conference on Artificial Intelligence, February 4–9, 2017, San Francisco, California, USA*, 1121–1127.
- KAZAKOV, Y. AND PRATT-HARTMANN, I. 2009. A note on the complexity of the satisfiability problem for graded modal logics. In *Proceedings of the 24th Annual IEEE Symposium on Logic in Computer Science, LICS 2009, 11–14 August 2009, Los Angeles, CA, USA*, 407–416.
- KAZAKOV, Y., SATTTLER, U. AND ZOLIN, E. 2007. How many legs do I have? non-simple roles in number restrictions revisited. In *Logic for Programming, Artificial Intelligence, and Reasoning, 14th International Conference, LPAR 2007, Yerevan, Armenia, October 15–19, 2007, Proceedings*, 303–317.
- LADNER, R. E. 1977. The computational complexity of provability in systems of modal propositional logic. *SIAM Journal on Computing* 6, 3, 467–480.
- LUTZ, C. 2002. *The Complexity of Reasoning with Concrete Domains*. Ph.D. thesis, LuFG Theoretical Computer Science, RWTH-Aachen, Germany.

- PRATT-HARTMANN, I. 2005. Complexity of the two-variable fragment with counting quantifiers. *Journal of Logic, Language and Information* 14, 3, 369–395.
- PRATT-HARTMANN, I. 2007. Complexity of the guarded two-variable fragment with counting quantifiers. *Journal of Logic and Computation* 17, 1, 133–155.
- PRATT-HARTMANN, I. 2008. On the computational complexity of the numerically definite syllogistic and related logics. *Bulletin of Symbolic Logic* 14, 1, 1–28.
- TOBIES, S. 2001a. *Complexity Results and Practical Algorithms for Logics in Knowledge Representation*. Ph.D. thesis, RWTH-Aachen, Germany.
- TOBIES, S. 2001b. PSPACE reasoning for graded modal logics. *Journal of Logic and Computation* 11, 1, 85–106.
- ZOLIN, E. 2017. Undecidability of the transitive graded modal logic with converse. *Journal of Logic and Computation* 27, 5, 1399–1420.