Vol. LXXXIII No. 2

February 1969

The Journal of Laryngology and Otology

EDITED BY G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

Contents

Foetal Stapes—I. General Features	RAMJI DASS, S. P. THAPAR and SEWA SINGH MAKHNI	IOI
The Treatment of Carcinoma of the Hypopharynx: the Gutter-Graft Operation	RONALD MACBETH	101
Actiopathogenesis of Scleroma	A. Sinha, S. C. Pandhi and Om Prakash	133
Clinical and Medico-Legal Aspects of Acute Epi- glottiditis	A. POTONDI and O. RIBÁRI	
Sodium and Potassium Concentrations in Post- mortem Human Labyrinthine Fluids.	T. Palva	141
Virus Studies in Secretory Otitis Media	and P. TIKANMÄKI PETER ADLINGTON and JOAN R. DAVIES	147 161
Deafness and the Klippel-Feil Syndrome Clinical Records—	K. McLay and A. G. D. MARAN	175
Saucerization of the Radical Mastoidectomy Cavity	T. S. Stewart	185
Neurilemmoma of the Larynx	ALUN B. THOMAS and LYN REES	189
A Case of Respiratory Obstruction from Birth Due to a Polyp of Nasal Origin	C. R. SALKELD	197
Neurilemmoma of the Tongue	O, Ö. BITITCI	201

General Notes

Headley Brothers Ltd Ashford, Kent

Annual Subscription £5/5/0 net, U.S.A. \$15 https://doi.org/10.1017/50022215100070110 Published online by Cambridge University Press Monthly, 12/6 net, post free

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the understanding that they are contributed to this *Journal* solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgment must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc. The Harvard system of recording references should be used, e.g. GREEN, C.,

The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in World Medical Periodicals, published by World Health Organization, 1952.

It is most important that authors should verify *personally* the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks will normally be held by the Printers for three years, after which they will be destroyed. Any author is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS LTD., The Invicta Press, Ashford, Kent. Coloured illustrations will be charged to authors, unless a special grant is authorized by the Editor.

5. Orders for reprints must be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Editorial communications may be addressed to THE EDITOR, Journal of Laryngology, c/o HEADLEY BROTHERS LTD., The Invicta Press, Ashford, Kent.

7. The annual subscription is five guineas sterling (U.S.A. \$15) post free, and payable in advance.

8. Single copies will be on sale at 128. 6d. each; copies of parts up to Vol. LXIII which are available may be purchased at 78. 6d. each.


9. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS LTD., The Invicta Press, Ashford, Kent.

United States of America

Orders for this Journal may be sent through local booksellers, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, HEADLEY BROTHERS LTD., The Invicta Press, Ashford, Kent, England.

C Journal of Laryngology and Otology, 1969

i



Tegretol[®] Anticonvulsant

Widens the scope of anti-epileptic therapy

Geigy


Tegretol is available as tablets each containing 200 mg 5-carbamoyl-5H-dibenz (b,f) azepine.

Detailed literature is available on request

Geigy (U.K.) Limited Pharmaceuticals Division Macclesfield, Cheshire

PETERS AP/5 CLINIC AUDIOMETER

WITH SYNCHRONOUS MASKING



Please mention The Journal of Laryngology and Otology when replying to advertisements

Refer your patients to Ingrams Independent HEARING AID SUPPLIERS

Ingrams have the different makes to choose from

LONDON	2 Shepherd Street, London, W.1. (2 minutes from Green Park)	Hyde Park 9041	
BIRMINGHAM	20 Auchinleck Sq, Broad St, Birmingham, 15. (In the Five Ways Shopping Centre)	Midland 4314	
BRIGHTON	5 Duke Street, Brighton, 1, (First turning off West Street)	Brighton 28699	
CROYDON	11 Essex House, College Road, Croydon. (Opposite the Technical College)	Croydon 4663	
MANCHESTER	Mansfield Chambers (Rooms 21 & 22), 17 St. Anne's Square, Manchester, 2. (Off Deansgate)	B'ackfriars 4104 (STD Code 061-834 4104)	
and in SCOTLAND it's BONOCHORD-INGRAMS			
ABERDEEN	461 Union Street, Aberdeen. (Near Holborn Junction)	Aberdeen 29632	
EDINBURGH	17 Stafford Street, Edinburgh. (Corner of Alva Street)	Caledonian 1942	
GLASGOW	38 Bath Street, Glasgow, C.2. (Next door to Glasgow Corporation Transport Office)	Douglas 7710	
Domiciliary visits in EVERY part of the country. INGRAMS FOR THE <u>LARGER</u> RANGE OF HEARING AIDS— Subscribers to the Code of Ethics of the Hearing Aid Industry Association.			

Specifically designed for you and your patients. Precision-engineered, functionally-built equipment for the Ear, Nose and Throat and other Specialists.

S.M.R. keeps you and your patients' comfort and satisfaction in mind. Whatever your needs . . . whether they be for one unit or a completely equipped treatment room, you are assured of the finest . . . a room that reflects professional good taste.



Picture shows what can be done with small treatment room areas. The rooms are complete from the compact treatment unit to the physician's stool. Yet, this equipment fits easily and functionally into a small area. This is just one of many room arrangements available from S.M.R.



Please mention The Journal of Laryngology and Otology when replying to advertisements

<section-header><section-header><section-header><section-header><section-header>

Clinicians have frequently testified to the success of Orbenin in the treatment of Gram-positive infections. World Literature now contains over 400 references to Orbenin.

Pneumonia 🗌 Lung Abscess 🗌 Tonsillitis 🗌 Pharyngitis 🗋 Acute bronchitis

Septicaemia

Boils Carbuncles Infected dermatoses

Osteomyelitis
Osteitis

Post-operative wound infections 🗌 Burns 📋 Skin graft protection

Acute endocarditis

Staphylococcal enterocolitis []] Staphylococcal urinary tract infections Staphylococcal meningitis

Dosage

Adults: 500mg (2 capsules) four times daily, by mouth. 250mg (1 vial) by intramuscular injection, four times daily.

Children: Under 2 years—quarter adult dose. 2-10 years—half adult dose.

In severe infections the dosage may be increased.

Contra-indications

Orbenin should not be given to patients with a penicillin allergy or administered by subconjunctival injection.

Side-effects

As with other penicillins. Additional information is available on

request.

Orbenin (Regd.) (cloxacillin sodium BP) is a product of British research at **Beecham Research Laboratories** Brentford, England. originators of the new penicillins.





THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT

Official organ for the American Laryngological Rhinological and Otological Society

Price \$18.00 per year Canada \$19.00 per year Foreign \$19.00 per year

ESTABLISHED 1896

THEODORE E. WALSH, M.D. EDITOR

517 SOUTH EUCLID AVENUE SAINT LOUIS MO. 63110.



P. C. WERTH LTD. Contractors to H.M. Government and Local Authorities

Specialist Supply, Maintenance, Calibration Services for Audiological Test and Teaching Equipment

17 STRATFORD PLACE, OXFORD ST., LONDON W1N ODH Telephone: 01-629 5559

Clinical, Diagnostic, Screening and Bekesy Audiometers. Acoustic Impedance Apparatus. Stereo Auditory Trainers. Visible Speech Apparatus. Artificial Mastoid, etc.

Very Advanced Specification incorporates: Filtered White Noise or Narrow Band Masking. Extended Hearing Loss Range (Up to 120 dB). Full facilities for Speech Audiometry (Live or Recorded). Fully Transistorised. Battery or Mains Powered. Available with Noise Insulated Headset. Supplied complete with Smart Practical Carrying Case. All Equipment Fully Guar-

anteed and Serviced/Calibrated on our own Premises. Kamplex Portable Diagnostic Audiometer Model TA 151



Please mention The Journal of Laryngology and Olology when replying to advertisements



'Otosporin' rapidly destroys virtually all bacteria commonly found in ear infections, reduces irritation, inflammation and pain. 'Otosporin' brand Drops contain polymyxin B sulphate, neomycin sulphate and hydrocortisone in a bland suspension designed to facilitate penetration. Issued in bottles of 5 ml.



Full information is available on request.



Burroughs Wellcome & Co. London (The Wellcome Foundation Ltd.)

eliminate the cross infection risk



Opulets sterile, single dose eye and ear preparations

To overcome the risk of contamination in eye and ear infections, Opulets have been developed as sterile, single dose eye and ear drop tubes and ointment capsules. Each is packed in a film sachet and sterilized by gamma radiation, ensuring effective sterilization with no loss of activity – the strict criteria of efficacy and safety being met in full.

After each application, these single dose units are discarded, eliminating cross infection risks – each treatment administered by either medical staff or the patient is thus under the most sterile conditions possible in practical medicine.

Opulets provide the lowest unit price available compared with all other preparations of a similar nature. Presentations available at present are: **Atropine Eye Ointment** Atropine Sulphate BP 0.5% w/w. Chloramphenicol Eye and Ear Drops Chloramphenicol BP 0.5% w/v. Chloramphenicol Eye and Ear Ointment Chloramphenicol BP 1.0% w/w. Fluorescein Eye Drops Fluorescein Sodium BP 1% w/v Hydrocortisone/Neomycin Eye and Ear Ointment Hydrocortisone Acetate BP 1.5% w/w Noemycin BP 0.5% w/w. **Physostigmine Eye Drops** Physostigmine (Eserine) Sulphate BPC 0.5% w/v. Sulphacetamide Eye Drops Sulphacetamide Sodium BP 10% w/v. Sulphacetamide Eye Ointment Sulphacetamide Sodium BP 6% w/w.

full descriptive technical literature is available on request Pharmax Limited Thames Road Crayford Kent

