

physiological response to these patients' abnormal eating habits or their low body weight.

The data on *treatment* are also highly equivocal. Whilst several reports have described the benefits of antidepressant medication in the treatment of patients with anorexia nervosa, no study has even met the minimal requirements of proper evaluation (Szmukler, 1982). There have also been claims that antidepressant medication helps patients with bulimia (DSM-III). However, the only controlled outcome study found that adequate dosages of mianserin had no specific effect on eating habits or attitudes, or indeed on mental state (Sabine *et al.*, 1983). It must be added that it is questionable whether response to antidepressant medication is a legitimate basis for assigning patients to a particular diagnostic category (Murray and Murphy, 1978).

With regard to *family history* of affective disorder, the study of Hudson and colleagues certainly substantiates earlier findings of a high prevalence of affective disorder amongst the relatives of patients with eating disorders. However, the only conclusion that can be drawn from this observation is that a family history of affective disorder predisposes individuals to develop an eating disorder.

In conclusion, an association between eating disorders and disturbances of mood is incontrovertible. However, the balance of evidence weighs against the suggestion that eating disorders are forms of affective disorder.

PETER J. COOPER  
CHRISTOPHER G. FAIRBURN

Warneford Hospital,  
Oxford OX3 7JX

#### References

- ECKERT, E. D., GOLDBERG, S. C., HALMI, K. A., CASPER, R. C. & DAVIS, J. M. (1982) Depression in anorexia nervosa. *Psychological Medicine*, **12**, 115–22.
- FAIRBURN, C. G. (1982) Binge eating and its management. *British Journal of Psychiatry*, **141**, 631–3.
- GERNER, R. H. & GWIRTSMAN H. E. (1981) Abnormalities of dexamethasone suppression test and urinary MHPG in anorexia nervosa. *American Journal of Psychiatry*, **138**, 650–3.
- Hsu, L. K. G. (1980) Outcome of anorexia nervosa: a review of the literature. *Archives of General Psychiatry*, **37**, 1014–6.
- HUDSON, J. I., LAFFER, P. S. & POPE, H. G. (1982) Bulimia related to affective disorder by family history and response to dexamethasone suppression test. *American Journal of Psychiatry*, **139**, 685–7.
- MURRAY, R. M. & MURPHY, D. L. (1978) Drug response and psychiatric nosology. *Psychological Medicine*, **8**, 667–81.
- RUSSELL, G. F. M. (1970) Anorexia nervosa: its identity as an illness and its treatment. In *Modern Trends in Psychological Medicine*, Volume 2 (ed. J. H. Price), pp 131–64. London: Butterworth.

SABINE, E. J., YONACE, A., FARRINGTON, A. J., BARRATT, K. H. & WAKELING, A. (1983) Bulimia nervosa: a placebo controlled double-blind therapeutic trial of mianserin. *British Journal of Clinical Pharmacology*, **5**, 195S–202S.

SZMUKLER, G. I. (1982) Drug treatment and anorexic states. In *Drugs and Appetite* (ed. T. Silverstone), pp. 159–81. London: Academic Press.

#### PHONEME DISCRIMINATION IN SCHIZOPHRENIA

DEAR SIR,

I read with interest the paper by Drs Kugler and Caudrey (*Journal*, January 1983, **142**, 53–59).

I have, however, found one point puzzling. It was reported in the article that all the patients met Feighner's diagnostic criteria of schizophrenia but, at the same time, the duration of their illness ranged from two months to 27 years. This seems to be contradictory since Feighner's criteria of schizophrenia requires the duration of at least six months.

T. KITAMURA

School of Medicine,  
Keio Gijuku University,  
Tokyo, Japan

#### Dr Kugler replies:

All patients in this study were initially tested on the basis of hospital diagnoses. Case-note summaries were made at the time of testing and were subsequently examined, using the syndrome checklist of the Present State Examination and Feighner's diagnostic criteria, to confirm diagnoses before the data were analysed. The mental state of the patients in the time between testing and data analysis was also referred to and the minimum duration of illness for any schizophrenic patient at the time of analysis was 18 months. Age, duration of illness, age at onset of the illness and medication data *at the time of testing* are reported in our paper.

I hope this answers Dr Kitamura's query and apologise for the lack of clarity on this point.

Harperbury Hospital, Herts

#### WEEKLY PIMOZIDE VERSUS FLUPHENAZINE DECANOATE IN SCHIZOPHRENIC OUT- AND DAY-PATIENTS

DEAR SIR,

Pimozide has been found to be effective as maintenance treatment in schizophrenics when given four days weekly to in- and day-patients (McCreadie *et al.*, 1980) and once weekly to in-patients (McCreadie *et al.*, 1982). A preliminary investigation has now been carried out in schizophrenics living in the community.

In a double-blind study, 27 male and female chronic