## NOTES FOR AUTHORS

The Bulletin of Entomological Research publishes original research papers concerning insects, mites, ticks or other arthropods of economic importance in agriculture, forestry, stored products, biological control, medicine, animal health and natural resource management. The geographical scope of the Bulletin is worldwide but with emphasis on the tropics. Taxonomic papers are accepted if relevant. Short review papers, although normally by invitation, will also be considered for publication.

**Page Format.** The *Bulletin* is printed in a two-column format (column width of three inches) with a text area of 170×240 mm.

**Text.** Papers should be typed, one one side of the paper only, with double line spacing and ample margins (at least 1.5 cm) on each side and with no underlining or bold in text except for scientific names. Draft quality print from a word-processor is not acceptable. Standard abbreviations (e.g. fig. and figs) and metric units must be used. Guidelines for taxonomic papers are available.

On acceptance, word-processed text stored on floppy disk is encouraged, providing the software is IBM/DOS compatible, but floppy discs must be accompanied by a hard copy. This will enable papers to be handled rapidly, and with less type-setting errors. Further instructions regarding coding of word-processor documents are available on request.

**Abstract.** Each paper must commence with a carefully prepared, accurate, informative abstract, in one paragraph, that is complete in itself and intelligible without reference to text or figures. It should not exceed 250 words.

**Tables.** Tables should be reduced to the simplest form, and should not be used where text or illustrations give the same information. They should be submitted on separate sheets at the end of the article and must fit conveniently into single column, full width or landscape (if absolutely necessary) format.

**Illustrations.** Text figures, line drawings and graphs should be of sufficient size and quality to allow for reduction by half or two-thirds. Half-tone photographs are acceptable where they are a real contribution to the text. Figure and Table captions should be typed on a <u>separate sheet</u> in the following format:

Figs 23–26. Figs 23–24, <u>Urophora</u> eggs: 23, <u>U. hispanica</u>; 24, <u>U. stigma</u>. Figs 25–26, spermathecae: 25, <u>U. maura</u>; 26, <u>U. stigma</u>; scale lines = 0.05 mm.

**References.** References must be based on the name and year system, give full journal titles and conform to the following styles:

Powell, W. (1986) Enhancing parasitoid activity in crops. pp. 319–340 in Waage, J. & Greathead, D. (Eds) <u>Insect parasitoids</u>. London, Academic Press (Symposium, Royal Entomological Society of London No. 13).

Southwood, T.R.E. (1978) Ecological methods with particular reference to the study of insect populations. 2nd edn. 524 pp. London, Chapman & Hall.

Zhou, X., Carter, N. & Mumford, J. (1989) A simulation model describing the population dynamics and damage potential of the rose grain aphid, <u>Metopolophium dirhodum</u> (Walker) (Hemiptera: Aphididae), in the UK. <u>Bulletin of Entomological Research</u> 79, 373–380.

Citation of authors in the text should appear in the form "Wilson (1986)" or "(Wilson, 1986)". More than one author should be cited as "(Holloway et al., 1987; Walker & Huddleston, 1988)".

**Offprints.** 50 copies of each paper are provided free to the author (or major author) of each paper. Further copies may be obtained on payment, and the number required should be specified and ordered at proof stage.

Manuscripts. The original manuscript and artwork plus two copies should be submitted to:

The Editors Bulletin of Entomological Research CAB International Institute of Entomology 56, Queen's Gate London SW7 5JR, United Kingdom.

## SEPTEMBER 1990

## **Bulletin of Entomological Research**

Guest Editorial	241
Bisset, J. A., Rodriguez, M. M., Diaz, C., Ortiz, E. & Marquetti, M. C. The mechanism of organosphosphate and carbamate resistance in <i>Culex quinquefasciatus</i> (Diptera: Culicidae) from Cuba	245
Chiera, J. W. & Punyua, D. K. Survival of unfed <i>Rhipicephalus appendiculatus</i> (Acarina: Ixodidae) in relation to host resistance and environmental factors in Kenya	251
Doube, B. M. & Giller, P. S. A comparison of two types of trap for sampling dung beetle populations (Coleoptera: Scarabaeidae)	259
Dransfield, R. D., Brightwell, R., Kyorku, C. & Williams, B. Control of tsetse fly (Diptera: Glossinidae) populations using traps at Nguruman, south-west Kenya	265
Duncan, F. D., Nel, A., Batzofin, S. H. & Hewitt, P. H. A mathematical approach to rating food acceptance of the harvester termite, <i>Hodotermes mossambicus</i> (Isoptera: Hodotermitidae) and the evaluation of baits for its control	277
Echendu, T. N. C. & Akingbohungbe, A. E. Intensive free-choice and no-choice cohort tests for evaluating resistance to <i>Maruca testulalis</i> (Lepidoptera: Pyralidae) in cowpea	289
Hunter, D. M. & Cosenzo, E. L. The origin of plagues and recent outbreaks of the South American locust, <i>Schistocerca cancellata</i> (Orthoptera: Acrididae) in Argentina	295
Islam, M. Influence of Scirpophaga incertulas (Lepidoptera: Pyralidae) on deepwater rice	301
Logan, W. M., Cowie, R. H. & Wood, T. G. Termite (Isoptera) control in agriculture and forestry by non-chemical methods: a review	309
Loxdale, H. D. Estimating levels of gene flow between natural populations of cereal aphids (Homoptera: Aphididae)	331
MacVicker, J. A. K., Moore, J. S., Molyneaux, D. H. & Maroli, M. Honeydew sugars in wild-caught Italian phlebotomine sandflies (Diptera: Psychodidae) as detected by high performance liquid chromatography	339
Mansour, F. & Karchi, Z. The evaluation of antibiosis of selected lines for resistance of melon to the carmine spider mite <i>Tetranychus cinnabarinus</i> (Acari: Tetranychidae)	345
Pagnocca, F. C., da Silva, O. A., Hebling-Beraldo, M. J., Bueno, O. C., Fernandes, J. B. &Vieira, P. C. Toxicity of sesame extracts to the symbiotic fungus of leaf cutting ants	349
Rees, D. P. & Walker, A. J. The effect of temperature and relative humidity on population growth of three <i>Liposcelis</i> species (Psocoptera: Liposcelidae) infesting stored products in tropical countries	353

© C·A·B International, 1990

All rights reserved. No part of this publication may be reproduced, in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, https://doi.org/10.1017/S0007485300050410 Published online by Cambridge University Pressr permission of the copyright owner.