

psychiatry's most expensive intervention. These omissions are frustrating because the results suggest that treatment does make a difference; patients who received more of it were less likely to be violent in the following 10 weeks, at least in the early stages of the study. But these are minor quibbles. The strength of this work is illustrated by the dilemma it poses for other researchers: what remains to be done in this field? Precious little, in elucidating the factors that distinguish violent patients from non-violent ones. There were few surprises here, and future surveys will recycle the main variables of personality, previous violence, substance misuse and cultural influences, until we all fall asleep. The unanswered questions are about intervention. How do we apply these findings in clinical practice? Will treatment reduce violence, and can the outcome justify the costs?

Anthony Maden

SWANSON, J. W., HOLZER, C. E., GANJU, V. K., *et al* (1990) Violence and psychiatric disorder in the community: evidence from the Epidemiologic Catchment Area surveys. *Hospital and Community Psychiatry*, **41**, 761–70.


Assertive Outreach in Mental Health. A Manual for Practitioners

Tom Burns & Mike Firn
Oxford University Press, 2002.
355 pp. £24.50 pb.
ISBN: 0-19-851615-0

'This is a handbook primarily for practitioners and not for academics or researchers.' So write Tom Burns & Mike Firn in the opening sentence of the final chapter in their book. I would agree. This is a readable, practical manual covering all aspects of the origins, development and operation of assertive outreach in mental health.

Part I covers 'Conceptual issues' and takes the reader through the origins, context and model for this type of mental health service. There are useful discussions around the target population, and referrals to and discharges from the caseload, with particular emphasis on model fidelity and also on medication, compulsion versus freedom and cultural sensitivity.

Part II on 'Health and social care practice' takes the reader through all the major diagnostic categories one would expect in a service where 'by definition' the target group will be those with severe and enduring mental illnesses, such as bipolar disorder, schizophrenia, personality disorders, substance abuse, and depression and anxiety. However, in addition, the authors usefully explore the problem areas that lie at the roots of why



individuals require assertive outreach: engagement, medication compliance, self-neglect, hospital, suicidality and homelessness.

Part III, 'Structural issues', looks at managing the team, training, service planning, and research and development. The information in this part of the book will be useful for commissioners and service managers, as the authors lay out in detail how to set up an Assertive Outreach Team and how it would fit into the wider mental health system. There are even suggested activities for team building days, such as ten-pin bowling or greyhound racing.

Each part, and indeed each chapter, could be read on its own. The book is an excellent source of material for teaching, learning and debate among practising clinicians of all disciplines and it would be a useful addition to all Mental Health Team libraries. Parts I and III will also help commissioners and managers developing this type of service, a key element of all the frameworks (England and Wales, Scotland and Northern Ireland).

Linda J Watt Consultant Psychiatrist and Medical Director for Mental Health and Learning Disabilities, Greater Glasgow Primary Care NHS Trust, Gartnavel Royal Hospital, Great Western Rd, Glasgow G12 0XH

What Works in Reducing Domestic Violence? A Comprehensive Guide for Professionals

Julie Taylor-Browne (ed.). London: Whiting and Birch, 2001. 396 pp.
£16.95 pb, £47.50 hb.
ISBN: 1-86177-037-5

Is domestic violence a psychiatric issue or only one for child psychiatrists? There are few articles on the subject in the British psychiatric literature and even this excellent little book, which systematically

covers many aspects of the subject, has little to say about psychiatric services.

The book is a collection of reports commissioned by the Home Office crime reduction programme Violence Against Women Initiative. While this occasionally leads to repetition, it ensures that each topic – such as women survivors' views, the needs of children, policing and prosecution, is complete in itself.

From the health point of view, it confirms what has emerged before – that a presentation to a health professional is an opportunity to make an enquiry or confirm a suspicion, which would probably be welcomed by the woman concerned. Professionals, however, are often reluctant to make these enquiries for fear of possibly making matters worse, and anyway, do not know how to offer help. But what has been found to help?

Women survivors of domestic violence found most help from the refuge system, even though these are often crowded and difficult to access. The contributors argue that while much has been done via the establishment of local domestic violence fora to promote interagency collaboration, these may simply add to the burden of work for small voluntary agencies, without supporting their core work. The provision of offender services, excellent in principle, can also be seen as an opportunity cost, especially when successful schemes are difficult to establish.

What should this have to do with a Community Mental Health Team? The sole reference I found on this topic merely suggested that domestic violence was an inappropriate ground for referral by general practitioners. Nevertheless, the psychological consequences may be grave and should be considered. Curiously, in both adult and child mental health, if the assault is sexual then it is more likely to be successfully referred and there is extensive literature in this field. Surely, however,

