

## Short Communication

# Vitamin C deficiency in weanling guinea pigs: differential expression of oxidative stress and DNA repair in liver and brain

Jens Lykkesfeldt<sup>1\*</sup>, Gilberto Perez Trueba<sup>2</sup>, Henrik E. Poulsen<sup>2</sup> and Stephan Christen<sup>3</sup>

<sup>1</sup>Section of Biomedicine, Department of Veterinary Pathobiology, Faculty of Life Sciences, University of Copenhagen, 9 Ridebanevej, DK-1870 Frederiksberg C, Denmark

<sup>2</sup>Department of Clinical Pharmacology Q, Copenhagen University Hospital, 20 Blegdamsvej, DK-2200, Kbh K, Copenhagen, Denmark

<sup>3</sup>Institute of Infectious Diseases, University of Berne, Friedbühlstrasse 51, CH-3010 Berne, Switzerland

(Received 1 February 2007 – Revised 29 March 2007 – Accepted 30 April 2007)

Neonates are particularly susceptible to malnutrition due to their limited reserves of micronutrients and their rapid growth. In the present study, we examined the effect of vitamin C deficiency on markers of oxidative stress in plasma, liver and brain of weanling guinea pigs. Vitamin C deficiency caused rapid and significant depletion of ascorbate ( $P < 0.001$ ), tocopherols ( $P < 0.001$ ) and glutathione ( $P < 0.001$ ), and a decrease in superoxide dismutase activity ( $P = 0.005$ ) in the liver, while protein oxidation was significantly increased ( $P = 0.011$ ). No changes in lipid oxidation or oxidatively damaged DNA were observed in this tissue. In the brain, the pattern was markedly different. Of the measured antioxidants, only ascorbate was significantly depleted ( $P < 0.001$ ), but in contrast to the liver, ascorbate oxidation ( $P = 0.034$ ), lipid oxidation ( $P < 0.001$ ), DNA oxidation ( $P = 0.13$ ) and DNA incision repair ( $P = 0.014$ ) were all increased, while protein oxidation decreased ( $P = 0.003$ ). The results show that the selective preservation of brain ascorbate and induction of DNA repair in vitamin C-deficient weanling guinea pigs is not sufficient to prevent oxidative damage. Vitamin C deficiency may therefore be particularly adverse during the neonatal period.

### Vitamin C deficiency: Weanling guinea pigs: Oxidative stress markers: DNA repair

Pre- and postnatal malnutrition may have serious consequences for neuronal development and growth<sup>1,2</sup>. The impact of vitamin C deficiency, beyond that of scurvy, has not been studied in detail. Vitamin C, for example, is known to play an important role in maintaining brain function<sup>3</sup>. The brain is particularly susceptible to oxidative damage and therefore highly dependent on proper maintenance of redox homeostasis, especially during development, when brain metabolism and growth are maximal<sup>4</sup>. An imbalance in redox homeostasis has also been implicated in neurodegenerative diseases of ageing and other neurological disorders such as schizophrenia<sup>5–8</sup>.

It has been known for a while that vitamin C is preferentially retained in the brain during deficiency<sup>9</sup>. As established recently, this is due to the high-affinity sodium ascorbate cotransporter SVCT2<sup>10</sup>. Experiments with homozygous SVCT2-knockout mice have shown that this transporter is essential for perinatal survival<sup>10</sup>, indicating that vitamin C is crucial for early brain development, although the precise mechanism of death is unknown. Moreover, studies of combined vitamin C and E deficiency in guinea pigs have revealed extensive neuronal damage despite a relatively modest increase in oxidative stress in the brain<sup>11,12</sup>.

Unfortunately, neonatal vitamin C deficiency is fairly common. In a study with 127 pregnant Brazilian women, it was found that 40 % of the smokers and 27 % of the non-smokers had hypovitaminosis C (i.e. a plasma concentration  $< 23 \mu\text{mol/l}$ ) and that this condition was passed on to their fetuses<sup>13</sup>. Major epidemiological studies have shown that this prevalence of hypovitaminosis C can largely be extended to the Western world<sup>14</sup>. Vitamin C deficiency due to inadequate perinatal feeding (e.g. with pasteurized milk) is also fairly common<sup>15</sup>.

Like man, guinea pigs completely depend on dietary vitamin C<sup>16,17</sup>. In the present study, we used newly weaned guinea pigs to test if neonates are susceptible to vitamin C deficiency and if even small changes in antioxidant status in the brain result in deleterious events as measured by oxidative damage.

### Methods

#### Animals and study design

The study was approved by the Danish Animal Experimentation Inspectorate. Dunkin Hartley guinea pigs were born in our animal facility by breeder guinea pigs obtained from

**Abbreviations:** Asc, ascorbate; MDA, malondialdehyde; TBA, thiobarbituric acid.


\* **Corresponding author:** Dr Jens Lykkesfeldt, fax +45 35 35 35 14, email jopl@life.ku.dk

Statens Serum Institut (Allerød, Denmark), that had been maintained at the animal facility for at least 2 weeks. After 2 d, the neonates were taken from their mothers and randomized into two weight-matched groups (control group,  $n$  8; deficient group,  $n$  12). All animals were housed in plastic cages (four per cage) and fed pathogen-free standard diets (Altromin International, Lage, Germany). The diets were essentially identical except for vitamin C content. Control animals were placed on a normal guinea pig diet (Diet #3010, 1036 mg vitamin C/kg)<sup>18</sup>, while deficient animals were supplied with a diet low in vitamin C (Diet #2010, 36 mg vitamin C/kg)<sup>19</sup>. The dietary regimen was continued for about 3 weeks. Animals were checked daily by educated staff and weighed twice a week (Fig. 1). No clinical signs of scurvy were observed during the experiment.

After 3 weeks, the animals were anaesthetized using pentobarbital (55 mg/kg containing 2% lidocaine, by intraperitoneal injection). Heparin (500 units; 100  $\mu$ l) was injected intracardially using a 27G needle and after 1 min a 2 ml blood sample was obtained by cardiac puncture using a 18G needle while carefully avoiding haemolysis. The animals were perfused with 100 ml ice-cold PBS, after which liver and brain were removed and immediately frozen at  $-80^{\circ}\text{C}$  until further analysis. Blood samples were immediately centrifuged (2000g, 5 min,  $4^{\circ}\text{C}$ ). One aliquot (200  $\mu$ l) of plasma was acidified with an equal volume of 10% *meta*-phosphoric acid containing 2 mM-EDTA, briefly vortex mixed and frozen at  $-80^{\circ}\text{C}$  for ascorbate (Asc) and dehydroascorbic acid analysis. The remaining plasma was stored neat in aliquots at  $-80^{\circ}\text{C}$ .

#### Biochemical analyses

Asc and dehydroascorbic acid in *meta*-phosphoric acid-stabilized plasma and tissues were analysed by HPLC with coulometric detection as described previously<sup>20,21</sup>. Because of the very low vitamin C levels in plasma of animals fed with the deficient diet, dehydroascorbic acid was below the detection limit of this assay.  $\alpha$ -Tocopherol and  $\gamma$ -tocopherol were analysed by HPLC with amperometric detection as described<sup>22</sup>. Glutathione was measured as described by Hissin & Hilf<sup>23</sup>. Superoxide dismutase activity was quantified by the pyrogallol method<sup>24</sup>. Plasma oxidizability was quantified essentially as



**Fig. 1.** Weights of 2-d-old neonatal guinea pigs maintained on either a control diet (○) or a vitamin C-deficient diet (●) for 3 weeks. No significant difference was observed over the course of the study period. Values are means with standard deviations depicted by vertical bars.

described by Kontush & Beisiegel<sup>25</sup> except that lag time was objectively based on a sigmoidal curve-fitting model with subsequent calculation of the inflexion. The intercept between the baseline and inflexion was used as lag time.

Malondialdehyde (MDA) was used as an index of lipid oxidation and was assessed by thiobarbituric acid (TBA) derivatization, followed by specific quantification of the genuine MDA(TBA)<sub>2</sub> adduct by HPLC with fluorescence detection as described previously<sup>26</sup>. Protein carbonyls were measured using the ZenTech ELISA kit (Alexis Corporation) based on the method by Buss & Winterbourn<sup>27</sup>. Oxidatively modified DNA as measured by 8-oxo-deoxyguanosine content was quantified by HPLC with electrochemical detection as described<sup>28</sup> and expressed per 10<sup>6</sup> unmodified guanosine residues. Base excision repair was estimated by the nicking technique as described earlier<sup>29</sup>. Protein concentrations were measured by using the Bradford method<sup>30</sup>.

#### Statistical analysis

Effects were analysed by one-way ANOVA followed by *post hoc t*-tests. A two-tailed *P* value less than 0.05 was considered statistically significant. All data are reported as means and standard deviations.

#### Results and discussion

In the present study, we examined the effect of vitamin C deficiency on liver and brain redox homeostasis in weanling guinea pigs. We were particularly interested in the brain, since it is highly susceptible to oxidative damage and known to accumulate high concentrations of Asc, the reduced and anti-oxidative form of vitamin C. Although no significant differences in growth were observed over the course of the study (Fig. 1), the deficient guinea pigs showed clear signs of growth arrest after 3 weeks and would have developed scurvy and died had the experiment been allowed to continue. Apart from its antiscorbutic effect, vitamin C plays a pivotal role in redox homeostasis<sup>3,31,32</sup>.

Weanling guinea pigs fed for 3 weeks on a vitamin C-deficient diet displayed severe depletion of plasma and liver Asc, showing less than 1% Asc compared to control animals ( $P < 0.001$ ; Table 1). In contrast, the depletion of Asc in the brain was only about 70%. This preferential preservation of vitamin C in the brain is in agreement with other studies<sup>9,33</sup> and is most likely the result of active transport of vitamin C across the blood-brain barrier.

Asc depletion in plasma and liver was accompanied by a general change in the antioxidant status. Thus, vitamin C deficiency caused a decrease of  $\alpha$ - and  $\gamma$ -tocopherol in plasma and liver, and resulted in lower glutathione levels and superoxide dismutase activity in the liver, suggesting a collapse of the redox homeostasis in this tissue (Table 1). In contrast, Asc was the only antioxidant in the brain that was significantly decreased. The significant 2.5-fold increase in dehydroascorbic acid observed in vitamin C-deficient animals ( $P = 0.034$ ) is indicative of elevated oxidative stress in the brain. Thus, increased antioxidant action of Asc combined with inadequate recycling capacity for vitamin C results in increased turnover as recognized by the increased presence of dehydroascorbic acid.

**Table 1.** Biomarkers of oxidative stress and damage in plasma, liver and brain of weanling guinea pigs after 3 weeks on a vitamin C-deficient diet compared to animals on a control diet

(Mean values and standard deviations)

	Controls ( <i>n</i> 8)		Deficient ( <i>n</i> 12)		<i>P</i> value*
	Mean	SD	Mean	SD	
<b>Plasma</b>					
Ascorbate ( $\mu\text{M}$ )	168.3	49.1	1.0	0.5	<0.001
$\alpha$ -Tocopherol ( $\mu\text{M}$ )	5.8	1.8	3.8	1.4	0.017
$\gamma$ -Tocopherol ( $\mu\text{M}$ )	0.10	0.04	0.02	0.01	<0.001
Malondialdehyde ( $\mu\text{mol/l}$ )	0.34	0.10	0.61	0.15	0.013
Plasma oxidizability lag time (min)	268	25	62	30	<0.001
<b>Liver</b>					
Ascorbate (nmol/g tissue)	2835	623	26.2	4.6	<0.001
Dehydroascorbic acid (% of total vitamin C)	3.3	1.8	5.3	9.7	NS
$\alpha$ -Tocopherol (nmol/g tissue)	33.8	9.9	15.0	3.2	<0.001
$\gamma$ -Tocopherol (nmol/g tissue)	1.0	0.4	0.2	0.1	<0.001
Glutathione (nmol/g tissue)	3511	490	1822	755	<0.001
Superoxide dismutase ( $\mu\text{g/mg}$ protein)	7.8	0.6	6.5	1.1	0.005
Malondialdehyde (nmol/g tissue)	128	15	127	23	NS
Protein carbonyls (nmol/mg protein)	151	5	171	18	0.011
8-Oxo-deoxyguanosine ( $1/10^6$ dG)	12.9	4.6	11.4	3.8	NS
Base excision repair (%; arbitrary units)	1.22	0.14	1.04	0.22	NS
<b>Brain</b>					
Ascorbate (nmol/g tissue)	1450	138	453	80	<0.001
Dehydroascorbic acid (% of total vitamin C)	6.5	8.2	14.8	4.0	0.034
$\alpha$ -Tocopherol (nmol/g tissue)	17.0	1.7	15.3	2.4	NS
$\gamma$ -Tocopherol (nmol/g tissue)	0.4	0.1	0.4	0.1	NS
Glutathione (nmol/g tissue)	1189	80	1146	92	NS
Superoxide dismutase ( $\mu\text{g/mg}$ protein)	2.6	0.3	2.5	0.5	NS
Malondialdehyde (nmol/g tissue)	186	47	429	79	<0.001
Protein carbonyls (pmol/mg protein)	77.6	5.5	64.3	9.0	0.003
8-Oxo-deoxyguanosine ( $1/10^6$ dG)†	3.18	0.52	3.71	0.51	NS
Base excision repair (%; arbitrary units)	0.24	0.05	0.33	0.09	0.014

dG, unmodified guanosine residues.

\* Compared to control group.

† Control, *n* 4; deficient, *n* 8.

Vitamin C deficiency was also associated with increased oxidative damage. In plasma, lipid oxidation was significantly increased in vitamin C-deficient animals ( $P=0.013$ ) and the resistance to lipid oxidation (lag time) was significantly diminished to less than one-quarter of that of animals fed with the normal diet ( $P<0.001$ ). In the liver, protein oxidation was increased ( $P=0.011$ ) while lipid and DNA oxidation were unchanged.

As with the antioxidants, the pattern of oxidative damage was different in the brain. Here, MDA was increased by 130% in deficient animals compared to that of controls ( $P<0.001$ ). In contrast, protein oxidation was significantly lower ( $P=0.003$ ). Whether this is related to vitamin C's role in protein synthesis is not known. With regard to DNA damage, both oxidatively damaged DNA and DNA repair capacity was assessed, the latter as 8-oxo-deoxyguanosine glycosylase activity. Since *de novo* synthesis of DNA is only possible during cell proliferation, the capacity to repair damaged DNA is of major importance for the ability of the cells to survive an oxidative insult. In the brain, DNA incision repair was 38% higher in deficient animals compared to controls ( $P=0.014$ ), suggesting an up-regulation of DNA repair in response to the increased oxidative stress. This response was apparently adequate, since 8-oxo-deoxyguanosine did not significantly increase in the vitamin C-deficient animals ( $P=0.13$ ).

Recently, it was reported that *in vitro*-cultured cells do not respond to oxidative DNA damage by inducing repair<sup>34</sup>. The authors concluded that DNA damage *per se* does not induce DNA repair<sup>34</sup>. However, it is possible that DNA repair was already maximally induced in their *in vitro* system, or that induction was no longer possible due to cell transformation. Our *in vivo* data, however, suggest that either the reduced levels of antioxidants *per se* or a secondary effect of such depletion indeed do induce DNA repair *in vivo*.

Neonatal guinea pigs appear to be more susceptible to malnutrition compared to their older counterparts, possibly due to their limited reserve of micronutrients and their rapid growth. This became apparent in the present study as the neonates began suffering growth arrest and even weight loss, usually among the first signs of emerging lameness in the hind limbs and clinical scurvy, already after 3 weeks (Fig. 1), while in older animals (e.g. 3 months old) the same condition is normally reached only after more than 5 weeks on a deficient diet<sup>35</sup>. Thus, neonates appear to be particularly susceptible to vitamin C deficiency.

## Conclusion

The present results show that in weanling guinea pigs, vitamin C deficiency results in altered brain redox homeostasis and increased lipid oxidation. Preferential preservation of vitamin C in the brain over other tissues, antioxidant function of Asc

and induction of DNA incision repair provide some protection against oxidative damage to DNA and proteins. To strengthen the link between oxidative stress and disease, future studies should include the evaluation of brain injury. Also, the expression of SVCT2 in the brain during vitamin C deficiency may be of importance in understanding the preferential preservation of vitamin C in the brain.

### Acknowledgements

Annie B. Kristensen, Jytte Nielsen, Corinne Siegenthaler, Jane Povlsen and the KVL animal facility are thanked for excellent technical assistance. This study was supported by National Research Council grant 21-01-0525 to J. L. The laboratory of S. C. is supported by grants from the Swiss National Science Foundation (3100-108236) and National Institute of Neurological Disorders and Stroke (R01 NS33997).

### References

- Guesry P (1998) The role of nutrition in brain development. *Prev Med* **27**, 189–194.
- Morgane PJ, Austin-LaFrance R, Bronzino J, Tonkiss J, az-Cintra S, Cintra L, Kemper T & Galler JR (1993) Prenatal malnutrition and development of the brain. *Neurosci Biobehav Rev* **17**, 91–128.
- Rice ME (2000) Ascorbate regulation and its neuroprotective role in the brain. *Trends Neurosci* **23**, 209–216.
- Rougemont M, Do KQ & Castagne V (2002) New model of glutathione deficit during development: effect on lipid peroxidation in the rat brain. *J Neurosci Res* **70**, 774–783.
- Bains JS & Shaw CA (1997) Neurodegenerative disorders in humans: the role of glutathione in oxidative stress-mediated neuronal death. *Brain Res Brain Res Rev* **25**, 335–358.
- Mahadik SP & Mukherjee S (1996) Free radical pathology and antioxidant defense in schizophrenia: a review. *Schizophr Res* **19**, 1–17.
- Mukerjee S, Mahadik SP, Scheffer R, Correnti EE & Kelkar H (1996) Impaired antioxidant defense at the onset of psychosis. *Schizophr Res* **19**, 19–26.
- Schulz JB, Lindenau J, Seyfried J & Dichgans J (2000) Glutathione, oxidative stress and neurodegeneration. *Eur J Biochem* **267**, 4904–4911.
- Kuo CH, Yonehara N & Yoshida H (1979) Subcellular ascorbic acid in scorbutic guinea pig brain. *J Nutr Sci Vitaminol (Tokyo)* **25**, 9–13.
- Sotiriou S, Gispert S, Cheng J, *et al.* (2002) Ascorbic-acid transporter Slc23a1 is essential for vitamin C transport into the brain and for perinatal survival. *Nat Med* **8**, 514–517.
- Burk RF, Christensen JM, Maguire MJ, Austin LM, Whetsell WO Jr, May JM, Hill KE & Ebner FF (2006) A combined deficiency of vitamins E and C causes severe central nervous system damage in guinea pigs. *J Nutr* **136**, 1576–1581.
- Hill KE, Montine TJ, Motley AK, Li X, May JM & Burk RF (2003) Combined deficiency of vitamins E and C causes paralysis and death in guinea pigs. *Am J Clin Nutr* **77**, 1484–1488.
- Madruga de Oliveira A, de Carvalho Rondo PH & Barros SB (2004) Concentrations of ascorbic acid in the plasma of pregnant smokers and nonsmokers and their newborns. *Int J Vitam Nutr Res* **74**, 193–198.
- Lykkesfeldt J (2006) Smoking depletes vitamin C: should smokers be recommended to take supplements? In *Cigarette Smoke and Oxidative Stress* pp. 237–260 [B Halliwell and HE Poulsen, editors]. Berlin: Springer Verlag.
- Ratanachu-Ek S, Sukswai P, Jeerathanyasakun Y & Wongtpradit L (2003) Scurvy in pediatric patients: a review of 28 cases. *J Med Assoc Thai* **86**, Suppl. 3, S734–S740.
- Chatterjee IB (1973) Evolution and the biosynthesis of ascorbic acid. *Science* **182**, 1271–1272.
- Nandi A, Mukhopadhyay CK, Ghosh MK, Chattopadhyay DJ & Chatterjee IB (1997) Evolutionary significance of vitamin C biosynthesis in terrestrial vertebrates. *Free Radic Biol Med* **22**, 1047–1054.
- Altromin (2007) Altromin Diet 3010 Composition. <http://www.altromin.de/content/db.php?nummer=3010&sprache=de>.
- Altromin (2007) Altromin Diet 2010 Composition. <http://www.altromin.de/content/db.php?nummer=2010&sprache=de>.
- Lykkesfeldt J (2000) Determination of ascorbic acid and dehydroascorbic acid in biological samples by high-performance liquid chromatography using subtraction methods: reliable reduction with Tris[2-carboxyethyl]phosphine hydrochloride. *Anal Biochem* **282**, 89–93. [Published erratum appears in *Anal Biochem* (2001) 292, 173].
- Lykkesfeldt J (2002) Measurement of ascorbic acid and dehydroascorbic acid in biological samples. In *Current Protocols in Toxicology*, pp. 761–7615 [M Maines, LG Costa, E Hodson and JC Reed, editors]. New York: John Wiley & Sons.
- Sattler W, Mohr D & Stocker R (1994) Rapid isolation of lipoproteins and assessment of their peroxidation by high-performance liquid chromatography postcolumn chemiluminescence. *Meth Enzymol* **233**, 469–489.
- Hissin PJ & Hilf R (1976) A fluorometric method for determination of oxidized and reduced glutathione in tissues. *Anal Biochem* **74**, 214–226.
- Marklund S & Marklund G (1974) Involvement of the superoxide anion radical in the autoxidation of pyrogallol and a convenient assay for superoxide dismutase. *Eur J Biochem* **47**, 469–474.
- Kontush A & Beisiegel U (1999) Measurement of oxidizability of blood plasma. *Methods Enzymol* **299**, 35–49.
- Lykkesfeldt J (2001) Determination of malondialdehyde as dithiobarbituric acid adduct in biological samples by HPLC with fluorescence detection: comparison with ultraviolet-visible spectrophotometry. *Clin Chem* **47**, 1725–1727.
- Buss IH & Winterbourn CC (2002) Protein carbonyl measurement by ELISA. *Methods Mol Biol* **186**, 123–128.
- Gedik CM & Collins A (2005) Establishing the background level of base oxidation in human lymphocyte DNA: results of an interlaboratory validation study. *FASEB J* **19**, 82–84.
- Riis B, Risom L, Loft S & Poulsen HE (2002) OGG1 mRNA expression and incision activity in rats are higher in foetal tissue than in adult liver tissue while 8-oxo-2'-deoxyguanosine levels are unchanged. *DNA Repair (Amst)* **1**, 709–717.
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**, 248–254.
- Buettner GR & Schafer FQ (2004) Ascorbate as an antioxidant. In *Vitamin C: Its Functions and Biochemistry in Animals and Plants*, pp. 173–188 [H Asard, JM May and N Smirnoff, editors]. Oxford: BIOS Scientific Publishers.
- Rice ME & Russo-Menna I (1998) Differential compartmentalization of brain ascorbate and glutathione between neurons and glia. *Neuroscience* **82**, 1213–1223.
- Hughes RE, Hurlley RJ & Jones PR (1971) The retention of ascorbic acid by guinea-pig tissues. *Br J Nutr* **26**, 433–438.
- Bercht M, Flohr-Beckhaus C, Osterod M, Runger TM, Radicella JP & Epe B (2007) Is the repair of oxidative DNA base modifications inducible by a preceding DNA damage induction? *DNA Repair (Amst)* **6**, 367–373.
- Lykkesfeldt J (2002) Increased oxidative damage in vitamin C deficiency is accompanied by induction of ascorbic acid recycling capacity in young but not mature guinea pigs. *Free Radic Res* **36**, 567–574.