

Animal performance and economic implications of Holstein-Friesian, Jersey and Jersey × Holstein-Friesian cows under seasonal pasture based systems

R. Prendiville^{1,2}, L. Shalloo¹, K.M. Pierce², F. Buckley¹

¹Teagasc, Moorepark, Dairy Production Research Centre, Fermoy, Co. Cork, Ireland

²School of Agriculture, Food Science and Veterinary Medicine, UCD, Belfield, Dublin 4, Ireland

Email: robert.prendiville@teagasc.ie

Introduction In seasonal production systems such as Ireland the aim is to optimise milk production from pasture with limited concentrate supplementation. Ideally cows in such systems should efficiently convert pasture to product and maintain a calving interval around 365 days. Crossbreeding, particularly with the Jersey (J), is common practice in New Zealand and Lopez-Villalobos *et al.* (2000) reported Jersey×Holstein-Friesian (F₁) cows to be more profitable than the parent breeds. The J breed would appear to offer potential for crossbreeding under Irish conditions with its small size, the prospect of improved reproductive performance, and high milk components which with the introduction of multi component price systems of payment is of particular interest in adding value to milk (Shalloo, 2007). The aim of this study was to provide comparative cow performance data and the implications of same for overall farm profitability of Holstein-Friesian (HF), J and F₁ cows under Irish seasonal pasture-based management.

Material and methods A total of 329 lactations from 65 HF, 48 J and 49 F₁ cows were available from a three year study; 2006, 2007 and 2008 (years 1, 2 and 3, respectively), conducted at the 'Ballydague' research farm. Cows were managed in a seasonal production system (Dillon *et al.*, 1995). Milk yield was recorded daily while milk components and bodyweight were determined weekly. A more detailed description of fertility measures is provided by Prendiville *et al.* (2010). The Moorepark Dairy Systems Model (MDSM) (Shalloo *et al.*, 2004), a stochastic budgetary simulation model was used to simulate a model farm integrating biological data for each breed group. This model incorporates animal inventory and valuation, milk production, feed requirement, land, labour and economic analysis. Variable costs including fertilizer, contractor charges, medical and veterinarian, artificial insemination, silage and reseeded, fixed costs (machinery maintenance and running costs, farm maintenance, car, telephone, electricity and insurance) and sales values (milk, cull cow and calf) were included at current prices (Teagasc, 2008). The model was limited to 40 hectare of land and 13t DM of grass/ha was assumed to be grown. Fertilizer application was assumed to be 250 kg of N/ha. A milk price of 27c/l at 33.0g/kg protein and 36.0g/kg fat with a ratio of the value of protein to fat of 2.6 to 1 was used. Cull cow price was assumed to be €366, €149 and €268 for the HF, J and F₁, respectively (for details see Prendiville *et al.*, 2010). All calves were assumed sold at 4 weeks of age. Male calf value was €80, €0 and €30 for HF, J and F₁, respectively, while female calves were valued at €330. Replacement rate was calculated as the proportion of cows that failed to become pregnant by the end of the 13-week breeding season plus a voluntary culling rate of 10% of the remaining cows. Fertility values obtained for the HF and J were not statistically different. Consequently, empty rates were 18% for the HF and J and 9% for the F₁. Due to the differences in replacement rates milk yield was adjusted for parity structure. Differences in calving date were also assumed based on differences in calving to conception interval.

Results According to the model 96 HF, 114 J and 97 F₁ cows would be facilitated on the 40 ha land base. Milk yield was highest with the HF (5,651 kg), intermediate with the F₁ (5,272 kg) and lowest with the J (4,220 kg). Milk fat and protein content were highest for the J (5.32% and 4.03%), intermediate with the F₁ (4.77% and 3.88%) and lowest with HF (4.12% and 3.49%). Consequently, milk solids production was highest for the F₁ (456 kg) intermediate for the HF (430kg) and lowest for the J (395 kg). Milk returns were highest for the J (€172,816), intermediate for the F₁ (€171,790) and lowest with the HF (€158,675). Due to additional animal numbers labour cost was higher with the J (€32,811) compared to the HF and F₁ (€27,760 and €28,463, respectively). Replacement costs were lowest with the F₁ (€26,935), intermediate with the HF (€38,904) and highest with the J (€45,982). Livestock sales were highest for the HF (€28,675) and similar for the J and F₁ (€22,696 and €21,674). Total costs were €149,852, €167,089 and €137,786 for the HF, J and F₁, respectively. However, overall farm profit was highest with the F₁ (€55,678) followed by the HF (€37,499) and J (€28,423). Farm profitability per hectare was highest for the F₁ (€1,392), intermediate for the HF (€938) and lowest for the J (€711). The additional profit generated by the F₁ is largely attributed to increased milk value and lower replacement costs i.e. improved reproductive efficiency.

Conclusions In Ireland the impending removal of EU milk quotas will result in land becoming the most limiting resource. Although many farmers view crossbreeding as a means of improving reproductive performance and herd health, results from this study indicate that despite the lower livestock sales (cull cow and calf revenue), overall farm profitability with F₁ cows may well be higher compared to HF cows, due to greater milk receipts coupled with improved reproductive efficiency and survival.

Acknowledgements Financial support from the Research Stimulus Fund (RSF-06-353) is gratefully acknowledged.

References

- Dillon, P., Crosse, S., Stakelum, G. and Flynn, F. 1995. *Grass Forage Science* 50, 286-299.
- Lopez-Villalobos N., Garrick, D.J., Holmes, C.W., Blair, H.T. and Spelman, R.J. 2000. *Journal of Dairy Science* 83, 144-153.
- Prendiville R. 2010. PhD. Thesis, National University of Ireland, Dublin, Ireland.
- Shalloo L., Dillon, P., Rath, M. and Wallace, M. 2004. *Journal of Dairy Science* 87, 1945-1959.
- Teagasc. 2008. Management data for farm planning. Teagasc, Oakpark, Carlow, Ireland.