

Pulling Out Threads from the Cosmic Tapestry: Defining Filaments of Galaxies

Kevin A. Pimbblet^A

^A Department of Physics, University of Queensland, Brisbane QLD 4072, Australia.
Email: pimbblet@physics.uq.edu.au

Received 2005 February 1, accepted 2005 March 9

Abstract: Filaments of galaxies are the dominant feature of modern large-scale redshift surveys. They can account for up to perhaps half of the baryonic mass budget of the Universe and their distribution and abundance can help constrain cosmological models. However, there remains no single, definitive way in which to detect, describe, and define what filaments are and their extent. This work examines a number of physically motivated, as well as statistical, methods that can be used to define filaments and examines their relative merits.

Keywords: large-scale structure of Universe — cosmology: observations — methods: observational

1 Motivation

What is a filament of galaxies (FOG)? Although at first glance, this is a seemingly innocuous, benign, and non-trivial question, there is not really a straightforward answer to it. Many authors, including the present one, have recently been searching for more concrete definitions and hence, also, methods of finding and detecting FOGs (e.g. Pimbblet 2005 and references therein).

The present work is therefore a timely review of the (growing) myriad of approaches that exist to define and detect FOGs in an attempt to answer the question of what a FOG actually is. In a lot of ways, investigations of FOGs nowadays are arguably analogous to where the investigations of galaxy clusters stood at about one half of a century ago (see Abell 1965 for an excellent, albeit somewhat dated by today's standards, review of galaxy clustering). Undoubtedly, the reason for the recent flurry of activity into investigating and characterising FOGs has to be the availability of modern, high-quality and, most importantly, wide-field redshift surveys such as the 2dF Galaxy Redshift Survey (2dFGRS; e.g. Colless et al. 2001), the Sloan Digital Sky Survey (SDSS; e.g. Abazajian et al. 2004), the 6dF Galaxy Survey (6dFGS; e.g. Jones et al. 2004), and the Las Campanas Redshift Survey (LCRS; e.g. Shectman et al. 1996).

Having written that, the discovery of significant segments of large-scale structure other than galaxy clusters — sheets, filaments, and walls of galaxies — is not a new thing. Geller & Huchra (1989) famously cartographed the 'Great Wall' from the CfA redshift survey (e.g. Huchra et al. 1983): a highly significant feature that stretches for at least $170 \times 60 h_{100}^{-1}$ Mpc at $cz \approx 7500 \text{ km s}^{-1}$ (Figure 1). So significant is this detection, that it should even have an imprint on the cosmic microwave background radiation (Atrio-Barandela & Kashlinsky 1992; see also Chodorowski 1994).

What else may we expect from FOGs? We know that in hierarchical structure formation modelling there has long been the prediction that galaxy clusters grow through repeated mergers with other galaxy clusters (and galaxy groups) together with continuous accretion of their surrounding matter (e.g. Zeldovich, Einasto, & Shandarin 1982; Katz et al. 1996; Jenkins et al. 1998; Colberg et al. 2000; see also Bond, Kofman, & Pogosyan 1996). We also know that the accretion process occurs in a highly non-isotropic manner: galaxy filaments funnel matter onto large clusters along preferred directions (see Pimbblet 2005; Ebeling, Barrett, & Donovan 2004; Kodama et al. 2001). Beyond a few virial radii from galaxy clusters centres, FOGs are predicted to weave a complex, web- or sponge-like tapestry that gives surveys like SDSS, 6dFGS, 2dFGRS, and LCRS their distinctive appearance (Figure 2; see also the 2dFGRS homepage at www.mso.anu.edu.au/2dFGRS).

We also know that FOGs are highly important for the mass budget of the Universe (e.g. Colberg et al. 1999). Indeed, Cen & Ostriker (1999) show that for a Λ cold dark matter (Λ CDM) Universe, a large fraction, perhaps as much as half (Fukugita, Hogan, & Peebles 1998), of baryonic material will not have been observed as it is situated in the inter-cluster media in a hot and tenuous gaseous phase. Along with the dark matter component and perhaps up to a quarter of the galaxian population, these baryons are preferentially situated in (inter-cluster) FOGs. Moreover, FOGs can provide tests of structure formation (cf. Colberg, Krughoff, & Connolly 2005 with Pimbblet, Drinkwater, & Hawkrigg 2004) and cluster evolution (see Colberg et al. 1999). They can also be useful in ascertaining the homogeneity scale of our Universe (if, indeed one considers there to be a homogeneity scale; e.g. Coleman & Pietronero 1992 and references therein). Certainly, given that objects with scale lengths $> 150 h_{100}^{-1}$ Mpc exist and




Figure 1 The Great Wall of Geller & Huchra (1989) reconstructed using J2000 coordinates from the November 1993 public data release of the CfA dataset. Following Geller & Huchra (1989), all galaxies within a declination range of $20 < \delta < 40$ are plotted with no cut in magnitude made. The radius of the circle is $15\,000\text{ km s}^{-1}$. The Great Wall can be seen extending outward from 13 hours.

are not chance superpositions, we should be questioning the validity of the cosmological assumption up to such lengths and beyond.

The rest of this paper plans out as follows. In Section 2, we investigate the numerous methods that one can employ to detect FOGs and explore their relative merits. In Section 3 we discuss the findings and present our conclusions.

2 Detection

Already we have seen a number of properties of FOGs. If sufficiently large, then they can cause a decrement in the cosmic microwave background radiation. They also possess multi-wavelength visibility (visual, X-ray, etc.). We describe below how one may take advantage of such properties to detect them in a given dataset.

2.1 Optical Overdensity

At a very simple level, a FOG is merely an overdensity of galaxies compared to the local field¹ (or void) level.

¹ Here, we use the term ‘field’ to mean the ‘average background’ level.

Pimblet & Drinkwater (2004) use this fact to find a relatively short ($\sim 6 h_{100}^{-1}$ Mpc) filament between the two close (both in redshift and spatially) galaxy clusters ACO1079 and ACO1084. Mathematically, one can readily compute this galaxy excess as

$$N_{\text{filament}} = N_{\text{filament+field}} - N_{\text{field}}. \quad (1)$$

Should the observed field sample be too close to the observed filament sample it will obviously contain some (small but non-negligible) amount of contamination:

$$N'_{\text{field}} = N_{\text{field}} + \gamma N_{\text{filament}}, \quad (2)$$

where γ is the ratio between the galaxy densities of the filament and field populations (Paolillo et al. 2001). Substituting N'_{field} instead of N_{field} from Eqn (2) into Eqn (1) gives

$$N'_{\text{filament}} = N_{\text{filament}}(1 - \gamma). \quad (3)$$

In Figure 3 we plot an adaptation of the result obtained by Pimblet & Drinkwater (2004) utilizing this particular method. Most of the excess galaxy population is faint,




Figure 2 Same for Figure 1, but using data from 2dFGRS. No cut has been made in declination or b_j magnitude, although the plot has been constrained to the same redshift limit as Figure 1, albeit with a different declination range. Note the complex manner in which FOGs weave through the structure and its qualitative similarity to the CfA survey.

with only a few brighter members. Moreover, only a small fraction ($\approx 30\%$) of these galaxies have colours consistent with early-type galaxies from the two clusters colour–magnitude relations (Pimblet et al. 2002). In shallow (perhaps, mono-colour) surveys with no supporting redshift information, therefore, such an approach is probably not very efficient nor exceedingly sensitive and may be somewhat prone to large errors.

2.2 X-ray

Given that a non-negligible fraction of baryonic material in a Λ CDM Universe may exist as hot inter-cluster gas, one can also consider looking for FOGs in X-ray band passes. Using the *ROSAT* All-Sky Survey data, Briel & Henry (1995) attempted just this by combining together the inter-cluster regions of 40 cluster pairs. Although they failed to find any X-ray emitting FOG, they did place an upper limit on the X-ray surface brightness of $4 \times 10^{-16} \text{ ergs cm}^{-2} \text{ s}^{-1}$ (0.5–2.0 keV).

Scharf et al. (2000) make a 5σ joint X-ray/optical detection of $>12 h_{50}^{-1} \text{ Mpc}$ (0.5 deg) FOG with a surface brightness of $1.6 \times 10^{-16} \text{ ergs cm}^{-2} \text{ s}^{-1}$. The count rate

for this filament, however, is $2 - 3\sigma$ above background levels.

In the Shapley supercluster meanwhile, Kull & Böhringer (1999) find a promising extended X-ray emission between a close cluster pair that is ~ 2.5 times brighter than Briel & Henry's (1995) bound. The only problem here is that Scharf et al. (2000) note that the X-ray emission could be ejecta due to the clusters interacting (merging) with one another rather than material falling in from an actual filament.

More promising progress on this front has been made by Tittley & Henriksen (2001) and Durret et al. (2003). The former detect a gaseous FOG between ACO3391 and ACO3395 that has a minimum flux of $1.3 \times 10^{-12} \text{ ergs cm}^{-2} \text{ s}^{-1}$ (0.8–10 keV) and represents at least 2% of the total mass of the system. The latter team study the ACO85 cluster complex and find a highly elongated filamentary structure. Again, however, this filament may be the result of cluster interactions and not a true FOG in the large-scale structure sense.

It would seem that whilst one may expect there to be significant X-ray emission from baryonic material contained in FOGs, we are not quite detecting it with sufficient




Figure 3 Number of galaxies in excess of the field population (i.e. the filament population) as a function of absolute magnitude (adapted from the investigation of Pimblet & Drinkwater 2004). The $\pm 1\sigma$ errorbars come from a consideration of Poissonian errors and the variance galaxy number density. Whilst there are only a few brighter galaxies, the FOG contains many more fainter ones.

regularity or confidence to use X-ray emission as the primary tool for FOG detection (unlike in the case of galaxy clusters where X-ray detections are made with much more confidence; e.g. Ebeling et al. 1996).

2.3 Lensing

Pogosyan et al. (1998) point out that FOGs that connect together neighbouring galaxy clusters should have sufficient surface mass density as to be detectable in the weak lensing regime. Indeed, weak lensing would only depend upon the projected density and not the square of the projected density like X-rays are (Pogosyan et al. 1998) and therefore it may constitute an altogether better way of detecting and defining FOGs. There are a small number of investigations that have been proceeding in this direction.

Kaiser et al. (1998) perform a lensing analysis on the $z = 0.4$ supercluster MS 0302+17 and find a FOG between two of its three component galaxy clusters. The detection has remained dubious, however, as there may be foreground structure interfering, perhaps some edge of chip effects and residual systematics in the point spread function anisotropy correction involved (Gavazzi et al. 2004). Indeed, Gavazzi et al. (2004) report that they cannot independently confirm the detection of this particular FOG.

Meanwhile, in other investigations, Clowe et al. (1998) report on the detection of a FOG extending from the $z = 0.8$ rich cluster RXJ 1716+67. However, the size of their imaging is small and it is thus unknown how far this filament extends in the direction of a nearby cluster. Gray et al. (2002) examine the ACO901/902 supercluster

and find a FOG present. The significance of the detection is, however, small. Superposed with this is the issue that the filament lies in the inter-chip region of the analysis. Nonetheless, this remains a relatively good detection when compared to the problems that Kaiser et al. (1998) encounter.

Yet to date, arguably one of the best weak lensing FOG detections has to be that of Dietrich et al. (2004) between ACO222 and ACO223. Not only is the filament detected by weak lensing, but Dietrich et al. (2004) supply supporting evidence from X-ray emission and increased galaxy density between the clusters. Dietrich et al. (2004) point out, however, that they could not find an objective way to define their filament and in this respect their filament candidate is not very different to that of Kaiser et al. (1998).

So, it would appear that given imaging of sufficient quality and depth, weak lensing could provide an excellent way of detecting FOGs, most especially in combination with other methods (e.g. X-ray; see above).

2.4 Redshifts

Redshifts of regions around galaxy clusters can provide concrete determinations of the presence of FOGs. For example, Ebeling, Barrett, & Donovan (2004) report on a $4 h_{70}^{-1}$ Mpc filament that is feeding the growth of the massive cluster MACS J0717.5+3745 at $z = 0.55$. Its extent beyond the virial radius of the cluster means that it cannot be the remnant of some previous interaction or merger whilst its colours are quite consistent with the colour-magnitude relation (CMR; e.g. Visvanathan & Sandage 1977; Bower, Lucey, & Ellis 1992). Indeed, the CMR and other photometric redshift techniques can also help to better define FOGs. Kodama et al. (2001) report several ‘octopus’-like tentacles around ACO851 ($z = 0.41$) which have colours entirely consistent with the CMR of the cluster itself. Pimblet, Edge, & Couch (2005) locate a large-scale wall covering at least $40 h_{100}^{-1}$ Mpc situated in front of ACO22 ($z = 0.14$; Figure 4). Not only does this wall exhibit a CMR similar to ACO22, but it also has a Butcher & Oemler (1984; see Pimblet 2003 for a review of the Butcher–Oemler effect) blue fraction that does not change significantly between the cluster and the wall (Figure 4).

The above are examples of FOGs around individual clusters. Of course, it is the modern Large-redshift surveys such as 2dFGRS, 6dFGS, SDSS, and LCRS that are providing the community with an unprecedented view of the very large-scale structure of the Universe (e.g. Figure 2). With such large datasets, finding individual filaments can become as easy as looking at the regions between two galaxy clusters in three dimensional space and making an appropriate cut at some galaxy density threshold² to determine if there is a significant

² Pimblet, Edge, & Couch (2005) note that the typical surface density of FOGs is of the order 10 bright (brighter than say $M^* + 2$) galaxies per square h^{-1} Mpc.




Figure 4 The wall in front of ACO22 discovered by Pimblet, Edge, & Couch (2005). The upper panel shows the redshift distribution of the cluster (right-hand peak; downward arrow) and the wall (left-hand side; $cz < 40\,000\text{ km s}^{-1}$). The lower panel shows how the $(B - R)$ colours of the galaxies vary with redshift. The colour distribution and Butcher–Oemler blue fraction is statistically the same (within 2σ) between the cluster and the wall when appropriate magnitudinal cuts are made (Pimblet, Edge, & Couch 2005).

overdensity of galaxies present (e.g. Pimblet, Drinkwater, & Hawkrigg 2004). One potential pitfall is that one may mistake a redshift space distortion (see Hawkins et al. 2003) for a FOG. Pimblet, Drinkwater, & Hawkrigg (2004) circumvent this by only considering cluster-cluster pairs within 1000 km s^{-1} of each other and check the FOG distribution angles along the line of sight to ensure that no ‘fingers of god’ are mistaken for FOGs (and conversely, no end-on FOG is mistaken for a cluster!). Also, surveys like 2dFGRS only cream off the very luminous galaxies (and even at bright magnitudes, are incomplete; Cross et al. 2004). They tell us little about the low surface brightness populations which likely contribute a non-negligible fraction of any FOG’s mass. The next-generation of deeper redshift surveys (e.g. using AAΩ on the Anglo-Australian Telescope) should help us to address this point.

We note in passing, however, that one potentially unanswered question is whether a FOG can exist that is not connected to any cluster? If they can, this would bias the results of these kind of investigations which focus exclusively on intercluster regions as the locations to search out FOGs. Recent work by Fairall et al. (2004) suggests, however, that there can be no isolated FOGs.

2.5 Further Statistical Methods

Given that the large-redshift surveys may contain several hundred FOGs, it can be better to approach the entire FOG population in a more consistent (and less time-consuming) manner. There are a multitude of statistics available for the analysis of large-scale structure and FOGs, and here

we only review a small, representative selection of these approaches.

Some readers may be familiar with the children’s game ‘connect the dots’ (CTD herein). The idea is to connect a series of points on a plane, in a particular sequence, in order to form some kind of picture at the end of it. One well known relative of CTD is the travelling salesman problem where the challenge is to connect some points in by the shortest possible route. Yet finding FOGs in redshift surveys like 2dFGRS can also be thought of as another variant of CTD types of games (Arias-Castro et al. 2004). Consider a toy-model: a two dimensional curve that one randomly³ samples points from along its length (i.e. galaxies contained in a FOG). Add in some random noise, and now the problem becomes whether we are able to recover the original curve (FOG) in the presence of clutter (field galaxies; chance superpositions; etc.). Donoho and collaborators (e.g. Arias-Castro et al. 2004) approach the CTD problem from a number of angles. One promising method is to make use of a multiscale adaptive geometric analysis. By using a zoo of parallelogram strips of various angles, lengths, widths, and eccentricities one can evaluate (count) the galaxy population in all such strips. Then all one has to do is identify strips with unusually high-count rates and search for long runs of such strips that would constitute a good continuation of a given curve. The problem with this analogy is that FOGs are not perfect lines or curves: they can be bumpy, lumpy, and clumpy in three dimensions (Figure 5). Finding a FOG that is only just above the density of the clutter can prove to be hard (Figure 5), especially if one is interested in the morphology of FOGs.

However, if one can make use of galaxy orientation angles (Pimblet 2005), this problem now becomes vectorized (a so-called ‘connect-the-darts’ problem) and potentially easier to solve (Figure 6; Arias-Castro et al. 2004). We know from early work by Binggeli (1982) that the major axis of galaxy clusters are generally aligned exceptionally well with their first-ranked (usually a cD-type) galaxy and that close cluster pairs generally point to each other. Moreover in Λ CDM, filamentary structure funnels material along preferred directions toward clusters. Since galaxy alignment tends to follow the orientation of clusters and the filaments that feed them, Pimblet (2005; and references therein) took advantage of this fact by computing the degree of galaxy angle anisotropy for a selected region of the sky. Regions that have significantly anisotropic distributions of galaxy angles (as in Figure 6) readily show up.

Since it is easy to think of galaxies as points within large-scale surveys, it is no surprise that one can apply many mathematical approaches to delineating FOGs and other intrinsic patterns within them. A popular approach is to use a minimal spanning tree formalism (MST herein; e.g. Barrow, Bhavsar, & Sonoda 1985; Bhavsar & Ling 1988; Krzewina & Saslaw 1996; Doroshkevich et al. 2000;

³ Random numbers are generated using the method of Pimblet & Bulmer (2005).




Figure 5 Top left: a simulated toy-model FOG as might be found in a redshift survey; generated as randomly sampled points along a vertical (thick) line. Bottom left: a smoothed distribution of this FOG. An isodensity contour cut could readily be employed to delineate it. Top right: adding in random noise results in the original signal becoming harder to pick out — a variant of the so-called connect the dots type of problem. Bottom right: without the noise, the isodensity contours are approximately straight, but with noise, they start to look much more curvy. Further, it may also appear that at lower densities there is a horizontal FOG passing through the overdensity in the lower portion of the plot; but it should still be possible to detect the original FOG by thresholding at a particular isodensity contour.




Figure 6 If one makes use of galaxy angles (i.e. vectors), then the problem posed by Figure 5 potentially becomes much easier to solve. Here, the members of the FOG have orientations of $\theta = 90 \pm 15$ deg, whereas the interlopers (red) have purely random orientation angles.

Doroshkevich et al. 2004 amongst others). One drawback of MST is that it will produce a unique graph for a given set of points. It is known that large-scale surveys such as SDSS and 2dFGRS are incomplete by about 10 to 20% at all magnitudes (Cross et al. 2004; Pimblet et al. 2001) — a fact that makes MST potentially a poorer choice for analysing large-scale structure with than other methods.

Such other statistical methods, that here we will mention only briefly, include the use of Voronoi (and the complementary Delaunay) tessellations (e.g. van de Weygaert 1994). Essentially, the Voronoi tessellation can be thought of as constructing a skeleton of the Universe by simply finding the bisection line between a single point and every other point. This process is repeated for each point and then the Voronoi tessellation is then the unification of all the halfplanes that have been created (see van de Weygaert 1994 for more indepth detail). Analysis of the cell-like structures of the Voronoi skeleton can inform one about the underlying galaxy distribution, although direct detection of FOGs from these tessellations remains a relatively unattempted task. Minkowski functionals such as SHAPEFINDERS (e.g. Bharadwaj et al.

2000; Pandey & Bharadwaj 2005; see also Shandarin, Sheth, & Sahni 2004) and the genus statistics (e.g. Hoyle et al. 2002; Hoyle, Vogeley, & Gott 2002) can also provide us with a direct way of analysing the structure of the galaxy distribution. Moreover, they can also provide a direct measure of the ‘filamentarity’ and ‘planarity’ of the Universe (e.g. Schmalzing et al. 1999) and one can readily delineate FOGs from them by using an isodensity contour cut. Finally, we should also mention that there are a host of other marked point processes (Stoica et al. 2005 and references therein) which are also capable of recovering individual FOGs. All of these methods highlight the presence of FOGs within redshift surveys to varying degrees.

3 Discussion and Conclusions

Given the above methods to detect FOGs, there is a large amount of literature dedicated to their dissemination. One issue that seems to be prominent in the literature is the mixed nomenclature for FOGs. Many authors refer to them as ‘walls’ (Geller & Huchra 1989), others call them ‘filaments’, some use the term ‘sheets’ or ‘pancakes’. So what is the difference between all these terms? Pimblet, Drinkwater, & Hawkrigg (2004) and Colberg, Krughoff, & Connolly (2005) attempt to refine these definitions by dividing detected FOGs into several categories based upon their visual morphology. So the difference between a filament and a wall then becomes a matter of how thick (or, equally, how wide) the FOG is in three-dimensional space (i.e. a filament will have depth \approx width). Sheets are then synonymous with walls. Is this kind of morphological classification useful? Given that walls appear to be much, much rarer than ‘normal’ filaments (Pimblet, Drinkwater, & Hawkrigg 2004) and unlike filaments, they do not possess non-isotropic galaxy orientations (Pimblet 2005) — yes. Their relative abundances (also filling factors) and lengths should help us to better constrain the ideal cosmological paradigm (e.g. in Λ CDM cosmologies, studies of FOGs can readily exclude bias parameters of $b > 1.5$; Bharadwaj & Pandey 2004) as can the number of FOGs connected to clusters of a given mass (Colberg, Krughoff, & Connolly 2005).

At the outset of this work, the question ‘what is a filament of Galaxies?’ was posed. This work has reviewed a number of methods for finding, detecting, and defining FOGs in datasets of varying complexity. Those that are more physically motivated (gravitational weak lensing searches; X-ray searches) appear to be an optimal way to detecting them (especially in unison), but yet, they remain a time-intensive method owing to the required amount of observing time to get down to sufficient limiting magnitudes and fluxes.

We have also investigated how FOGs are detected in large-redshift surveys using a variety of methods ranging from simple isodensity thresholding to more involved statistics like the MST. Here, it seems that even the very simple approaches can yield useful results, such as the

distribution and abundances of FOG lengths, that are in remarkable agreement with theory.

Acknowledgments

K.A.P. thanks Michael Drinkwater, Alastair Edge, and Mary Hawkrigg for their support. This work is financed through the award of a University of Queensland EPSA Research Fellowship and a UQRSF grant.

On a final note, I also wish to express my sincere gratitude to the two referees, Dominique Proust and Tony Fairall, who provided quick and useful reports that have improved the quality of this work.

References

- Abazajian, K., et al. 2004, *AJ*, 128, 502
 Abell, G. O. 1965, *ARA&A*, 3, 1
 Arias-Castro, E., Donoho, D., Huo, X., & Tovey, C. 2004, *Advances in Applied Probability*, submitted (www-stat.stanford.edu/~donoho/Reports/2004/CTD-Arias-et-al.pdf)
 Atrio-Barandela, F., & Kashlinsky, A. 1992, *ApJ*, 390, 322
 Barrow, J. D., Bhavsar, S. P., & Sonoda, D. H. 1985, *MNRAS*, 216, 17
 Bharadwaj, S., & Pandey, B. 2004, *ApJ*, 615, 1
 Bharadwaj, S., Sahni, V., Sathyaprakash, B. S., Shandarin, S. F., & Yess, C. 2000, *ApJ*, 528, 21
 Bhavsar, S. P., & Ling, E. N. 1988, *ApJ*, 331, L63
 Binggeli, B. 1982, *A&A*, 107, 338
 Bond, J. R., Kofman, L., & Pogosyan, D. 1996, *Natur*, 380, 603
 Bower, R. G., Lucey, J. R., & Ellis, R. S. 1992, *MNRAS*, 254, 601
 Briel, U. G., & Henry, J. P. 1995, *A&A*, 302, L9
 Butcher, H., & Oemler, A. 1984, *ApJ*, 285, 426
 Cen, R., & Ostriker, J. P. 1999, *ApJ*, 514, 1
 Chodorowski, M. 1994, *MNRAS*, 266, 897
 Clowe, D., Luppino, G. A., Kaiser, N., Henry, J. P., & Gioia, I. M. 1998, *ApJ*, 497, L61
 Colberg, J. M., Krughoff, K. S., & Connolly, A. J. 2005, *MNRAS*, 359, 272
 Colberg, J. M., et al. 2000, *MNRAS*, 319, 209
 Colberg, J. M., White, S. D. M., Jenkins, A., & Pearce, F. R. 1999, *MNRAS*, 308, 593
 Coleman, P. H., & Pietronero, L. 1992, *PhR*, 213, 311
 Cross, N. J. G., Driver, S. P., Liske, J., Lemon, D. J., Peacock, J. A., Cole, S., Norberg, P., & Sutherland, W. J. 2004, *MNRAS*, 349, 576
 Dietrich, J. P., Schneider, P., Clowe, D., Romano-Diaz, E., & Kerp, J. 2004 (astro-ph/0406541)
 Doroshkevich, A. G., Fong, R., McCracken, H. J., Ratcliffe, A., Shanks, T., & Turchaninov, V. I. 2000, *MNRAS*, 315, 767
 Doroshkevich, A., Tucker, D. L., Allam, S., & Way, M. J. 2004, *A&A*, 418, 7
 Durret, F., Lima Neto, G. B., Forman, W., & Churazov, E. 2003, *A&A*, 403, L29
 Ebeling, H., Barrett, E., & Donovan, D. 2004, *ApJ*, 609, L49
 Ebeling, H., Voges, W., Bohringer, H., Edge, A. C., Huchra, J. P., & Briel, U. G. 1996, *MNRAS*, 281, 799
 Fairall, A., Turner, D., Pretorius, M. L., Wiehahn, M., McBride, V., de Vaux, G., & Woudt, P. A. 2004 (astro-ph/0411437)
 Fukugita, M., Hogan, C. J., & Peebles, P. J. E. 1998, *ApJ*, 503, 518
 Gavazzi, R., Mellier, Y., Fort, B., Cuillandre, J.-C., & Dantel-Fort, M. 2004, *A&A*, 422, 407
 Geller, M. J., & Huchra, J. P. 1989, *Sci*, 246, 897
 Gray, M. E., Taylor, A. N., Meisenheimer, K., Dye, S., Wolf, C., & Thommes, E. 2002, *ApJ*, 568, 141
 Hawkins, E., et al. 2003, *MNRAS*, 346, 78
 Hoyle, F., et al. 2002, *ApJ*, 580, 663
 Hoyle, F., Vogeley, M. S., & Gott, J. R. I. 2002, *ApJ*, 570, 44

- Huchra, J., Davis, M., Latham, D., & Tonry, J. 1983, *ApJS*, 52, 89
- Jenkins, A., et al. 1998, *ApJ*, 499, 20
- Jones, D. H., et al. 2004, *MNRAS*, 355, 747
- Kaiser, N., Wilson, G., Luppino, G., Kofman, L., Gioia, I., Metzger, M., & Dahle, H. 1998 (astro-ph/9809268)
- Katz, N., Weinberg, D. H., Hernquist, L., & Miralda-Escude, J. 1996, *ApJ*, 457, L57
- Kodama, T., Smail, I., Nakata, F., Okamura, S., & Bower, R. G. 2001, *ApJ*, 562, L9
- Krzewina, L. G., & Saslaw, W. C. 1996, *MNRAS*, 278, 869
- Kull, A., & Böhringer, H. 1999, *A&A*, 341, 23
- Pandey, B., & Bharadwaj, S. 2005, *MNRAS*, 357, 1068
- Paolillo, M., Andreon, S., Longo, G., Puddu, E., Gal, R. R., Scaramella, R., Djorgovski, S. G., & de Carvalho, R. 2001, *A&A*, 367, 59
- Pimbblet, K. A. 2003, *PASA*, 20, 294
- Pimbblet, K. A. 2005, *MNRAS*, 358, 256
- Pimbblet, K. A., & Bulmer, M. 2005, *PASA*, 22, 1
- Pimbblet, K. A., & Drinkwater, M. J. 2004, *MNRAS*, 347, 137
- Pimbblet, K. A., Drinkwater, M. J., & Hawkrigg, M. C. 2004, *MNRAS*, 354, L61
- Pimbblet, K. A., Edge, A. C., & Couch, W. J. 2005, *MNRAS*, 357, L45
- Pimbblet, K. A., Smail, I., Kodama, T., Couch, W. J., Edge, A. C., Zabludoff, A. I., & O'Hely, E. 2002, *MNRAS*, 331, 333
- Pimbblet, K. A., Smail, I., Edge, A. C., Couch, W. J., O'Hely, E., & Zabludoff, A. I. 2001, *MNRAS*, 327, 588
- Pogosyan, D., Bond, J. R., Kofman, L., & Wadsley, J. 1998, in *Wide Field Surveys in Cosmology*, 14th IAP Meet., 1998 (Paris: Editions Frontieres), 61
- Scharf, C., Donahue, M., Voit, G. M., Rosati, P., & Postman, M. 2000, *ApJ*, 528, L73
- Schmalzing, J., Buchert, T., Melott, A. L., Sahni, V., Sathyaprakash, B. S., & Shandarin, S. F. 1999, *ApJ*, 526, 568
- Shandarin, S. F., Sheth, J. V., & Sahni, V. 2004, *MNRAS*, 353, 162
- Shectman, S. A., Landy, S. D., Oemler, A., Tucker, D. L., Lin, H., Kirshner, R. P., & Schechter, P. L. 1996, *ApJ*, 470, 172
- Stoica, R. S., Martinez, V. J., Mateu, J., & Saar, E. 2005, *A&A*, 434, 423
- Tittley, E. R., & Henriksen, M. 2001, *ApJ*, 563, 673
- van de Weygaert, R. 1994, *A&A*, 283, 361
- Visvanathan, N., & Sandage, A. 1977, *ApJ*, 216, 214
- Zeldovich, I. B., Einasto, J., & Shandarin, S. F. 1982, *Natur*, 300, 407