INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, http://www.editorialmanager.com/psm/. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

- 1. S.I. units should be used throughout in text, figures and tables.
- 2. Authors should spell out in full any abbreviations used in their manuscripts.
- 3. Foreign quotations and phrases should be followed by a translation.
- 4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* 286, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* 32, 173–179.

Cleckley HJ (1941). The Mask of Sanity, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank. org/INTPAH/ Resources/Publications/Quantitative-Techniques/health.eq tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. Line artwork: Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; Combination artwork (line/tone): Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; Black and white halftone artwork: Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; Colour halftone artwork: Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. Tables Tables should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

REVIEW ARTICLE		Reciprocal effects of stable and temporary components of	
Response, remission and drop-out rates following		neuroticism and affective disorders: results of a longitudinal	
high-frequency repetitive transcranial magnetic stimulation		cohort study	
(rTMS) for treating major depression: a systematic review and		Spinhoven P, Penelo E, de Rooij M, Penninx BW & Ormel J	337
meta-analysis of randomized, double-blind and		Do positive or negative stressful events predict the development	t
sham-controlled trials		of new brain lesions in people with multiple sclerosis?	
Berlim MT, van den Eynde F, Tovar-Perdomo S & Daskalakis ZJ	225	Burns MN, Nawacki E, Kwasny MJ, Pelletier D & Mohr DC	349
ORIGINAL ARTICLES		Latent structure of social fears and social anxiety disorders	
Paradoxical effects of short-term antidepressant treatment in fMRI emotional processing models in volunteers with high neuroticism		Iza M, Wall MM, Heimberg RG, Rodebaugh TL, Schneier FR, Liu S-M & Blanco C	361
Di Simplicio M, Norbury R, Reinecke A & Harmer CJ	241	Environmental influences on familial resemblance for drug	
Peer problems are associated with elevated serum leptin levels in children		abuse in first-cousin pairs: a Swedish national study Kendler KS, Ohlsson H, Sundquist K & Sundquist J	371
Kohlboeck G, Romanos M, Tiesler C, Koletzko S, Kratzsch J,		Altered top-down and bottom-up processing of fear	
Thiery J, Bauer C-P, von Berg A, Berdel D, Hoffmann B,		conditioning in panic disorder with agoraphobia	
Schaaf B, Lehmann I, Herbarth O, Heinrich J, GINI-plus &	055	Lueken U, Straube B, Reinhardt I, Maslowski NI, Wittchen H-U,	
LISA-plus study groups	255	Ströhle A, Wittmann A, Pfleiderer B, Konrad C, Ewert A,	
Individual- and area-level influence on suicide risk: a		Uhlmann C, Arolt V, Jansen A & Kircher T	381
multilevel longitudinal study of Swedish schoolchildren		Occupitation of the section of the s	
Zammit S, Gunnell D, Lewis G, Leckie G, Dalman C & Allebeck P	267	Cognitive subtypes in non-affected siblings of schizophrenia	
		patients: characteristics and profile congruency with affected family members	
Sociodemographic, psychiatric, and somatic risk factors for		Quee PJ, Alizadeh BZ, Aleman A, van den Heuvel & GROUP	
suicide: a Swedish national cohort study		Investigators	395
Crump C, Sundquist K, Sundquist J & Winkleby MA	279	livestigators	393
Incidence and predictors of suicide attempts among		Modelling the interplay between childhood and adult	
primary-care patients with depressive disorders: a 5-year		adversity in pathways to psychosis: initial evidence from the	
prospective study		AESOP Study	
Riihimäki K, Vuorilehto M, Melartin T, Haukka J & Isometsä E	291	Morgan C, Reininghaus U, Fearon P, Hutchinson G, Morgan K,	
	20.	Dazzan P, Boydell J, Kirkbride JB, Doody GA, Jones PB,	
Causal beliefs of the public and social acceptance of persons		Murray RM & Craig T	407
with mental illness: a comparative analysis of schizophrenia,		Sexual minority status and psychotic symptoms: findings	
depression and alcohol dependence		from the Netherlands Mental Health Survey and Incidence	
Schomerus G, Matschinger H & Angermeyer MC	303	Studies (NEMESIS)	
Lifetime prevalence and co-morbidity of externalizing		Gevonden MJ, Selten JP, Myin-Germeys I, de Graaf R,	
disorders and depression in prospective assessment		ten Have M, van Dorsselaer S, van Os J & Veling W	421
Hamdi NR & Jacono WG	315		
		Rapid visual information processing as a cognitive	
Subclinical psychotic experiences and bipolar		endophenotype for attention deficit hyperactivity disorder	405
spectrum features in depression: association with		Gau SS-F & Huang W-L	435
outcome of psychotherapy		Correspondence	447
Wigman JTW, van Os J, Abidi L, Huibers MJH, Roelofs J,	005	Out the state of the	050
Arntz A Kelleher I & Peeters FPMI	325	Corrigendum	253





