

KRASNOSELSKI–MANN ITERATION FOR HIERARCHICAL FIXED POINTS AND EQUILIBRIUM PROBLEM

GIUSEPPE MARINO[✉], VITTORIO COLAO, LUIGI MUGLIA
and YONGHONG YAO

(Received 30 April 2008)

Abstract

We give an explicit Krasnoselski–Mann type method for finding common solutions of the following system of equilibrium and hierarchical fixed points:

$$\begin{cases} G(x^*, y) \geq 0, & \forall y \in C, \\ \text{find } x^* \in \text{Fix}(T) \text{ such that } \langle x^* - f(x^*), x - x^* \rangle \geq 0, & \forall x \in \text{Fix}(T), \end{cases}$$

where C is a closed convex subset of a Hilbert space H , $G : C \times C \rightarrow \mathbb{R}$ is an equilibrium function, $T : C \rightarrow C$ is a nonexpansive mapping with $\text{Fix}(T)$ its set of fixed points and $f : C \rightarrow C$ is a ρ -contraction. Our algorithm is constructed and proved using the idea of the paper of [Y. Yao and Y.-C. Liou, ‘Weak and strong convergence of Krasnosel’skiĭ–Mann iteration for hierarchical fixed point problems’, *Inverse Problems* 24 (2008), 501–508], in which only the variational inequality problem of finding hierarchically a fixed point of a nonexpansive mapping T with respect to a ρ -contraction f was considered. The paper follows the lines of research of corresponding results of Moudafi and Théra.

2000 *Mathematics subject classification*: primary 47H09, 47H10; secondary 58E35.

Keywords and phrases: hierarchical fixed points, equilibrium problem, nonexpansive map, contraction, variational inequality problem, projection.

1. Introduction

Let T, V be two nonexpansive mappings from C to C , where C is a closed and convex subset of a Hilbert space H . Consider the variational inequality problem (VIP) of finding hierarchically a fixed point of a nonexpansive mapping T with respect to another nonexpansive mapping V , that is,

$$\text{find } x^* \in \text{Fix}(T) \text{ such that } \langle x^* - Vx^*, y - x^* \rangle \geq 0 \quad y \in \text{Fix}(T). \quad (1.1)$$

(Equivalently, $x^* = P_{\text{Fix}(T)} Vx^*$ – that is, x^* is a fixed point of the nonexpansive map $P_{\text{Fix}(T)} V$ – where for K closed convex subset of H , P_K is the metric projection of H on K).

Supported by Ministero dell’Università e della Ricerca of Italy.

© 2009 Australian Mathematical Society 0004-9727/2009 \$16.00

Of course if $V = I$, the solution set S of (1.1) is just $\text{Fix}(T)$.

The VIP (1.1) covers several topics investigated in literature, among them the following:

(1) (Monotone inclusions) Yamada [32] studies the VIP (1.1) assuming $V = I - \gamma F$, where $\gamma > 0$ is sufficiently small and the operator F is Lipschitzian and strongly monotone.

(2) (Convex optimization [4, 23]) Let φ be a proper lower semicontinuous convex function on H and let ψ be a convex function on H so that $\nabla\psi$ is strongly monotone. Take

$$T = \text{prox}_{\lambda\varphi} := \operatorname{argmin} \left\{ \varphi(z) + \frac{1}{2\lambda} \|\cdot - z\|^2 \right\}.$$

Then the VIP (1.1) reduces to the hierarchical minimization problem

$$\min_{x \in \operatorname{argmin} \varphi} \psi(x).$$

(3) (Quadratic minimizations over a fixed point set [14]) If A is a linear bounded strongly positive operator on H , f is a ρ -contraction on H and h is a potential for γf (that is, $h'(x) = \gamma f(x)$) where $\gamma > 0$ is a constant, consider the minimization problem

$$\min_{x \in \text{Fix}(T)} \frac{1}{2} \langle Ax, x \rangle - h(x). \tag{1.2}$$

The optimality condition to minimize (1.2) is to find a fixed point of T so that

$$\langle (A - \gamma f)x^*, x - x^* \rangle \geq 0, \quad x \in \text{Fix}(T).$$

Taking $V = I - \lambda(A - \gamma f)$, where $\gamma > 0$ is appropriately chosen so that V is nonexpansive, we find that the previous VIP reduced to (1.1).

(4) Let A be a maximal monotone operator. Take $T = J_\lambda^A := (I + \lambda A)^{-1}$ and $V = I - \gamma \nabla\psi$, where ψ is a convex function such that $\nabla\psi$ is η -Lipschitzian (which is equivalent to the fact that $\nabla\psi$ is η^{-1} co-coercive), with $\gamma \in (0, 2/\eta]$ and $\text{Fix}(J_\lambda^A) = A^{-1}(0)$. So VIP (1.1) reduces to the mathematical program with generalized equation constraint,

$$\min_{0 \in A(x)} \psi(x),$$

considered in [13].

A very particular case of the VIP (1.1) occurs when V is a constant mapping, that is, given $u \in H$,

$$\text{find } x^* \in \text{Fix}(T) \text{ such that } \langle x^* - u, x - x^* \rangle \geq 0, \quad x \in \text{Fix}(T), \tag{1.3}$$

or, equivalently, find the fixed point of T closest to u , that is,

$$x^* = P_{\text{Fix}(T)}u = \operatorname{argmin}_{x \in \text{Fix}(T)} \frac{1}{2} \|u - x\|^2.$$

This problem was widely investigated in [2, 9, 12, 22, 26, 28, 29]. The explicit method, initiated by Halpern in [9], generates a sequence $(x_n)_n$ by iterating

$$x_{n+1} = \alpha_n u + (1 - \alpha_n)Tx_n, \quad (1.4)$$

where $u, x_0 \in C$ and $(\alpha_n)_n \subset [0, 1]$.

The next result is well known.

THEOREM 1.1 [2, 9, 20, 21, 25–27]. *Assume that $\text{Fix}(T)$ is nonempty. Suppose that the sequence $(\alpha_n)_n$ satisfies the following:*

- (1) $\lim_n \alpha_n = 0$;
- (2) $\sum_n \alpha_n = \infty$;
- (3) $\sum_n |\alpha_{n+1} - \alpha_n| < \infty$ or $\lim_n ((\alpha_{n+1} - \alpha_n)/\alpha_n) = 0$.

Then the sequence $(x_n)_n$ generated by the algorithm (1.4) converges in norm to $P_{\text{Fix}(T)}u$.

A more general case than V constant is that one $V = f$ with f is a ρ -contraction, that is, $\|f(x) - f(y)\| \leq \rho\|x - y\|$, $\rho \in (0, 1)$. In this case we call (1.1) the contractive VIP and the method is also known as viscosity approximation. It was first studied by Moudafi [15] and further developed by Xu [30].

In this method, the explicit scheme (1.4) is replaced by Mann-type scheme

$$x_{n+1} = \lambda_n f(x_n) + (1 - \lambda_n)Tx_n \quad (1.5)$$

where $(\lambda_n)_n$ is a sequence in $[0, 1]$.

THEOREM 1.2 [15, 30]. *Assume that $\text{Fix}(T)$ is nonempty and let $(x_n)_n$ be the sequence by the algorithm (1.5). Assume that:*

- (1) $\lim_n \lambda_n = 0$;
- (2) $\sum_n \lambda_n = \infty$;
- (3) $\sum_n |\lambda_{n+1} - \lambda_n| < \infty$ or $\lim_n ((\lambda_{n+1} - \lambda_n)/\lambda_n) = 0$.

Then $\lim_n x_n = x^$ exists and x^* is the unique solution of the variational inequality*

$$\langle (I - f)x^*, x - x^* \rangle \geq 0, \quad x \in \text{Fix}(T).$$

Very recently, Yao and Liou [35] replaced the Mann-type scheme (1.5) with the Krasnoselski–Mann type scheme

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n(\lambda_n f(x_n) + (1 - \lambda_n)Tx_n)$$

and proved the following theorem.

THEOREM 1.3 [35]. *Let C be a nonempty closed convex subset of a real Hilbert space H . Let T be a nonexpansive mapping of C into itself such that $\text{Fix}(T) \neq \emptyset$. Let $P : C \rightarrow C$ be a ρ -contraction. Let $(x_n)_n$ be a sequence generated by*

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n(\sigma_n Px_n + (1 - \sigma_n)Tx_n), \quad n \geq 0.$$

Let $(\alpha_n)_n, (\sigma_n)_n$ be two real number sequences in $(0, 1)$ satisfying the following conditions:

- (i) $0 < \liminf_{n \rightarrow \infty} \alpha_n \leq \limsup_{n \rightarrow \infty} \alpha_n < 1$;
- (ii) $\lim_{n \rightarrow \infty} \sigma_n = 0$ and $\sum_n \sigma_n = \infty$.

Then:

- (1) $(x_n)_n$ converges strongly to a fixed point of T ;
- (2) $(x_n)_n$ is asymptotically regular, namely $\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0$;
- (3) $(x_n)_n$ converges strongly to a solution of the problem

$$\text{find } x^* \in \text{Fix}(T) \text{ such that } \langle x^* - f(x^*), x - x^* \rangle \geq 0, \quad \forall x \in \text{Fix}(T).$$

The above scheme is a particular case of the Krasnoselski–Mann algorithm

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n(\lambda_n V(x_n) + (1 - \lambda_n)Tx_n)$$

with V a nonexpansive mapping, introduced by Moudafi [17].

Some algorithms in signal processing and image reconstruction may be written as the well-known Krasnoselski–Mann (K–M) iteration. The main feature of (K–M)-iteration convergence theorems provided a unified framework for analyzing various concrete algorithms. For details, see [3, 5, 31–34].

On the other hand, note that if we put $C = \text{Fix}(T)$ and $G(x, y) := \langle (I - V)x, y - x \rangle$, then the VIP (1.1) can be rewritten as

$$\text{find } x^* \in C \text{ such that } G(x^*, y) \geq 0, \quad y \in C, \tag{1.6}$$

that is, as an equilibrium problem. More generally, following [6], we can have a countable family of bifunctions from $C \times C$ to \mathbb{R} . The basic formulation of this class of problems reduces to solving the system of equilibrium problems

$$\text{find } x \in C \text{ such that } G_i(x, y) \geq 0, \quad \forall i \in I, \forall y \in C. \tag{1.7}$$

Blum and Oettli [1, 19] show that, in the case of a single equilibrium problem, the formulation (1.6) covers monotone inclusion problems, saddlepoint problems, VIPs, minimization problems, Nash equilibria in noncooperative games, vector equilibrium problems and certain fixed point problems (see [8]).

It is also worth remarking that, in the case of VIP (1.1), the induced bifunction $G(x, y) := \langle (I - V)x, y - x \rangle$ satisfies the following condition.

CONDITION (1).

- (E1) $G(x, x) = 0$ for all $x \in H$.
- (E2) $G(x, y) + G(y, x) \leq 0$ for all $(x, y) \in H \times H$ (that is, G is monotone).
- (E3) For each $x, y, z \in H$,

$$\limsup_{t \rightarrow 0} G(tz + (1 - t)x, y) \leq G(x, y).$$

- (E4) The function $y \rightarrow G(x, y)$ is convex and lower semicontinuous for each $x \in H$.

While many methods have been proposed to solve (1.6) (see [7, 10, 11, 16, 18]), we are not aware of so many results for systems of equilibrium problems. For some partial results on these topics see [6].

Here we study a particular case of a system of two equilibrium functions, one induced by a contractive VIP and one satisfying Condition (1), namely

$$\begin{cases} G(x^*, y) \geq 0, & \forall y \in C, \\ \text{find } x^* \in \text{Fix}(T) \text{ such that } \langle x^* - f(x^*), x - x^* \rangle \geq 0, & \forall x \in \text{Fix}(T). \end{cases} \quad (1.8)$$

Of course such systems include the systems given by a VIP and a contractive VIP.

We show that the following Krasnoselski–Mann-type scheme for the VIP and equilibrium function

$$\begin{cases} x_0 \in C, \\ G(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, & \forall y \in C, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n(\lambda_n f(x_n) + (1 - \lambda_n)Tu_n), & n \geq 1, \end{cases} \quad (1.9)$$

solves the system.

2. Preliminaries

We give several known results that are fundamental for our proof.

LEMMA 2.1 [24]. *Let $(x_n)_{n \in \mathbb{N}}$ and $(z_n)_{n \in \mathbb{N}}$ be bounded sequences in a Banach space X and let $(\beta_n)_{n \in \mathbb{N}}$ be a sequence in $[0, 1]$ with $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$. Suppose that*

$$x_{n+1} = \beta_n x_n + (1 - \beta_n)z_n,$$

for all integers $n \geq 0$, and

$$\limsup_{n \rightarrow \infty} (\|z_{n+1} - z_n\| - \|x_{n+1} - x_n\|) \leq 0.$$

Then $\lim_{n \rightarrow \infty} \|x_n - z_n\| = 0$.

LEMMA 2.2 [29]. *Assume $(a_n)_n$ is a sequence of nonnegative numbers such that*

$$a_{n+1} \leq (1 - \gamma_n)a_n + \delta_n, \quad n \geq 0,$$

where $(\gamma_n)_n$ is a sequence in $(0, 1)$ and $(\delta_n)_n$ is a sequence in \mathbb{R} such that:

- (1) $\sum_{n=1}^{\infty} \gamma_n = \infty$;
- (2) $\limsup_{n \rightarrow \infty} \delta_n / \gamma_n \leq 0$ or $\sum_{n=1}^{\infty} |\delta_n| < \infty$.

Then $\lim_{n \rightarrow \infty} a_n = 0$.

The next lemmas concern the equilibrium function G and the set of equilibrium points

$$EP(G) = \{x \in C \mid G(x, y) \geq 0, \forall y \in C\}.$$

LEMMA 2.3 [6]. *Let C be a nonempty closed convex subset of H and $G : C \times C \rightarrow \mathbb{R}$ satisfy Condition (1). For $x \in C$ and $r > 0$, let $S_r : H \rightarrow C$ be the r -resolvent of G ,*

$$S_r(x) := \left\{ z \in C \mid G(z, y) + \frac{1}{r} \langle y - z, z - x \rangle \geq 0, \forall y \in C \right\}.$$

Then S_r is well defined and the following hold:

- (1) S_r is single-valued;
- (2) S_r is firmly nonexpansive, that is,

$$\|S_r x - S_r y\|^2 \leq \langle S_r x - S_r y, x - y \rangle,$$

for all $x, y \in H$;

- (3) $\text{Fix}(S_r) = EP(G)$;
- (4) $EP(G)$ is closed and convex.

LEMMA 2.4 [6]. *Suppose that $G : C \times C \rightarrow \mathbb{R}$ is an equilibrium function satisfying Condition (1). Let $(x_n)_{n \in \mathbb{N}}$ be a sequence in H and $(r_n)_{n \in \mathbb{N}}$ a sequence in $(0, +\infty)$. Define, for all $n \in \mathbb{N}$, $u_n := S_{r_n} x_n$ and suppose that $u_n \rightarrow p$ and $(x_n - u_n) \rightarrow z$. Then $p \in C$ and for all $y \in C$, $G(p, y) + \langle z, p - y \rangle \geq 0$.*

REMARK 2.5. Note that in Lemma 2.4, if $z = 0$, then the weak cluster point p for $(u_n)_{n \in \mathbb{N}}$ is a weak cluster point for $(x_n)_{n \in \mathbb{N}}$ and also an equilibrium point for G .

LEMMA 2.6. *Let $G : C \times C \rightarrow \mathbb{R}$ be a bifunction such that Condition (1) holds. Let $(w_n)_n$ be a bounded sequence and $z_n := S_{r_n} w_n$. Let $(r_n)_n$ be a sequence of positive numbers such that $\liminf_n r_n = r > 0$. Then there exists a constant $L > 0$ such that*

$$\|z_{n+1} - z_n\| \leq \|w_{n+1} - w_n\| + L \left| 1 - \frac{r_n}{r_{n+1}} \right|. \tag{2.1}$$

PROOF. Since $z_n := S_{r_n} w_n$ and $z_{n+1} := S_{r_{n+1}} w_{n+1}$, we obtain that

$$G(z_{n+1}, y) + \frac{1}{r_{n+1}} \langle y - z_{n+1}, z_{n+1} - w_{n+1} \rangle \geq 0, \quad \forall y \in C,$$

and

$$G(z_n, y) + \frac{1}{r_n} \langle y - z_n, z_n - w_n \rangle \geq 0, \quad \forall y \in C.$$

In particular,

$$G(z_{n+1}, z_n) + \frac{1}{r_{n+1}} \langle z_n - z_{n+1}, z_{n+1} - w_{n+1} \rangle \geq 0$$

and

$$G(z_n, z_{n+1}) + \frac{1}{r_n} \langle z_{n+1} - z_n, z_n - w_n \rangle \geq 0.$$

Hence, summing up these two inequalities and using (E2),

$$\frac{1}{r_n} \langle z_{n+1} - z_n, z_n - w_n \rangle + \frac{1}{r_{n+1}} \langle z_n - z_{n+1}, z_{n+1} - w_{n+1} \rangle \geq 0,$$

so it follows that

$$\left\langle z_{n+1} - z_n, \frac{z_n - w_n}{r_n} - \frac{z_{n+1} - w_{n+1}}{r_{n+1}} \right\rangle \geq 0. \tag{2.2}$$

We derive from (2.2) that

$$\begin{aligned} & \left\langle z_{n+1} - z_n, z_n - w_n - \frac{r_n}{r_{n+1}}(z_{n+1} - w_{n+1}) \right\rangle \geq 0 \\ & \Rightarrow \left\langle z_{n+1} - z_n, z_n - z_{n+1} - w_n + z_{n+1} - \frac{r_n}{r_{n+1}}(z_{n+1} - w_{n+1}) \right\rangle \geq 0 \\ & \Rightarrow -\|z_{n+1} - z_n\|^2 + \left\langle z_{n+1} - z_n, (z_{n+1} - w_{n+1}) \left(1 - \frac{r_n}{r_{n+1}}\right) \right. \\ & \quad \left. + (w_{n+1} - w_n) \right\rangle \geq 0. \end{aligned}$$

Then

$$\begin{aligned} \|z_{n+1} - z_n\|^2 & \leq \left\langle z_{n+1} - z_n, (z_{n+1} - w_{n+1}) \left(1 - \frac{r_n}{r_{n+1}}\right) + (w_{n+1} - w_n) \right\rangle \\ & \leq \|z_{n+1} - z_n\| \left(\|w_{n+1} - w_n\| + \left|1 - \frac{r_n}{r_{n+1}}\right| \|z_{n+1} - w_{n+1}\| \right), \end{aligned}$$

and so

$$\|z_{n+1} - z_n\| \leq \|w_{n+1} - w_n\| + \left|1 - \frac{r_n}{r_{n+1}}\right| \|z_{n+1} - w_{n+1}\|.$$

By hypothesis on $(r_n)_n$, if $L := \sup_n \|z_{n+1} - w_{n+1}\|$, we conclude that

$$\|z_{n+1} - z_n\| \leq \|w_{n+1} - w_n\| + L \left|1 - \frac{r_n}{r_{n+1}}\right|. \quad \square$$

3. Main result

THEOREM 3.1. *Let C be a closed convex subset of a Hilbert space H . Let $T : C \rightarrow C$ be a nonexpansive mapping with $\text{Fix}(T) \cap EP(G) \neq \emptyset$. Let $f : C \rightarrow C$ be a ρ -contraction. Let $(\lambda_n)_n$ be a sequence in $(0, 1)$ such that $\lambda_n \rightarrow 0$ and $\sum_n \lambda_n = \infty$. Let $(\alpha_n)_n$ be a sequence in $(0, 1)$ such that $0 < \liminf_n \alpha_n \leq \limsup_n \alpha_n < 1$.*

Let $(r_n)_n$ be a sequence of positive real numbers such that $\liminf_n r_n = r > 0$ and $\lim_n |1 - ((r_n)/(r_{n+1}))| = 0$. Let $(x_n)_n, (u_n)_n$ be the sequences defined by

$$\begin{cases} x_0 \in C, \\ G(u_n, y) + \frac{1}{r_n} \langle y - u_n, u_n - x_n \rangle \geq 0, & \forall y \in C, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n(\lambda_n f(x_n) + (1 - \lambda_n)Tu_n), & n \geq 1. \end{cases} \quad (3.1)$$

Then the sequences both converge to a point $z \in \text{Fix}(T) \cap EP(G)$ which is the unique solution in $\text{Fix}(T) \cap EP(G)$ of the variational inequality

$$\langle z - f(z), z - x \rangle \leq 0, \quad \forall x \in \text{Fix}(T) \cap EP(G). \quad (3.2)$$

Equivalently, $z = P_{\text{Fix}(T) \cap EP(G)} f z$.

PROOF. Since the inequality

$$\|u_n - z\| = \|S_{r_n}x_n - S_{r_n}z\| \leq \|x_n - z\|$$

holds, we only prove that $x_n \rightarrow z$. We divide the proof into several steps.

STEP 1. We prove that the sequence $(x_n)_n$ is bounded. Let $v \in \text{Fix}(T) \cap EP(G)$. Then

$$\begin{aligned} \|x_{n+1} - v\| &= \|(1 - \alpha_n)(x_n - v) + \alpha_n[\lambda_n(f(x_n) - v) + (1 - \lambda_n)(Tu_n - v)]\| \\ &\leq (1 - \alpha_n)\|x_n - v\| + \alpha_n[\lambda_n(\|f(x_n) - f(v)\| \\ &\quad + \|f(v) - v\|) + (1 - \lambda_n)\|x_n - v\|] \\ &\leq (1 - \alpha_n)\|x_n - v\| + \alpha_n\lambda_n\rho\|x_n - v\| \\ &\quad + \alpha_n\lambda_n\|f(v) - v\| + \alpha_n(1 - \lambda_n)\|x_n - v\| \\ &= (1 - (1 - \rho)\lambda_n\alpha_n)\|x_n - v\| + \alpha_n\lambda_n\|f(v) - v\| \\ &\leq \max\left\{\|x_n - v\|, \frac{\|f(v) - v\|}{1 - \rho}\right\}. \end{aligned}$$

By induction we obtain that

$$\|x_n - v\| \leq \max\left\{\|x_0 - v\|, \frac{\|f(v) - v\|}{1 - \rho}\right\}.$$

STEP 2. We prove that the sequence $(x_n)_n$ is asymptotically regular, that is, $\|x_n - x_{n+1}\| \rightarrow 0$, as $n \rightarrow \infty$. Set $y_n = \lambda_n f(x_n) + (1 - \lambda_n)Tu_n$ and note that

$$\begin{aligned} y_{n+1} - y_n &= \lambda_{n+1}f(x_{n+1}) + (1 - \lambda_{n+1})Tu_{n+1} - \lambda_n f(x_n) - (1 - \lambda_n)Tu_n \\ &= \lambda_{n+1}(f(x_{n+1}) - f(x_n)) + (\lambda_{n+1} - \lambda_n)f(x_n) \\ &\quad + (1 - \lambda_{n+1})(Tu_{n+1} - Tu_n) - (\lambda_{n+1} - \lambda_n)Tu_n. \end{aligned}$$

So, to apply Lemma 2.1 (due to Suzuki), we observe that $x_{n+1} = \alpha_n x_n + (1 - \alpha_n)y_n$ and

$$\begin{aligned} & \limsup_n (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \\ & \leq \limsup_n [\lambda_{n+1} \|f(x_{n+1}) - f(x_n)\| + |\lambda_{n+1} - \lambda_n| \|f(x_n) - Tu_n\| \\ & \quad + (1 - \lambda_{n+1}) \|Tu_{n+1} - Tu_n\| - \|x_{n+1} - x_n\|] \\ & \leq \limsup_n [\lambda_{n+1} \rho \|x_{n+1} - x_n\| + |\lambda_{n+1} - \lambda_n| \|f(x_n) - Tu_n\| \\ & \quad + (1 - \lambda_{n+1}) \|u_{n+1} - u_n\| - \|x_{n+1} - x_n\|] \\ & \leq \limsup_n \left[\lambda_{n+1} \|x_{n+1} - x_n\| + (1 - \lambda_{n+1}) \left(\|x_{n+1} - x_n\| + L \left| 1 - \frac{r_n}{r_{n+1}} \right| \right) \right. \\ & \quad \left. + |\lambda_{n+1} - \lambda_n| (\|Tu_n\| + \|f(x_n)\|) - \|x_{n+1} - x_n\| \right], \end{aligned}$$

where the second inequality holds by (2.1) in Lemma 2.6. By the boundedness of $(x_n)_n$ and the hypotheses on the sequences $(\lambda_n)_n, (r_n)_n$ we conclude that

$$\begin{aligned} & \limsup_n (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \\ & \leq \limsup_n \left[\|x_{n+1} - x_n\| + L \left| 1 - \frac{r_n}{r_{n+1}} \right| \right. \\ & \quad \left. + |\lambda_{n+1} - \lambda_n| (\|Tu_n\| + \|f(x_n)\|) - \|x_{n+1} - x_n\| \right] \\ & = \limsup_n \left[L \left| 1 - \frac{r_n}{r_{n+1}} \right| + |\lambda_{n+1} - \lambda_n| (\|Tu_n\| + \|f(x_n)\|) \right] = 0. \end{aligned}$$

We can apply Lemma 2.1 to derive

$$\lim_n \|x_n - y_n\| = 0. \tag{3.3}$$

On the other hand, a straightforward computation leads to

$$\lim_n \|x_{n+1} - x_n\| = \lim_n \alpha_n \|x_n - y_n\| = 0. \tag{3.4}$$

STEP 3. We prove that $\lim_n \|x_n - u_n\| = 0$. First of all we note that, by the firm nonexpansivity of S_{r_n} , if $p \in EP(G)$, then

$$\begin{aligned} \|u_n - p\|^2 &= \langle u_n - p, S_{r_n}x_n - S_{r_n}p \rangle \leq \langle u_n - p, x_n - p \rangle \\ &= \frac{1}{2} (\|u_n - p\|^2 + \|x_n - p\|^2 - \|x_n - u_n\|^2), \end{aligned}$$

from which

$$\|u_n - p\|^2 \leq \|x_n - p\|^2 - \|x_n - u_n\|^2. \tag{3.5}$$

On the other hand, if $v \in \text{Fix}(T) \cap EP(G)$, then

$$\begin{aligned}
 \|x_{n+1} - v\|^2 &= \|(1 - \alpha_n)(x_n - v) + \alpha_n(\lambda_n f(x_n) + (1 - \lambda_n)Tu_n - v)\|^2 \\
 &= \|(1 - \alpha_n)(x_n - v) + \alpha_n(Tu_n - v) + \alpha_n\lambda_n(f(x_n) - Tu_n)\|^2 \\
 &\leq \|(1 - \alpha_n)(x_n - v) + \alpha_n(Tu_n - v)\|^2 \\
 &\quad + 2\lambda_n \langle f(x_n) - Tu_n, x_{n+1} - v \rangle \\
 &\leq (1 - \alpha_n)\|x_n - v\|^2 + \alpha_n\|Tu_n - Tv\|^2 \\
 &\quad + 2\lambda_n \langle f(x_n) - Tu_n, x_{n+1} - v \rangle \\
 &\leq (1 - \alpha_n)\|x_n - v\|^2 + \alpha_n\|u_n - v\|^2 \\
 &\quad + 2\lambda_n \langle f(x_n) - Tu_n, x_{n+1} - v \rangle.
 \end{aligned}
 \tag{3.6}$$

Combining (3.5) with (3.6) and setting

$$z_n = 2\lambda_n \langle f(x_n) - Tu_n, x_{n+1} - v \rangle$$

leads to

$$\begin{aligned}
 \|x_{n+1} - v\|^2 &\leq (1 - \alpha_n)\|x_n - v\|^2 + \alpha_n(\|x_n - v\|^2 - \|x_n - u_n\|^2) + z_n \\
 &\leq \|x_n - v\|^2 - \alpha_n\|x_n - u_n\|^2 + z_n.
 \end{aligned}
 \tag{3.7}$$

Thus,

$$\begin{aligned}
 \alpha_n\|x_n - u_n\|^2 &\leq \|x_n - v\|^2 - \|x_{n+1} - v\|^2 + z_n \\
 &\leq \|x_n - x_{n+1}\|^2 + 2\|x_n - x_{n+1}\|\|x_{n+1} - v\| + z_n.
 \end{aligned}
 \tag{3.8}$$

Since $(x_n)_n$ is bounded, $z_n \rightarrow 0$. Moreover, by asymptotically regularity of $(x_n)_n$ and by the hypothesis on $(\alpha_n)_n$, from the latter it follows that

$$\lim_n \|x_n - u_n\| = 0, \tag{3.9}$$

as required.

STEP 4. We now prove that the set of weak cluster points $\omega_w(x_n)$ is a subset of $\text{Fix}(T) \cap EP(G)$. Let $(x_{n_k})_k$ be a subsequence of $(x_n)_n$ weakly converging to a point $p \in C$. Since (3.9) holds, we can apply Lemma 2.4 to ensure that p lies in $EP(G)$.

To show that $p \in \text{Fix}(T)$, we observe that

$$\begin{aligned}
 \|x_{n_k} - Tx_{n_k}\| &\leq \|x_{n_k+1} - x_{n_k}\| + \|x_{n_k+1} - Tu_{n_k}\| + \|Tu_{n_k} - Tx_{n_k}\| \\
 &\leq \|x_{n_k+1} - x_{n_k}\| + (1 - \alpha_{n_k})\|x_{n_k} - Tu_{n_k}\| \\
 &\quad + \alpha_{n_k}\lambda_{n_k}\|f(x_{n_k}) - Tu_{n_k}\| + \|u_{n_k} - x_{n_k}\|,
 \end{aligned}$$

thus by hypotheses and by Steps 2 and 3,

$$\begin{aligned}
 \lim_k \|x_{n_k} - Tx_{n_k}\| &\leq \lim_k \frac{\|x_{n_k+1} - x_{n_k}\| + (2 - \alpha_{n_k})\|u_{n_k} - x_{n_k}\|}{\alpha_{n_k}} \\
 &\quad + \lambda_{n_k}\|f(x_{n_k}) - Tu_{n_k}\| = 0.
 \end{aligned}$$

Since $x_{n_k} \rightarrow p$, by the demiclosedness principle for nonexpansive mappings [2], we have $p \in \text{Fix}(T)$.

REMARK 3.2. Note that from Step 4 it follows that

$$\limsup_n \langle f(z) - z, x_n - z \rangle \leq 0, \tag{3.10}$$

where $z \in \text{Fix}(T) \cap EP(G)$ is the unique solution of the variational inequality (3.2). To show this, let $(x_{n_j})_j$ be such that

$$\limsup_n \langle f(z) - z, x_n - z \rangle = \lim_j \langle f(z) - z, x_{n_j} - z \rangle.$$

By eventually passing to subsequences, we may assume that $x_{n_j} \rightarrow p$. Then

$$\lim_j \langle f(z) - z, x_{n_j} - z \rangle = \langle f(z) - z, p - z \rangle \leq 0$$

since $p \in \text{Fix}(T) \cap EP(G)$.

STEP 5. Finally, we show that $x_n, u_n \rightarrow z$, as $n \rightarrow \infty$. Since the inequality

$$\|u_n - z\| = \|S_{r_n}x_n - S_{r_n}z\| \leq \|x_n - z\|$$

holds, it is enough to prove that $x_n \rightarrow z$:

$$\begin{aligned} \|x_{n+1} - z\|^2 &= \|(1 - \alpha_n)(x_n - z) + \alpha_n(\lambda_n f(x_n) + (1 - \lambda_n)Tu_n - z)\|^2 \\ &= \|(1 - \alpha_n)(x_n - z) + \alpha_n(1 - \lambda_n)(Tu_n - z) + \lambda_n\alpha_n(f(x_n) - z)\|^2 \\ &\leq \|(1 - \alpha_n)(x_n - z) + \alpha_n(1 - \lambda_n)(Tu_n - z)\|^2 \\ &\quad + 2\alpha_n\lambda_n \langle f(x_n) - z, x_{n+1} - z \rangle \\ &\leq (1 - \alpha_n)\|x_n - z\|^2 + \alpha_n(1 - \lambda_n)^2\|x_n - z\|^2 \\ &\quad + 2\alpha_n\lambda_n \langle f(x_n) - z, x_{n+1} - z \rangle \\ &\leq (1 - \alpha_n + \alpha_n(1 - \lambda_n))^2\|x_n - z\|^2 \\ &\quad + 2\alpha_n\lambda_n \langle f(x_n) - f(z), x_{n+1} - z \rangle + 2\alpha_n\lambda_n \langle f(z) - z, x_{n+1} - z \rangle. \end{aligned}$$

The Cauchy-Schwartz inequality gives

$$\begin{aligned} 2\alpha_n\lambda_n \langle f(x_n) - f(z), x_{n+1} - z \rangle &\leq 2\alpha_n\lambda_n \|f(x_n) - f(z)\| \|x_{n+1} - z\| \\ &\leq \alpha_n\lambda_n [\|f(x_n) - f(z)\|^2 + \|x_{n+1} - z\|^2]. \end{aligned}$$

So,

$$\begin{aligned} \|x_{n+1} - z\|^2 &\leq (1 - \alpha_n + \alpha_n(1 - \lambda_n)^2)\|x_n - z\|^2 \\ &\quad + \alpha_n\lambda_n (\|f(x_n) - f(z)\|^2 + \|x_{n+1} - z\|^2) \\ &\quad + 2\alpha_n\lambda_n \langle f(z) - z, x_{n+1} - z \rangle \\ &\leq (1 - \alpha_n + \alpha_n(1 - \lambda_n)^2 + \alpha_n\lambda_n\rho)\|x_n - z\|^2 \\ &\quad + \alpha_n\lambda_n \|x_{n+1} - z\|^2 + 2\alpha_n\lambda_n \langle f(z) - z, x_{n+1} - z \rangle. \end{aligned}$$

We can compute $(1 - \alpha_n + \alpha_n(1 - \lambda_n)^2 + \alpha_n\lambda_n\rho)$ and simplify:

$$\begin{aligned} \|x_{n+1} - z\|^2 &= (1 - \alpha_n\lambda_n - \alpha_n\lambda_n(1 - \rho - \lambda_n))\|x_n - z\|^2 \\ &\quad + \alpha_n\lambda_n\|x_{n+1} - z\|^2 + 2\alpha_n\lambda_n\langle f(z) - z, x_{n+1} - z \rangle. \end{aligned}$$

Then from the foregoing it follows that

$$\begin{aligned} \|x_{n+1} - z\|^2 &\leq \left(1 - \frac{\alpha_n\lambda_n(1 - \rho - \lambda_n)}{1 - \alpha_n\lambda_n}\right)\|x_n - z\|^2 \\ &\quad + 2\frac{\alpha_n\lambda_n}{1 - \alpha_n\lambda_n}\langle f(z) - z, x_{n+1} - z \rangle. \end{aligned} \quad (3.11)$$

Putting

$$\begin{aligned} a_n &= \|x_n - z\|^2, \\ \gamma_n &= \frac{\alpha_n\lambda_n(1 - \rho - \lambda_n)}{1 - \alpha_n\lambda_n} \end{aligned}$$

and

$$\delta_n = 2\frac{\alpha_n\lambda_n}{1 - \alpha_n\lambda_n}\langle f(z) - z, x_{n+1} - z \rangle,$$

then (3.11) becomes

$$a_{n+1} \leq (1 - \gamma_n)a_n + \delta_n.$$

Note that $\lim_n \gamma_n = 0$ and

$$\limsup_n \frac{\delta_n}{\gamma_n} = \limsup_n 2\frac{\langle f(z) - z, x_{n+1} - z \rangle}{(1 - \rho - \lambda_n)} \leq 0,$$

by (3.10). Thus we may apply Lemma 2.2 to conclude that

$$\lim_n a_n = \lim_n \|x_n - z\| = 0. \quad \square$$

References

- [1] E. Blum and E. Oettli, 'From optimization and variational inequalities to equilibrium problems', *Math. Student* **63** (1994), 123–145.
- [2] F. E. Browder, 'Convergence of approximation to fixed points of nonexpansive nonlinear mappings in Hilbert spaces', *Arch. Ration. Mech. Anal.* **24** (1967), 82–90.
- [3] C. Byrne, 'A unified treatment of some iterative algorithms in signal processing and image reconstruction', *Inverse Problems* **20** (2004), 103–120.
- [4] A. Cabot, 'Proximal point algorithm controlled by a slowly vanishing term: applications to hierarchical minimization', *SIAM J. Optim.* **15** (2005), 555–572.
- [5] Y. Censor, A. Motova and A. Segal, 'Perturbed projections and subgradient projections for the multiple-sets split feasibility problem', *J. Math. Anal. Appl.* **327** (2007), 1244–1256.
- [6] P. L. Combettes and S. A. Hirstoaga, 'Equilibrium programming in Hilbert spaces', *J. Nonlinear Convex Anal.* **6**(1) (2005), 117–136.
- [7] S. D. Flåm and A. S. Antipin, 'Equilibrium programming using proximal-like algorithms', *Math. Program.* **78** (1997), 29–41.

- [8] A. Göpfert, H. Riahi, C. Tammer and C. Zălinescu, *Variational Methods in Partially Ordered Spaces* (Springer, New York, 2003).
- [9] B. Halpern, 'Fixed points of nonexpansive maps', *Bull. Amer. Math. Soc.* **73** (1967), 957–961.
- [10] A. N. Iusem and W. Sosa, 'Iterative algorithms for equilibrium problems', *Optimization* **52** (2003), 301–316.
- [11] I. V. Konnov, 'Application of the proximal point method to nonmonotone equilibrium problems', *J. Optim. Theory Appl.* **119** (2003), 317–333.
- [12] P. L. Lions, 'Approximation de points fixes de contractions', *C. R. Acad. Sci. Sér. A-B Paris* **284** (1977), 1357–1359.
- [13] Z.-Q. Luo, J.-S. Pang and D. Ralph, *Mathematical Programs with Equilibrium Constraints* (Cambridge University Press, Cambridge, 1996).
- [14] G. Marino and H. K. Xu, 'A general iterative method for nonexpansive mappings in Hilbert spaces', *J. Math. Anal. Appl.* **318**(1) (2006), 43–52.
- [15] A. Moudafi, 'Viscosity approximation methods for fixed-points problems', *J. Math. Anal. Appl.* **241** (2000), 46–55.
- [16] ———, 'Second-order differential proximal methods for equilibrium problems', *J. Inequal. Pure Appl. Math.* **4** (2003), art. 18.
- [17] ———, 'Krasnoselski–Mann iteration for hierarchical fixed-point problems', *Inverse Problems* **23** (2007), 1635–1640.
- [18] A. Moudafi and M. Théra, 'Proximal and dynamical approaches to equilibrium problems', in: *Ill-Posed Variational Problems and Regularization Techniques*, Lecture Notes in Economics and Math. Sys., 477 (Springer, New York, 1999).
- [19] W. Oettli, 'A remark on vector-valued equilibria and generalized monotonicity', *Acta Math. Vietnamica* **22** (1997), 215–221.
- [20] S. Reich, 'Strong convergence theorems for resolvents of accretive operators in Banach spaces', *J. Math. Anal. Appl.* **75** (1980), 287–292.
- [21] ———, 'Approximating fixed points of nonexpansive mappings', *Panamerican. Math. J.* **4**(2) (1994), 23–28.
- [22] N. Shioji and W. Takahashi, 'Strong convergence of approximated sequences for nonexpansive mappings in Banach spaces', *Proc. Amer. Math. Soc.* **125** (1997), 3641–3645.
- [23] M. Solodov, 'An explicit descent method for bilevel convex optimization', *J. Convex Anal.* **14** (2007), 227–237.
- [24] T. Suzuki, 'Strong convergence of Krasnoselskii and Mann's type sequences for one-parameter nonexpansive semigroups without Bochner integrals', *J. Math. Anal. Appl.* **305**(1) (2005), 227–239.
- [25] R. Wittmann, 'Approximation of fixed points of nonexpansive mappings', *Arch. Math.* **58** (1992), 486–491.
- [26] H. K. Xu, 'Iterative algorithms for nonlinear operators', *J. London Math. Soc.* **66** (2002), 240–256.
- [27] ———, 'Another control condition in an iterative method for nonexpansive mappings', *Bull. Austral. Math. Soc.* **65** (2002), 109–113.
- [28] ———, 'Remarks on an iterative method for nonexpansive mappings', *Comm. Appl. Nonlinear Anal.* **10**(1) (2003), 67–75.
- [29] ———, 'An iterative approach to quadratic optimization', *J. Optim. Theory Appl.* **116**(3) (2003), 659–678.
- [30] ———, 'Viscosity approximation methods for nonexpansive mappings', *J. Math. Anal. Appl.* **298** (2004), 279–291.
- [31] ———, 'A variable Krasnoselski–Mann algorithm and the multiple-set split feasibility problem', *Inverse Problems* **22** (2006), 2021–2034.
- [32] I. Yamada, 'The hybrid steepest descent method for the variational inequality problem over the intersection of fixed point sets of nonexpansive mappings', in: *Inherently Parallel Algorithm for Feasibility and Optimization and their Applications* (eds. D. Butnariu, Y. Censor and S. Reich) (Elsevier, 2001), pp. 473–504.

- [33] I. Yamada, N. Ogura and N. Shirakawa, 'A numerically robust hybrid steepest descent method for the convexity constrained generalized inverse problems', *Contemp. Math.* **313** (2002), 269–305.
- [34] Q. Yang and J. Zhao, 'Generalized KM theorems and their applications', *Inverse Problems* **22** (2006), 833–844.
- [35] Y. Yao and Y.-C. Liou, 'Weak and strong convergence of Krasnosel'skiĭ–Mann iteration for hierarchical fixed point problems', *Inverse Problems* **24** (2008), 501–508.

GIUSEPPE MARINO, Dipartimento di Matematica, Università della Calabria,
87036, Arcavacata di Rende (CS), Italy
e-mail: gmarino@unical.it

VITTORIO COLAO, Dipartimento di Matematica, Università della Calabria,
87036, Arcavacata di Rende (CS), Italy
e-mail: colao@mat.unical.it

LUIGI MUGLIA, Dipartimento di Matematica, Università della Calabria,
87036, Arcavacata di Rende (CS), Italy
e-mail: muglia@mat.unical.it

YONGHONG YAO, Department of Mathematics, Tianjin Polytechnic University,
Tianjin 300160, People's Republic of China
e-mail: yaoyonghong@yahoo.cn