

# Bison with benefits: towards integrating wildlife and ranching sectors on a public rangeland in the western USA

DUSTIN H. RANGLACK and JOHAN T. DU TOIT

**Abstract** The North American model of wildlife conservation, based on the public trust doctrine, is credited for the recovery of several charismatic wildlife species, including the plains bison *Bison bison*. In that model, wildlife is a public resource from which the private sector may not profit either individually or collectively. In recent years, however, resilience thinking is driving changes in the traditional state-run wildlife management model to allow for integrated multi-sector approaches at the landscape scale. A free-ranging herd of bison on public land in the Henry Mountains of Utah is used as a case study to consider if and how a community-based conservation programme could be developed for a state-managed wildlife resource to benefit all stakeholders. The Henry Mountains bison, which are disease-free, share the rangeland with cattle that are privately owned by individual ranchers and corporations with various economic goals and environmental values. The ranchers currently derive no benefits from the bison and have concerns regarding competition between bison and cattle. However, a threshold harvesting strategy with community participation could generate revenue to offset these concerns. It could also provide benefits to the local community, increase state revenue, and increase the size of the bison population while securing its long-term genetic viability. Implementation would initially require facilitation by policy specialists, after which we suggest a Henry Mountains bison partnership could serve as a model for bison recovery efforts elsewhere in North America.

**Keywords** Adaptive management, community-based conservation, human-wildlife conflict, North American model, public trust doctrine

## Introduction

The North American model of wildlife conservation (Geist et al., 2001) is based on the doctrine that

wildlife resources are held in the public trust and managed by government agencies according to the law. This public trust doctrine, which is rooted in Roman law and was first invoked by the U.S. Supreme Court in the mid 19th century, is now enshrined in laws that preclude personal or communal profit from wildlife (Organ & Mahoney, 2007). Implementation of management and enforcement of laws by state and provincial wildlife agencies is funded by license revenue, so although the purpose of the model is to protect wildlife from overexploitation, it is also dependent upon state-regulated hunting and angling. As a consequence, populations of wildlife species with high trophy value are typically managed according to hunting objectives, subject to resistance from ranchers concerned about competition with their livestock. This command-and-control, top-down management approach is credited with having brought several iconic wildlife species back from the brink of extinction, yet it represents the traditional population-focused paradigm of wildlife management, which is now being challenged (Chapin et al., 2009; Allen et al., 2011; Curtin & Parker, 2014). Reconciling conservation, agriculture and other competing land uses involves changing the culture of the relevant management agencies, and calls for an integrated approach at the landscape scale (Sayer et al., 2013).

Lessons in integrating community participation into wildlife management can be learnt from the developing world (Hill, 2009), where experience has shown that conservation efforts generally fail if local communities are denied access to, or benefits from, their traditional resources (Newmark & Hough, 2000). In such situations community-based conservation approaches have sought to incentivize local communities to become active participants in conservation by enabling them to derive direct sustainable benefits in the process (Spiteri & Nepal, 2006). For decades, integrated conservation and development projects have been advocated, funded and technically supported by foreign aid agencies and international NGOs but the best practices are seldom applied in developed countries (Garnett et al., 2007). Here we suggest how cattle ranching on public rangeland could be at least partially integrated with the management of American bison *Bison bison*, the iconic species of the North American model (Organ et al., 2010).

Once numbering in the millions, bison populations across North America declined to < 100 wild individuals in total by the late 1800s (Hedrick, 2009). Of the c. 500,000 bison in

DUSTIN H. RANGLACK (Corresponding author) Department of Wildland Resources and Ecology Center, Utah State University, Logan, UT 84322-5230, USA. E-mail [dhrranglack@gmail.com](mailto:dhrranglack@gmail.com)

JOHAN T. DU TOIT Department of Wildland Resources, Utah State University, Logan, USA

Received 1 October 2014. Revision requested 11 November 2014.

Accepted 9 December 2014. First published online 8 May 2015.

This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (<http://creativecommons.org/licenses/by-nc-nd/3.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited. The written permission of Cambridge University Press must be obtained for commercial re-use. This article is subject to the Cambridge Core terms of use, available at <http://www.cambridge.org/core/terms>. <https://doi.org/10.1017/S0030605314001197>




FIG. 1 The Henry Mountains area of southern Utah, USA, with the area used by the bison *Bison bison* herd delineated by the black line. The shaded rectangle on the inset indicates the location of the main map in Utah.

North America at present, only c. 20,000 are in conservation herds (Gates & Ellison, 2010). Despite the success of many conservation herds and the large number of commercial herds, bison as a wildlife species has not recovered ecologically (Freese et al., 2007; Sanderson et al., 2008). Competition, disease and genetic introgression give rise to conflict between bison conservationists and livestock managers, and thus most bison herds are separated from cattle by fences and hazing practices (Meagher, 1989; Aune & Wallen, 2010). One of the few places in the USA where free-ranging bison share a public rangeland with cattle is in the Henry Mountains of southern Utah (Fig. 1).

Established in the early 1940s with bison from Yellowstone National Park (Popov & Low, 1950; Nelson, 1965), the Henry Mountains bison herd comprises c. 325 adults (post-hunt) and is controlled primarily by sport hunting. The presence of bison on public allotments leased for cattle grazing has become a source of contention between local cattle ranchers and the state and federal management agencies (UDWR, 2007; Ranglack et al., 2015). A search for mentions of the Henry Mountains bison in a major daily newspaper in Utah (*Deseret News*), and the archives of the Utah State Legislature, revealed that the contention diminishes when rangeland conditions are good but returns when annual rainfall returns to average or below-average levels (Fig. 2). Earlier work showed that bison and cattle have 91% dietary similarity in the Henry Mountains (van Vuren & Bray, 1983). This, combined with their conspicuous presence on the landscape as a result of their herding behaviour, dust wallowing, trampling, and




FIG. 2 The number of mentions of the Henry Mountains bison conflict in a major Salt Lake City newspaper (*Deseret News*) and the Utah State Legislature during 1991–2013, plotted with annual precipitation and mean annual precipitation at the nearby Hanksville airport. The conflict was in the news in the early 1990s but became quiescent during a decade of mostly favourable precipitation. The re-emergence of the conflict, now in the legislature as well as the news, coincided with the onset of a dry period in 2007.

large dung pats, has led to a local perception that bison are important competitors with cattle for grazing resources.

The Henry Mountains lie within one of the most important conservation priority areas of the roadless Bureau of Land Management lands in the western USA (Dickson et al., 2014). With the only free-ranging, disease-free, genetically pure, and huntable wild bison herd in the world, the rugged environment offers a unique opportunity to sportsmen, and thus there is high demand for hunting permits (tags). At the Western Hunt Expo in Salt Lake City in 2014, Sportsmen for Fish and Wildlife sold two Henry Mountains bison tags under the state’s conservation permit programme at auction for USD 19,000 and USD 18,500. These funds were primarily for use by the state in projects that benefit bison, with oversight from Sportsmen for Fish and Wildlife. In 2013, 8,135 prospective hunters (5,618 Utah residents, 2,517 non-residents) applied for permits to hunt bison on the Henry Mountains, and 95 permits were issued through a lottery programme, generating almost USD 50,000 of license revenue for the state.

The current management objective for the Henry Mountains bison is to maintain a stable population size by harvesting, with an escapement threshold (Lande et al., 1997) of 325 adults as agreed upon by state and federal agencies and the Henry Mountains Grazing Association (UDWR, 2007). At the current population size, the Henry Mountains bison are neither major consumers of the forage resources used by cattle nor competitors for the habitats that local ranchers consider to be most important for their cattle (Ranglack et al., 2015; D.H. Ranglack & J.T. du Toit, unpubl. data). Nevertheless, while they continue to derive no

benefits from the bison, local ranchers can be expected to continue using their contention of bison–cattle competition to attract political attention to their ranching interests. Allowing a fair system for the local ranching community to benefit directly from the bison that share the public rangeland with their cattle, while also securing bison conservation, may be a solution. Implementation would require the negotiation of legal, political, organizational and economic barriers but that process would be contingent upon an ecologically and economically feasible plan. Our objective is to investigate the feasibility of incrementally raising the escapement threshold for the Henry Mountains bison population, commensurately reducing the stocking rate of cattle, and using benefits from bison hunting to compensate the affected ranchers and provide a fund for community development.

### Study area

The Henry Mountains study area in south-central Utah (Fig. 1) includes arid, semi-arid and subalpine habitats for bison, which migrate seasonally between northern (summer) and southern (winter) parts of the range and utilize almost all vegetation types and elevations. Apart from bison, cattle are the only other large grazers in the region. Mule deer *Odocoileus hemionus* are present but their preference for forbs suggests competition with the grazers is negligible (van Vuren & Bray, 1983). Black-tailed jackrabbits *Lepus californicus* and desert cottontail rabbits *Sylvilagus audubonii* are common at low and mid elevations. Mountain lions *Puma concolor* and coyotes *Canis latrans* are the only large predators but their population densities are limited by well-established predator control efforts implemented by both government and private entities. For a description of the study area see Nelson (1965) and van Vuren & Bray (1986).

### Methods

**Bison population data** The Henry Mountains bison population has been surveyed annually by the Utah Division of Wildlife Resources (in July or August) since 1949 to estimate the pre-hunt population size ( $N$ ). Since 1983 the population has been harvested according to a fixed escapement strategy; the escapement threshold ( $c$ ) has been adjusted several times by agreement among representatives of the Utah Division of Wildlife Resources, the Bureau of Land Management, and the Henry Mountains Grazing Association. The number of bison harvested ( $H$ ) each year is the sum of the number killed by hunters and the number of live bison removed by the Utah Division of Wildlife Resources for translocation (Fig. 3).




FIG. 3 The Henry Mountains bison population trajectory during 1949–2012, with the total pre-hunt population count from the summer bison surveys conducted by the Utah Division of Wildlife Resources, the total harvest of hunted and removed (live) bison, and the escapement threshold as it was adjusted during 1983–2012.

**Harvest calculations** We performed a hypothetical exercise in which we allowed the pre-hunt bison population to increase above the level at which it is maintained by the current escapement threshold ( $c = 325$ ), considering incremental escapement thresholds up to  $c = 500$ . To quantify the actual harvesting rate ( $h$ ) that had been applied to the population during its history, we calculated  $h = H/N$  and then the mean  $h$  over the 30 years (1983–2012) during which an escapement threshold strategy has been in place. This yielded  $h = 0.20 \pm \text{SE } 0.015$ , or 20% of the adult population harvested, on average, each year. By comparison, the maximum intrinsic population growth rate ( $r_{max}$ ) for bison can be estimated from mean adult body mass ( $M$ ) as  $r_{max} = 1.5M^{-0.36}$  (Caughley & Krebs, 1983); using  $M = 500$  kg (Reynolds et al., 2003) yields  $r_{max} = 0.16$ . Applying the historical harvesting rate annually ( $h = 0.20$ ) would be unsustainable in the long term, therefore, and so we made the conservative assumption that  $h = r_{max}$  and used this theoretical value ( $h = 0.16$ ) for our calculations of future potential harvesting. With a fixed harvesting rate ( $h$ ) and a series of incremental values of escapement threshold ( $c$ ) the hypothetical pre-hunt population size ( $N$ ) was calculated for each threshold as  $N = c/(1 - h)$ , and the annual potential yield ( $H$ ) with that threshold as  $H = N - c$ .

**Financial calculations** With the current escapement threshold ( $c = 325$ ) holding the bison population at the maximum size tolerated by the Henry Mountains Grazing Association because of concerns regarding competition with cattle (the veracity of which is immaterial to this exercise), we assumed that any raising of the threshold ( $c > 325$ ) would have to be accompanied by destocking of cattle, for

which the ranchers would have to be compensated. For simplicity we followed Bureau of Land Management practice in considering bison and cattle as equivalent animal units, so one bison on the Henry Mountains rangeland for 1 year is equivalent to 12 animal unit months in terms of grazing effect. Thus, each additional bison allowed to remain on the Henry Mountains rangeland for 1 year would require 12 animal unit months to be released from cattle grazing, and those animal unit months would have to be purchased elsewhere to maintain the rancher's level of animal production. We assumed those animal unit months would be purchased on the open market from private landowners, and therefore we used the 2013 market rate for Utah of USD 14.50 per animal unit month (NASS, 2014). To provide the revenue to compensate for destocking cattle on the Henry Mountains rangeland, we arbitrarily allocated one bison tag for auction each year if the escapement threshold was raised to  $c = 350$ , and one additional tag per year for each increment of 50 above that (0 tags for  $c = 325$ , 1 tag for  $c = 350$ , 4 tags for  $c = 500$ ). We used the 2014 auction value of USD 19,000 per tag. As a notional benefit distribution scheme, we assumed all of the auction proceeds would go to the Henry Mountains Grazing Association, from where they would be distributed on a pro rata basis to affected permittees as direct compensation in market terms for the animal unit months they gave up, with the remainder going into a community fund such that no individual would derive personal financial profit.

## Results

Allowing the bison population to increase by raising the escapement threshold and reducing the number of cattle on the rangeland commensurately could ultimately provide a larger annual yield of bison. This could provide financial compensation for affected ranchers, and an annual community dividend (Table 1). Once the bison population had become regulated at a new escapement threshold, and assuming current market conditions, the local community could derive a minimum net benefit of USD 10,300 annually (after ranchers had been compensated for animal unit month reductions) for every increment of 50 bison in the escapement threshold. The state wildlife management agency could also increase its annual offtake of bison through any combination of hunting and live removal.

## Discussion

Even with a more conservative harvesting rate than has been applied in the Henry Mountains historically, each increase of 50 bison in the escapement threshold could result in nine more bison harvested each year. Of those, at least one could be allocated to the community for auction and up to eight could be added to those available for the state wildlife

TABLE 1 The required harvest ( $H$ ) to maintain the Henry Mountains bison *Bison bison* population under various potential escapement thresholds, using a conservative harvesting rate ( $h = 0.16$ ), where  $N$  is the hypothetical pre-hunt population size, with number of animal unit months, cost, number of tags, and net benefit.

Escapement threshold	$N$	$H$	Animal unit months <sup>1</sup>	Cost (USD)	Tags <sup>2</sup>	Net benefit (USD) <sup>3</sup>
325	387	62	0	0	0	0
350	417	67	300	4,350	1	14,650
400	476	76	900	13,050	2	24,950
450	536	86	1,500	21,750	3	35,250
500	595	95	2,100	30,450	4	45,550

<sup>1</sup>The commensurate reduction in cattle grazing (animal unit months) would incur the cost of purchasing grazing rights elsewhere on the open market, calculated using 2013 mean market rates for Utah. Compensation could be from the proceeds of an annual public auction of bison hunts (tags), which could be distributed on a pro-rata basis to affected ranchers, after which a net benefit could go to the community for local projects.

<sup>2</sup>An illustrative minimal number of tags for the community for each harvesting threshold

<sup>3</sup>Calculated assuming the 2014 auction value of USD 19,000 per tag

management agency to allocate to hunters via their lottery system or to remove for metapopulation management. As the Henry Mountains bison population is currently  $< 10\%$  of the size of the cattle population permitted on the Henry Mountains rangeland, and with each increase in the bison escapement threshold matched with a commensurate decrease in the cattle population, we are assuming no density dependent effects. As with all harvesting strategies involving community-based wildlife management, the initial calculations are to establish a starting point from where adaptive management could proceed (du Toit, 2002). Nevertheless, our conservative calculations demonstrate that a policy change could benefit both the local community and the state wildlife management agency.

From a conservation perspective the implications are that the Henry Mountains bison population could be increased to reduce the risk of small population syndrome (Gross & Wang, 2005). Additional animals could then become available for translocation as founders in other suitable habitats, to augment existing subpopulations, as exchange for genetically pure bison from elsewhere (metapopulation management), or as exchange for batches of animals of other species of concern. Genetically pure, disease-free bison, such as those in the Henry Mountains, are valuable to conservation, with few other populations meeting those requirements (Halbert & Derr, 2007; Aune et al., 2010).

If some financial benefits from the Henry Mountains bison were channelled directly to the local community this could incentivize local ranchers to adjust their cattle management to improve conditions for the bison population. There could also be opportunities for the local community to diversify operations to protect against future changes in

the livestock market or as a buffer against changing environmental conditions; for example, the community could stipulate that a hunter who purchases their tag(s) at auction must use one of the local ranchers as a guide. The community fund could also be used for projects that benefit the entire community, such as scholarships for local students, or improving community infrastructure (Frost & Bond, 2008).

From global analyses it is evident that conserving wildlife on rangelands depends on the active involvement of local communities (Deutsch, 2010). Integrated science-based management planning that takes into account the complexities of the social–ecological system is key for managing populations of large ungulates on rangelands shared with domestic livestock (Weisberg et al., 2002). Also, even where wildlife resources are publicly owned, wildlife management planning should take into account negative impacts (whether perceived or actual) on the communities that live with the wildlife resource (Madden, 2004). Offsetting such negative impacts is best achieved with a direct and tangible benefit that varies depending on the overall health or size of the wildlife population(s) in question (Berkes, 2007). Allowing a local community to become an active partner in, and beneficiary of, a wildlife management programme builds trust among partners and enhances the adaptive capacity of the social–ecological system (Berkes, 2004).

Although the Henry Mountains bison example demonstrates the ecological and financial feasibility of a staggered incentive system, we recognize the challenges to implementation within local, state and federal institutions. This represents a significant change to the present management system for wildlife resources on public lands in North America. Our purpose is to explore and describe the possibilities so that representatives from the community and state and federal agencies have a basis for ongoing dialogue and planning. Management of the Henry Mountains bison is a contentious issue that needs additional inputs from policy specialists, resource economists and social scientists, but our simple population analysis (Table 1) points to a solution. It is crucial that wildlife managers and cattle ranchers work together for American bison conservation (Freese et al., 2007). The same applies for European bison *Bison bonasus*, for which there is abundant suitable habitat outside protected areas (Kuemmerle et al., 2011). In a broader context we present the Henry Mountains bison as a case study of how wildlife management in North America could begin to adapt at a time of increasing challenge to the historically successful North American model (Duda et al., 1998; Manfredo et al., 2003; Teel & Manfredo, 2009; Organ et al., 2010).

## Acknowledgements

Bison population data were provided by the Utah Division of Wildlife Resources. P. Adler, M. Fisher, F. Howe,

D. Koons, K. Redford and an anonymous reviewer provided valuable comments.

## References

- ALLEN, C.R., CUMMING, G.S., GARMESTANI, A.S., TAYLOR, P.D. & WALKER, B.H. (2011) Managing for resilience. *Wildlife Biology*, 17, 337–349.
- AUNE, K., GATES, C.C. & BOYD, D. (2010) Reportable or notifiable diseases. In *American Bison: Status Survey and Conservation Guidelines 2010* (eds C.C. Gates, C.H. Freese, P.J.P. Gogan & M. Kotzman), pp. 27–38. IUCN, Gland, Switzerland.
- AUNE, K. & WALLEN, R. (2010) Legal status, policy issues and listings. In *American Bison: Status Survey and Conservation Guidelines 2010* (eds C.C. Gates, C.H. Freese, P.J.P. Gogan & M. Kotzman), pp. 63–84. IUCN, Gland, Switzerland.
- BERKES, F. (2004) Rethinking community-based conservation. *Conservation Biology*, 18, 621–630.
- BERKES, F. (2007) Community-based conservation in a globalized world. *PNAS*, 104, 15188–15193.
- CAUGHLEY, G. & KREBS, C.J. (1983) Are big mammals simply little mammals writ large? *Oecologia*, 59, 7–17.
- CHAPIN, F.S., CARPENTER, S.R., KOFINAS, G.P., FOLKE, C., ABEL, N., CLARK, W.C. et al. (2009) Ecosystem stewardship: sustainability strategies for a rapidly changing planet. *Trends in Ecology & Evolution*, 25, 241–249.
- CURTIN, C.G. & PARKER, J.P. (2014) Foundations of resilience thinking. *Conservation Biology*, 28, 912–923.
- DEUTSCH, J.C. (2010) Synthesis: local and global solutions to the challenge of keeping rangelands wild. In *Wild Rangelands: Conserving Wildlife While Maintaining Livestock in Semi-Arid Ecosystems* (eds J.T. du Toit, R. Kock & J.C. Deutsch), pp. 393–402. Wiley-Blackwell and Zoological Society of London, Chichester, UK.
- DICKSON, B.G., ZACHMANN, L.J. & ALBANO, C.M. (2014) Systematic identification of potential conservation priority areas on roadless Bureau of Land Management lands in the western United States. *Biological Conservation*, 178, 117–127.
- DUDA, M.D., BISSELL, S.J. & YOUNG, K.C. (1998) *Wildlife and the American Mind: Public Opinion on and Attitudes Toward Fish and Wildlife Management*. Responsive Management, Harrisburg, USA.
- DU TOIT, J.T. (2002) Wildlife harvesting guidelines for community-based wildlife management: a southern African perspective. *Biodiversity and Conservation*, 11, 1403–1416.
- FREESE, C.H., AUNE, K.E., BOYD, D.P., DERR, J.N., FORREST, S.C., GATES, C.C. et al. (2007) Second chance for the plains bison. *Biological Conservation*, 136, 175–184.
- FROST, P.G.H. & BOND, I. (2008) The CAMPFIRE programme in Zimbabwe: payments for wildlife services. *Ecological Economics*, 65, 776–787.
- GARNETT, S.T., SAYER, J. & DU TOIT, J. (2007) Improving the effectiveness of interventions to balance conservation and development: a conceptual framework. *Ecology and Society*, 12, 2.
- GATES, C.C. & ELLISON, K. (2010) Numerical and geographic status. In *American Bison: Status Survey and Conservation Guidelines 2010* (eds C.C. Gates, C.H. Freese, P.J.P. Gogan & M. Kotzman), pp. 55–62. IUCN, Gland, Switzerland.
- GEIST, V., MAHONEY, S.P. & ORGAN, J.F. (2001) Why hunting has defined the North American model of wildlife conservation. *Transactions of the North American Wildlife and Natural Resources Conference*, 66, 175–185.
- GROSS, J.E. & WANG, G. (2005) Effects of population control strategies on retention of genetic diversity in National Park Service bison

- (*Bison bison*) herds. Final report submitted to Yellowstone Research Group, USGS-BRD. United States Geological Survey, Bozeman, USA.
- HALBERT, N.D. & DERR, J.N. (2007) A comprehensive evaluation of cattle introgression into US federal bison herds. *Journal of Heredity*, 98, 1–12.
- HEDRICK, P.W. (2009) Conservation genetics and North American bison (*Bison bison*). *Journal of Heredity*, 100, 411–420.
- HILL, C.M. (2009) Working with communities to achieve conservation goals. In *Wildlife and Society: The Science of Human Dimensions* (eds M.J. Manfredo, J.J. Vaske, P.J. Brown, D.J. Decker & E.A. Duke), pp. 117–128. Island Press, Washington, DC, USA.
- KUEMMERLE, T., RADELOFF, V.C., PERZANOWSKI, K., KOZLO, P., SIPKO, T., KHOYETSKYY, P. et al. (2011) Predicting potential European bison habitat across its former range. *Ecological Applications*, 21, 830–843.
- LANDE, R., SÆTHER, B.E. & ENGEN, S. (1997) Threshold harvesting for sustainability of fluctuating resources. *Ecology*, 78, 1341–1350.
- MADDEN, F. (2004) Creating coexistence between humans and wildlife: global perspectives on local efforts to address human–wildlife conflict. *Human Dimensions of Wildlife*, 9, 247–257.
- MANFREDO, M.J., TEEL, T.L. & BRIGHT, A.D. (2003) Why are public values toward wildlife changing? *Human Dimensions of Wildlife*, 8, 287–306.
- MEAGHER, M. (1989) Evaluation of boundary control for bison of Yellowstone National Park. *Wildlife Society Bulletin*, 17, 15–19.
- NASS (NATIONAL AGRICULTURAL STATISTICS SERVICE) (2014) *17 State Grazing Fees: Animal Unit*. USDA-NASS, Washington, D.C., USA.
- NELSON, K.L. (1965) *Status and Habits of the American Buffalo* (*Bison bison*) in the Henry Mountain Area of Utah. Utah State Department of Fish and Game, Logan, USA.
- NEWMARK, W.D. & HOUGH, J.L. (2000) Conserving wildlife in Africa: integrated conservation and development projects and beyond. *BioScience*, 50, 585–592.
- ORGAN, J. & MAHONEY, S. (2007) The future of public trust: the legal status of the Public Trust Doctrine. *The Wildlife Professional*, 1, 18–22.
- ORGAN, J., MAHONEY, S. & GEIST, V. (2010) Born in the hands of hunters: the North American model of wildlife conservation. *The Wildlife Professional*, 4, 22–27.
- POPOV, B.H. & LOW, J.B. (1950) *Game, Fur Animal and Fish: Introductions into Utah*. Utah State Department of Fish and Game, Logan, USA.
- RANGLACK, D.H., DURHAM, S. & DU TOIT, J.T. (2015) Competition on the range: science vs perception in a bison–cattle conflict in the western USA. *Journal of Applied Ecology*, 52, 467–474.
- REYNOLDS, H.W., GATES, C.C. & GLAHOLT, R.D. (2003) Bison (*Bison bison*). In *Wild Mammals of North America: Biology, Management, and Conservation*, 2nd edition (eds G.A. Feldhamer, B.C. Thompson & J.A. Chapman), pp. 1009–1060. The Johns Hopkins University Press, Baltimore, USA.
- SANDERSON, E.W., REDFORD, K.H., WEBER, B., AUNE, K., BALDES, D., BERGER, J. et al. (2008) The ecological future of the North American bison: conceiving long-term, large-scale conservation of wildlife. *Conservation Biology*, 22, 252–266.
- SAYER, J., SUNDERLAND, T., GHAZOUL, J., PFUND, J.-L., SHEIL, D., MEIJAARD, E. et al. (2013) Ten principles for a landscape approach to reconciling agriculture, conservation, and other competing land uses. *PNAS*, 110, 8349–8356.
- SPITERI, A. & NEPAL, S.K. (2006) Incentive-based conservation programs in developing countries: a review of some key issues and suggestions for improvements. *Environmental Management*, 37, 1–14.
- TEEL, T.L. & MANFREDO, M.J. (2009) Understanding the diversity of public interests in wildlife conservation. *Conservation Biology*, 24, 128–139.
- UDWR (UTAH DIVISION OF WILDLIFE RESOURCES) (2007) *Bison Unit Management Plan: Unit 15, Henry Mountains*. Utah Division of Wildlife Resources, Salt Lake City, USA.
- VAN VUREN, D. & BRAY, M.P. (1983) Diets of bison and cattle on a seeded range in southern Utah. *Journal of Range Management*, 36, 499–500.
- VAN VUREN, D. & BRAY, M.P. (1986) Population dynamics of bison in the Henry Mountains, Utah. *Journal of Mammalogy*, 67, 503–511.
- WEISBERG, P.J., HOBBS, N.T., ELLIS, J.E. & COUGHENOUR, M.B. (2002) An ecosystem approach to population management of ungulates. *Journal of Environmental Management*, 65, 181–198.

## Biographical sketches

DUSTIN RANGLACK is interested in the ecology and conservation of large mammals in terrestrial ecosystems. He aims to help solve conservation problems by studying use of space by animals, predator–prey interactions, and human–wildlife interactions. JOHAN DU TOIT is interested in the conservation of terrestrial ecosystem processes through the fusion of science and management.