

March 1988

ISSN 0007-4853
Vol. 78(1), 1-171

Bulletin of Entomological Research



C·A·B International Institute of Entomology

C·A·B INTERNATIONAL INSTITUTE OF ENTOMOLOGY

Director

K. M. HARRIS, BSc, DSc, DipAgrSc, DTA, FIBiol

Editor

J. M. B. HARLEY, BSc

Headquarters, Publication Office and Library—56 Queen's Gate, London SW7 5JR, UK.
Telephone: 01-584-0067. Telegrams: ENTINFO LONDON SW7. BT GOLD/DIALCOM:
84: CAU006. Telex: 265871 MONREF G (Box CAU006). Fax: 01-581-1676.

Identification Service—c/o British Museum (Natural History), Cromwell Road, London SW7 5BD, UK.

The C·A·B International Institute of Entomology is one of 4 Institutes and 11 Bureaux that together form C·A·B International (CABI) a not-for-profit, inter-governmental organisation which exists to provide information, scientific and development services for agriculture and allied disciplines throughout the world.

The *Bulletin of Entomological Research* is a quarterly journal prepared by the Institute and published by C·A·B International.

Annual Subscription Rate 1988

| | |
|----------------------------------------------------------|--------|
| To subscribers in countries that are members of CABI | £74.00 |
| To subscribers in countries that are not members of CABI | £97.00 |

Back Numbers

Back volumes of the *Bulletin of Entomological Research* are available at a discount.

Orders and enquiries concerning subscriptions and back numbers should be sent to

Marketing and Sales Division,
C·A·B INTERNATIONAL,
WALLINGFORD, OXON, OX10 8DE, UK.
TELEPHONE: WALLINGFORD (0491) 32111
TELEX: 847964 (COMAGG G)
FAX: (0491) 33508
TELECOM GOLD/INTERNATIONAL DIALCOM: 84: CAU001

[Note: Prices are subject to change without notice.]

© C·A·B International, 1988. All rights reserved. No part of this publication may be reproduced in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without the prior permission of the copyright owner.

Disclaimer

The information contained herein, including any expression of opinion and any projection or forecast, has been obtained from or is based upon sources believed by us to be reliable but is not guaranteed as to accuracy or completeness. The information is supplied without obligation and on the understanding that any person who acts upon it or otherwise changes his/her position in reliance thereon does so entirely at his/her own risk.

C·A·B International does not accept responsibility for any trade advertisement included in this publication.

**BULLETIN
OF
ENTOMOLOGICAL
RESEARCH**

VOL. 78

**ISSUED BY THE C·A·B INTERNATIONAL
INSTITUTE OF ENTOMOLOGY**

EDITOR: J. M. B. HARLEY

LONDON

**C·A·B INTERNATIONAL INSTITUTE OF ENTOMOLOGY
56, QUEEN'S GATE, SW7 5JR**

1988

C·A·B INTERNATIONAL INSTITUTE OF ENTOMOLOGY

Director

K. M. HARRIS, BSc, DSc, DipAgrSc, DTA, FIBiol

Editor

J. M. B. HARLEY, BSc

Headquarters, Publication Office and Library—56 Queen's Gate, London SW7 5JR, UK.

Telephone: 01-584-0067. Telegrams: ENTINFO LONDON SW7. BT GOLD/DIALCOM: 84: CAU006. Telex: 265871 MONREF G (Box CAU006). Fax: 01-581-1676.

Identification Service—c/o British Museum (Natural History), Cromwell Road, London SW7 5BD, UK.

The C·A·B International Institute of Entomology is one of 4 Institutes and 11 Bureaux that together form C·A·B International (CABI) a not-for-profit, inter-governmental organisation which exists to provide information, scientific and development services for agriculture and allied disciplines throughout the world.

The *Bulletin of Entomological Research* is a quarterly journal prepared by the Institute and published by C·A·B International.

Annual Subscription Rate 1989

| | |
|----------------------------------------------------------|---------|
| To subscribers in countries that are members of CABI | £77.00 |
| To subscribers in countries that are not members of CABI | £103.00 |

Back Numbers

Back volumes of the *Bulletin of Entomological Research* are available at a discount.

Orders and enquiries concerning subscriptions and back numbers should be sent to

Marketing and Sales Division,
C·A·B INTERNATIONAL,
WALLINGFORD, OXON, OX10 8DE, UK.
TELEPHONE: WALLINGFORD (0491) 32111
TELEX: 847964 (COMAGG G)
FAX: (0491) 33508
TELECOM GOLD/INTERNATIONAL DIALCOM: 84: CAU001

[Note: Prices are subject to change without notice.]

© C·A·B International, 1988. All rights reserved. No part of this publication may be reproduced in any form or by any means, electronically, mechanically, by photocopying, recording or otherwise, without the prior permission of the copyright owner.

Disclaimer

The information contained herein, including any expression of opinion and any projection or forecast, has been obtained from or is based upon sources believed by us to be reliable but is not guaranteed as to accuracy or completeness. The information is supplied without obligation and on the understanding that any person who acts upon it or otherwise changes his/her position in reliance thereon does so entirely at his/her own risk.

C·A·B International does not accept responsibility for any trade advertisement included in this publication.

CONTENTS

PART 1 MARCH 1988

Review article

PAGE

- YANINEK, J. S. & HERREN, H. R. Introduction and spread of the cassava green mite, *Mononychellus tanajoae* (Bondar) (Acaria: Tetranychidae), an exotic pest in Africa and the search for appropriate control methods: a review 1

Original articles

| | | |
|---------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| QUICKE, D. L. J. | A new genus and species of Braconinae (Hymenoptera: Braconidae) parasitic on <i>Diatraea</i> sp. (Lepidoptera: Pyralidae) from Ecuador | 15 |
| FFRENCH-CONSTANT, R. H., CLARK, S. J. & DEVONSHIRE, A. L. | Effect of decline of insecticide residues on selection for insecticide resistance in <i>Myzus persicae</i> (Sulzer) (Hemiptera: Aphididae) | 19 |
| VALE, G. A., LOVEMORE, D. F., FLINT, S. & COCKBILL, G. F. | Odour-baited targets to control tsetse flies, <i>Glossina</i> spp. (Diptera: Glossinidae), in Zimbabwe | 31 |
| VAN SICKLE, J. & PHELPS, R. J. | Age distributions and reproductive status of declining and stationary populations of <i>Glossina pallidipes</i> Austen (Diptera: Glossinidae) in Zimbabwe | 51 |
| DOUBE, B. M. & MOOLA, F. | The effect of the activity of the African dung beetle <i>Catharsius tricornutus</i> De Geer (Coleoptera: Scarabaeidae) on the survival and size of the African buffalo fly, <i>Haematobia thirouxi potans</i> (Bezzi) (Diptera: Muscidae), in bovine dung in the laboratory | 63 |
| NEWTON, C. & DIXON, A. F. G. | A preliminary study of variation and inheritance of life-history traits and the occurrence of hybrid vigour in <i>Sitobion avenae</i> (F.) (Hemiptera: Aphididae) | 75 |
| NEWTON, P. J. | Movement and impact of <i>Trichogrammatoidea cryptophlebiae</i> Nagaraja (Hymenoptera: Trichogrammatidae) in citrus orchards after inundative releases against the false codling moth, <i>Cryptophlebia leucotreta</i> (Meyrick) (Lepidoptera: Tortricidae) | 85 |
| VAN RENSBURG, J. B. J., GILIOMEE, J. H. & WALTERS, M. C. | Aspects of the injuriousness of the maize stalk borer, <i>Busseola fusca</i> (Fuller) (Lepidoptera: Noctuidae) | 101 |
| AALBERSBERG, Y. K., VAN DER WESTHUIZEN, M. C. & HEWITT, P. H. | Natural enemies and their impact on <i>Diuraphis noxia</i> (Mordvilko) (Hemiptera: Aphididae) populations | 111 |
| MACAULAY, E. D. M., TATCHELL, G. M. & TAYLOR, L. R. | The Rothamsted Insect Survey '12-metre' suction trap | 121 |
| NOYES, J. S. | <i>Metaphycus baruensis</i> sp.n. (Hymenoptera: Encyrtidae) parasitic in green scales, <i>Coccus</i> spp. (Hemiptera: Coccidae), on coffee in Kenya | 131 |
| TINGLE, C. C. D. & COPLAND, M. J. W. | Effects of temperature and host-plant on regulation of glasshouse mealybug (Hemiptera: Pseudococcidae) populations by introduced parasitoids (Hymenoptera: Encyrtidae) | 135 |
| MADUBUNYI, L. C. | The seasonal breeding rate of <i>Glossina tachinoides</i> Westwood (Diptera: Glossinidae) in peridomestic agroecosystems of the Nsukka area, Nigeria | 143 |

CONTENTS

| | PAGE |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| PONT, A. C. A shoot-fly, <i>Atherigona ramu</i> sp. n. (Diptera: Muscidae), attacking sugarcane in Papua New Guinea | 151 |
| VAN SICKLE, J. Invalid estimates of rate of population increase from <i>Glossina</i> (Diptera: Glossinidae) age distributions | 155 |
| FFRENCH-CONSTANT, R. H. & DEVONSHIRE, A. L. Monitoring frequencies of insecticide resistance in <i>Myzus persicae</i> (Sulzer) (Hemiptera: Aphididae) in England during 1985–86 by immunoassay | 163 |

PART 2 JUNE 1988

Original articles

| | PAGE |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| LUTWAMA, J. J. & MATANMI, B. A. Efficacy of <i>Bacillus thuringiensis</i> subsp. <i>kurstaki</i> and <i>Baculovirus heliothis</i> foliar applications for suppression of <i>Helicoverpa armiger</i> (Hübner) (Noctuidae) and other lepidopterous larvae on tomato in south-western Nigeria | 173 |
| PAGE, W. W. Varying durations of arrested oocyte development in relation to migration in the African armyworm moth, <i>Spodoptera exempta</i> (Walker) (Lepidoptera: Noctuidae) | 181 |
| VAN DRIESCHE, R. G. Field measurement of population recruitment of <i>Apanteles glomeratus</i> (L.) (Hymenoptera: Braconidae), a parasitoid of <i>Pieris rapae</i> (L.) (Lepidoptera: Pieridae), and factors influencing adult parasitoid foraging success in kale | 199 |
| TAYLOR, B. The impact of yellow stem-borer, <i>Scirpophaga incertulas</i> (Walker) (Lepidoptera: Pyralidae), on deepwater rice, with special reference to Bangladesh | 209 |
| KELLY, P. M., STERLING, P. H., SPEIGHT, M. R. & ENTWISTLE, P. F. Preliminary spray trials of a nuclear polyhedrosis virus as a control agent for the brown-tail moth, <i>Euproctis chrysorrhoea</i> (L.) (Lepidoptera: Lymantriidae) | 227 |
| MATTHIESSEN, J. N. & PALMER, M. J. Prediction of temperatures in cattle dung for estimating development times of coprophilous organisms | 235 |
| LOXDALE, H. D. & BROOKES, C. P. Electrophoretic study of enzymes from cereal aphid populations. V. Spatial and temporal genetic similarity of holocyclic populations of the bird-cherry oat aphid, <i>Rhopalosiphum padi</i> (L.) (Hemiptera: Aphididae), in Britain | 241 |
| REYNOLDS, D. R. & RILEY, J. R. A migration of grasshoppers, particularly <i>Diabolocatantops axillaris</i> (Thunberg) (Orthoptera: Acrididae), in the West African Sahel | 251 |
| EISEMANN, C. H. Upwind flight by gravid Australian sheep blowflies, <i>Lucilia cuprina</i> (Wiedemann) (Diptera: Calliphoridae), in response to stimuli from sheep | 273 |
| BURSELL, E., GOUGH, A. J. E., BEEVOR, P. S., CORK, A., HALL, D. R. & VALE, G. A. Identification of components of cattle urine attractive to tsetse flies, <i>Glossina</i> spp. (Diptera: Glossinidae) | 281 |
| VALE, G. A., HALL, D. R. & GOUGH, A. J. E. The olfactory responses of tsetse flies, <i>Glossina</i> spp. (Diptera: Glossinidae), to phenols and urine in the field | 293 |
| BRADY, J. & SHERENI, W. Landing responses of the tsetse fly <i>Glossina morsitans morsitans</i> Westwood and the stable fly <i>Stomoxys calcitrans</i> (L.) (Diptera: Glossinidae & Muscidae) to black-and-white patterns: a laboratory study | 301 |
| NOYES, J. S. <i>Gyranusoidea tebygi</i> sp. n. (Hymenoptera: Encyrtidae), a parasitoid of <i>Rastrococcus</i> (Hemiptera: Pseudococcidae) on mango in India | 313 |

CONTENTS

| | PAGE |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| LANGTON, P. H., CRANSTON, P. S. & ARMITAGE, P. The parthenogenetic midge of water supply systems, <i>Paratanytarsus grimmii</i> (Schneider) (Diptera: Chironomidae) | 317 |
| RAHBÉ, Y., FEBVAY, G. & KERMARREC, A. Foraging activity of the attine ant <i>Acromyrmex octospinosus</i> (Reich) (Hymenoptera: Formicidae) on resistant and susceptible yam varieties | 329 |
| PADGHAM, D. E. & WOODHEAD, S. Variety-related feeding patterns in the brown planthopper, <i>Nilaparvata lugens</i> (Stål) (Hemiptera: Delphacidae), on its host, the rice plant | 339 |
| BROWN, P. A. & BLACKMAN, R. L. Karyotype variation in the corn leaf aphid, <i>Rhopalosiphum maidis</i> (Fitch), species complex (Hemiptera: Aphididae) in relation to host-plant and morphology | 351 |

PART 3 SEPTEMBER 1988

Original articles

| | PAGE |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| DENT, D. R. & PAWAR, C. S. The influence of moonlight and weather on catches of <i>Helicoverpa armigera</i> (Hübner) (Lepidoptera: Noctuidae) in light and pheromone traps | 365 |
| MCGEACHIE, W. J. A remote sensing method for the estimation of light-trap efficiency | 379 |
| LANGLEY, P. A., HALL, M. J. R., FELTON, T. & CEESAY, M. Determining the age of tsetse flies, <i>Glossina</i> spp. (Diptera: Glossinidae): an appraisal of the pteridine fluorescence technique | 387 |
| VAN DRIESCHE, R. G. Survivorship patterns of larvae of <i>Pieris rapae</i> (L.) (Lepidoptera: Pieridae) in Massachusetts kale, with special reference to mortality due to <i>Apanteles glomeratus</i> (L.) (Hymenoptera: Braconidae) | 397 |
| TEEL, P. D., BAY, D. E. & AJIDAGBA, P. A. Ecology, distribution and host relationships of ticks (Acaria: Ixodidae) infesting livestock in Mali | 407 |
| DAVIS, A. L. V., DOUBE, B. M. & McLENNAN, P. D. Habitat associations and seasonal abundance of coprophilous Coleoptera (Staphylinidae, Hydrophilidae and Histeridae) in the Hluhluwe region of South Africa | 425 |
| HUDDLESTON, T. & WALKER, A. K. <i>Cardiochiles</i> (Hymenoptera: Braconidae), a parasitoid of lepidopterous larvae, in the Sahel of Africa, with a review of the biology and host relationships of the genus | 435 |
| OTIENO, W. A., ONYANGO, T. O., PILE, M. M., LAURENCE, B. R., DAWSON, G. W., WADHAMS, L. J. & PICKETT, J. A. A field trial of the synthetic oviposition pheromone with <i>Culex quinquefasciatus</i> Say (Diptera: Culicidae) in Kenya ... | 463 |
| HEMINGWAY, J., BONNING, B. C., JAYAWARDENA, K. G. I., WEERASINGHE, I. S., HERATH, P. R. J. & OOUCHI, H. Possible selective advantage of <i>Anopheles</i> spp. (Diptera: Culicidae) with the oxidase- and acetylcholinesterase-based insecticide resistance genes after exposure to organophosphates or an insect growth regulator in Sri Lankan rice fields | 471 |
| PERERA, P. A. C. R., HASSELL, M. P. & GODFRAY, H. C. J. Population dynamics of the coconut caterpillar, <i>Opisina arenosella</i> Walker (Lepidoptera: Xyloryctidae), in Sri Lanka | 479 |
| MCDONALD, G. & FARROW, R. A. Migration and dispersal of the Rutherglen bug, <i>Nysius vinitor</i> Bergroth (Hemiptera: Lygaeidae), in eastern Australia | 493 |

CONTENTS

| | PAGE |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| WINDER, J. A., SANDS, D. P. A. & KASSULKE, R. C. The life history, host specificity and potential of <i>Alagoasa parana</i> Samuelson (Coleoptera: Chrysomelidae) for biological control of <i>Lantana camara</i> in Australia | 511 |
| WILSON, M. R. The genus <i>Myndus</i> (Hemiptera: Cixiidae) in the Solomon Islands and Vanuatu and its relation to foliar decay of coconut palms in Vanuatu | 519 |
| TINGEY, W. M. & LAMONT, W. J., JR Insect abundance in field beans altered by intercropping | 527 |
| MOORES, G. D., DEVONSHIRE, A. L. & DENHOLM, I. A microtitre plate assay for characterizing insensitive acetylcholinesterase genotypes of insecticide-resistant insects | 537 |
| BOUČEK, Z. <i>Tamarixia leucaenae</i> sp. n. (Hymenoptera: Eulophidae) parasitic on the leucaena psyllid <i>Heteropsylla cubana</i> Crawford (Hemiptera) in Trinidad | 545 |

PART 4 DECEMBER 1988

Original articles

| | PAGE |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| LE SUEUR, D. & SHARP, B. L. The breeding requirements of three members of the <i>Anopheles gambiae</i> Giles complex (Diptera: Culicidae) in the endemic malaria area of Natal, South Africa | 549 |
| COX, J. M. & WETTON, M. N. Identification of the mealybug <i>Planococcus halli</i> Ezzat & McConnell (Hemiptera: Pseudococcidae) commonly occurring on yams (<i>Dioscorea</i> spp.) in Africa and the West Indies | 561 |
| BARCLAY, H. J. Models for combining methods of pest control: food-baited and pheromone-baited traps containing either insecticide or chemosterilant | 573 |
| GREEN, C. H. The effect of colour on trap- and screen-orientated responses in <i>Glossina palpalis palpalis</i> (Robineau-Desvoidy) (Diptera: Glossinidae) | 591 |
| UMEH, E. D. N. Development, oviposition, host feeding and sex determination in <i>Epidinocarsis lopezi</i> (De Santis) (Hymenoptera: Encyrtidae) | 605 |
| BLACKWELL, A. The significance of proleg disruption by chlordimeform to sublethal control of larvae of <i>Pieris brassicae</i> (L.) (Lepidoptera: Pieridae) in the field ... | 613 |
| GILLETT, S. D. Solitarization in the desert locust, <i>Schistocerca gregaria</i> (Forskål) (Orthoptera: Acrididae) | 623 |
| RIDSDILL-SMITH, T. J. & MATTHIESSEN, J. N. Bush fly, <i>Musca vetustissima</i> Walker (Diptera: Muscidae), control in relation to seasonal abundance of scarabaeine dung beetles (Coleoptera: Scarabaeidae) in south-western Australia | 633 |
| JEPSON, P. C. & HEALY, T. P. The location of floral nectar sources by mosquitoes: an advanced bioassay for volatile plant odours and initial studies with <i>Aedes aegypti</i> (L.) (Diptera: Culicidae) | 641 |
| HEALY, T. P. & JEPSON, P. C. The location of floral nectar sources by mosquitoes: the long-range responses of <i>Anopheles arabiensis</i> Patton (Diptera: Culicidae) to <i>Achillea millefolium</i> flowers and isolated floral odour | 651 |
| BARKER, G. M., PRESTIDGE, R. A. & POTTINGER, R. P. Reproductive phenology of <i>Listronotus bonariensis</i> (Kuschel) (Coleoptera: Curculionidae) in northern New Zealand pastures | 659 |
| GILBERT, N. & COAKER, T. H. Differential survival of British and Mediterranean strains of <i>Pieris rapae</i> (L.) (Lepidoptera: Pieridae) on different brassica cultivars | 669 |

CONTENTS

PAGE

| | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| McDONALD, G. & SMITH, A. M. Phenological development and seasonal distribution of the Rutherglen bug, <i>Nysius vinitor</i> Bergroth (Hemiptera: Lygaeidae), on various hosts in Victoria, south-eastern Australia | 673 |
| POWELL, W. & WRIGHT, A. F. The abilities of the aphid parasitoids <i>Aphidius ervi</i> Haliday and <i>A. rhopalosiphii</i> De Stefani Perez (Hymenoptera: Braconidae) to transfer between different known host species and the implications for the use of alternative hosts in pest control strategies | 683 |
| AGOUNKÉ, D., AGRICOLA, U. & BOKONON-GANTA, H. A. <i>Rastrococcus invadens</i> Williams (Hemiptera: Pseudococcidae), a serious exotic pest of fruit trees and other plants in West Africa | 695 |
| NARASIMHAM, A. U. & CHACKO, M. J. <i>Rastrococcus</i> spp. (Hemiptera: Pseudococcidae) and their natural enemies in India as potential biocontrol agents for <i>R. invadens</i> Williams | 703 |
| WILLINK, E. & MOORE, D. Aspects of the biology of <i>Rastrococcus invadens</i> Williams (Hemiptera: Pseudococcidae), a pest of fruit crops in West Africa, and one of its primary parasitoids, <i>Gyranusoidea tebygi</i> Noyes (Hymenoptera: Encyrtidae) | 709 |
| HERATH, P. R. J., JAYAWARDENA, K. G. I., HEMINGWAY, J. & HARRIS, J. DDT resistance in <i>Anopheles culicifacies</i> Giles and <i>A. subpictus</i> Grassi (Diptera: Culicidae) from Sri Lanka: a field study on the mechanisms and changes in gene frequency after cessation of DDT spraying | 717 |

DATES OF PUBLICATION IN PARTS

| | |
|---------------------------|------------------|
| PART 1, pp. 1–171 | 19 February 1988 |
| PART 2, pp. 173–363 | 24 May 1988 |
| PART 3, pp. 365–547 | 25 August 1988 |
| PART 4, pp. 549–747 | 18 November 1988 |

ERRATUM (VOL. 77)

p. 658, line 40, for 'rainy season' read 'dry season'.

ERRATA (VOL. 78)

- p. 35, line 4, for 'km' read 'km²'.
- p. 43, line 10, for 'south-west' read 'north-east'.
- p. 148, line 30, after '*morsitans*' add 'Westwood'.
- p. 199, line 2, before '**Hymenoptera**' add '('.
- p. 420, line 27, after '*microplus*' add '(Canestrini)'.
- p. 438, lines 27, 37, 40, 44, 46, delete numbers '217' to '263'.