

*Corresponding author.
doi: 10.1192/j.eurpsy.2021.1696

Introduction: Brain heterotopia represent a group of rare malformations with a heterogeneous phenotype, ranging from asymptomatic to severe clinical picture (resistant epilepsy, severe developmental delay). The etiology is multifactorial, including both genetic and environmental factors.

Objectives: In this paper we present our experience regarding behavior problems in patients with heterotopia.

Methods: A cohort of 16 pediatric patients with brain heterotopia, six females and ten males, with age at last follow-up ranging from 2 months to 24 years were investigated by clinical examination, electroencephalographic studies, brain imaging, and genomic tests. Specific psychological tests and psychiatric evaluation were performed in all children for behavior problems assessment.

Results: Six individuals presented behavioral problems: autism (three patients) and hyperkinesia with attention deficit (three patients). All of them had intellectual disability or learning problems; five patients had epilepsy, with drug-resistant seizures in four cases. In two cases the behavioral problems occurred before the onset of epileptic seizures.

Conclusions: Behavior problems are important features in patients with brain heterotopia, making the management of these patients more difficult, especially when they occur in association with drug-resistant epilepsy. Acknowledgements: This work was supported partially by grants of the Romanian National Authority for Scientific Research and Innovation CCCDI – UEFISCDI, Projects COFUND-ERANET E-RARE 3-HETER-OMICS-2 Number 87/2019 and 88/2019 within PNCDI III.

Disclosure: No significant relationships.
Keywords: hyperkinesia with attention deficit; autism; Epilepsy; brain heterotopia

EPV0090

Suicidal ideation during the first wave of COVID-19 pandemic in a child and adolescent psychiatry emergency care sample

F. Cordeiro*, V. Santos and T. Cartaxo
Child And Adolescent Psychiatry, Pediatric Hospital of Coimbra, Coimbra Hospital and University Centre, Coimbra, Portugal, Coimbra, Portugal

*Corresponding author.
doi: 10.1192/j.eurpsy.2021.1697

Introduction: The COVID-19 pandemic forced many countries to apply restrictive measures. During the first wave Portugal went through a lockdown, and all the child and adolescents had to stay home and could only contact with the one’s they lived with for several months.

Objectives: This study aimed to evaluate the impact of those restrictions on suicidal ideation in the pediatric population evaluated in a child and adolescent psychiatry emergency care of a tertiary referral hospital.

Methods: We conducted an exploratory retrospective study. All the data from discharge notes were collected between March 15th and June 15th of 2020 (n=59), and in the homologous period of the

previous year (n=178). The referral after evaluation (primary care, child and adolescent psychiatry consultation, inpatient unit) was considered a measure of severity.

Results: The demographic variables (sex, age) were homogeneous between the two groups (p ≥ 0,05). 17,4% (n=31) of the sample from 2019, and 16,9% (n=10) of the sample of 2020 had suicidal ideation, which was not statistically different between groups (p=1,000). The referral, after evaluation between groups were also not statistically different (p=0,186).

Conclusions: Even though the proportion of patients with suicidal ideation was homogenous during the two periods, the total number of patients evaluated in the emergency room were lower during the first wave of Covid-19 pandemic. We assume that the population had fear of seeking help in hospital facilities, but we also believe that the pause on school burdens and the reconnection between some families could have function as protective factors.

Disclosure: No significant relationships.
Keywords: Restrictions; pandemic; COVID-19; Suicidal ideation

EPV0091

The prevalence of oppositional defiant disorders among young people in europe: A systematic review and meta-analysis

R. Sacco^{1*}, N. Camilleri² and K. Umla-Runge³
¹Psychiatry, Cardiff University, Teesside University, Malta Mental Health Services, Attard, Malta; ²Child And Young People’s Services, Malta Mental Health Services, Pieta, Malta and ³Psychiatry, Cardiff University, Cardiff, United Kingdom

*Corresponding author.
doi: 10.1192/j.eurpsy.2021.1698

Introduction: This systematic review estimates the pooled prevalence (PP) of oppositional defiant disorders (ODD) among 5-to-18-year-old YP living in Europe, based on prevalence rates established in the last five years (LFY).

Objectives: Trends of prevalence rates across countries, gender and level of education were analysed. The random effects pooled prevalence rate (REPPR) for ODD was calculated.

Methods: A search strategy was conducted on three databases. Studies were also identified from reference lists and grey literature. Eligible studies were evaluated for reliability, validity, bias, and the REPPR for ODD was calculated.

Results: The European REPPR for ODD is calculated at 1.9% (Figure 1). The REPPR among males is 4.8%, whereas the rate among females is 2.7% (95% CI: 0.7%- 1.4%). The prevalence rate of ODD among primary school children is 1.8 times higher than the prevalence of secondary school children (Figure 2).




Figure 1: Forest plot showing the prevalence rates of oppositional defiant disorder