

for depressed fathers through a process evaluation from the perspective of fathers.

Methods. This qualitative study was nested within a cluster randomized controlled trial, with depressed fathers of young children (0–3 years) recruited from 18 towns in Karachi, Pakistan. Face to face or digitally audio-recorded interviews were conducted with depressed fathers (N = 24) from the intervention arm of the trial at end of intervention period (i.e., 4-month post-baseline). The intervention involved twelve sessions of a parenting intervention called Learning Through Play integrated with group Cognitive Behaviour Therapy and manual content from “Focus on Fathers”. Interviews were guided by a semi-structured topic guide to explore perceived usefulness of the intervention with particular focus on exploring any additional benefits or challenges of engaging partners into the intervention. Interviews lasted approximately an hour. Data were analysed using the principles of Framework Analysis. A concurrent analysis of initial interviews directed towards further interviews until data saturation.

Results. Analysis highlighted perceived usefulness of the intervention such as improvement in mood, engagement in routine tasks, healthy thinking patterns, increased attachment with child, improved relationship with the family, improvement in parenting knowledge and more positive attitudes towards child development, supporting partners in household chores, and recommendation to promote this partner inclusive parenting program throughout the country.

Conclusion. Addressing depression in parents is hugely important due to its adverse impact on both parents and children. This low-cost parenting program supported fathers in their parenting role along with improvement in psychological well-being. This has also informed barriers and facilitators to implement the LTP plus parenting program and the possibilities to roll out the intervention at national level.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Cognitive Behavioral Therapy Optimizing Post-Operative Outcomes Among Coronary Artery Bypass Graft Surgery Patients: A Systematic Review

Ms Danya Ibrahim^{1*}, Dr Ibrahim H. Elkhidir^{1,2}, Ms Zainab Mohammed¹, Ms Da'ad Abdalla¹ and Mr Omer A. Mohammed¹

¹Khartoum University, Faculty of Medicine, Khartoum, Sudan and

²Internal medicine, Ribat University Hospital, Khartoum, Sudan

*Presenting author.

doi: 10.1192/bjo.2024.167

Aims. This review aims to evaluate the lasting advantages of cognitive behavioral therapy (CBT) in alleviating anxiety and depression and improving overall health outcomes post-CABG.

Methods. A comprehensive search across databases including Science Direct, PsycINFO, PubMed, Google Scholar, VHL, Cochrane, and Scopus was conducted up to October 2023. The study adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement and Cochrane guidelines. Inclusion criteria involved randomized controlled trials reporting on CBT or CBT-based interventions tailored for CABG patients and control groups had usual care, with anxiety and depression symptoms, as well as quality of life, as primary

outcomes. Outcome variations were analyzed through standard deviation, while efficacy was measured via confidence intervals. Evaluation of the intervention process included examining feasibility, adherence, acceptability, inclusion rates, safety, and tolerability.

Results. Three randomized controlled trials including a pilot study in America and Denmark, encompassing a total of 286 patients, were reported in this review. The participants' pooled mean age was 63.19 years (intervention) and 63.9 years (control), the male-to-female ratio was approximately 2:1 [males n = 174 (intervention n = 101; control = 73) while females n = 91 (intervention n = 60; control n = 31)], and cardiac as well as non-cardiac comorbidities including psychiatric diagnosis have been reported. The array of therapies ranged from education on anxiety and depression management skills to a combination of CBT and supportive stress management (SSM), and psychoeducational interventions paired with physical therapy. Results indicated that both CBT and SSM led to improvements in anxiety and depression symptoms, accompanied by reduced hospital stays, decreased hopelessness, lower scores in dysmorphic mood and irritability, lowered perceived cognitive impairment and stress, and increased satisfaction with therapy. The cognitive behavioral therapy demonstrated commendable feasibility, acceptability, safety, and efficacy, with some concerns raised about potential issues of low adherence.

Conclusion. This systematic review emphasizes the positive impact of CBT on depression, anxiety, and quality of life in individuals post-CABG surgery. Future studies should adopt standardized CBT protocols, comprehensively evaluating CBT's influence on overall patient prognosis, considering cardiovascular outcomes across diverse ethnic groups, exploring cost-effectiveness, and specific patient cohorts that could benefit the most from CBT interventions.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Behavioral Addictions Prevalence and Impact on Medical Sciences Students' Mental Health: A Systematic Review and Meta-Analysis

Mr Omer A. Mohammed¹, Ms Danya Ibrahim^{1*}, Ms Shima Algam Mohamed², Ms Shaza K. Babikir¹ and Dr Ayman Zuhair³

¹Khartoum University, Faculty of Medicine, Khartoum, Sudan;

²University of Al-Neelain, Faculty of Medicine, Khartoum, Sudan and

³Norfolk and Suffolk NHS Foundation Trust, Norwich, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.168

Aims. This study aims to illuminate the prevalence of various behavioral addictions among health professions students and examine their negative effects on mental health.

Methods. In March 2023, a systematic literature search was conducted, encompassing randomized controlled trials, cohort, case-control, and cross-sectional studies from the past five years in PubMed and ScienceDirect. Adhering to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines, 19 papers underwent qualitative analysis, while 15 studies were subjected to quantitative analysis following a quality assessment review.