

## e-Poster viewing: Consultation liaison psychiatry and psychosomatics

### EV0253

#### Sleep disorders associated with interferon in chronic viral hepatitis C

C. Aarab\*, A. Bout, R. Aalouane, I. Rammouz  
 Psychiatrist hospital Ibn Alhassan, psychiatry, FEZ, Morocco  
 \* Corresponding author.

Chronic infection with hepatitis C is a public health problem in Morocco and in the world. The objective of the work screens for sleep problems during the disease and its management to better guide the monitoring and psychiatric interventions.

**Methodology** Sixty-eight patients followed for hepatitis C were recruited. The treatment was based on the combination of pegylated IFNa or standard IFNa and ribavirin for 6 to 12 months. The evaluation was done by a questionnaire containing demographic information, medical and psychiatric history, substance use. Sleep assessment was made by Pittsburgh Sleep Quality Index, and measuring the quality of life by SF36. The assessment before starting treatment and at 1 month and 4 months. Statistical analysis by SPSS 21 software read.

**Results** Fifty-six patients were able to complete the monitoring. 36 women to 20 men. The average age is 59.1 years old (standard deviation 9.88). A quarter of patients still gainfully employed. 64.3% of participants live in couple. Seventy-six point eight percent of patients live in the city. Seven per cent consume tobacco and single patient use cannabis occasionally. Sleep disorders increased from 37.5% before the start of the treatment to 55.4% after its start. While 44.5% have kept a sleep disturbed to 4 months. At first, the quality of life had more impaired scores on psychological than physical scores. The overall score declined significantly way after initiation of treatment (0.04). A larger decline on the psychological component (0.000) while the physical score did not significantly decrease.  
**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.583>

### EV0254

#### The age of accountability of consultation liaison, an example from a big referral hospital

Sobhan Aarabi<sup>1,3,\*</sup>, Seyedeh Elham Sharafi<sup>2</sup>,  
 Mohammad-Hosein Pourgharib Shahi<sup>2</sup>

<sup>1</sup> Neuroscience institute, sports medicine research center, Tehran university of medical sciences, Iran

<sup>2</sup> Psychosomatic research center, Imam Khomeini hospital, Tehran university of medical sciences, Tehran, Iran

\* Corresponding author.

E-mail address: [Sobhan.ae@gmail.com](mailto:Sobhan.ae@gmail.com) (S. Aarabi)

CL is a subspecialty in psychiatry and provides a variety of consultation services including diagnosis, therapeutic and teaching efforts of a psychiatrist in none psychiatric units or hospitals.

**Objectives** This is a retrospective descriptive study that evaluated the consultation liaison (CL) psychiatry services in Imam Khomeini hospital, during two recent years.

**Methods** The information has been obtained from reviews of medical profiles of patients who attended the hospital.

**Results** A total of 681 patients (365 patients were male) received CL services during the study period. The most prevalent diagnosis groups were mood disorder (37.91%), delirium (13.6%) and anxiety (12.64%). Our study revealed that cardiovascular unit, infectious disease unit and general surgery units frequently requested for consultations among all hospital wards. Pediatrics unit had the lowest request rate for psychiatric consultation.

**Conclusion** CL is on the rise in general hospitals, specifically in internal medicine and surgery units which alarm us to pay more attention to preventive strategies focused on the most referred problems including mood disorders, delirium and anxiety disorders in patients who attended general hospitals.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<sup>3</sup> Department of Sports Medicine (<http://smrc.tums.ac.ir>), Tehran University of Medical Sciences (<http://www.tums.ac.ir>), No. 7, Jalale Ale Ahmad Highway, PO Box: 14395-578, Tehran, Iran.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.584>

### EV0255

#### Low frequency of request for liaison psychiatry: A difficulty in the early detection of mental disorders by medical assistants?

M. Aguiar Machado\*, J. Gonçalves Oliveira, E. Oliveira Cunha, E. Vieira de Melo, E. Oliva-Costa  
 Universidade Federal de Sergipe, departamento de medicina, Aracaju-SE, Brazil  
 \* Corresponding author.

**Introduction** The liaison psychiatry (LP) is a feature used by the psychiatrist in order to improve the management of patients with mental suffering and/or mental disorder admitted to general hospital.

**Objectives** To characterize the epidemiological profile of hospitalized patients at the university hospital of the federal university of Sergipe (HU-UFS) submitted to LP.

**Methods** retrospective and observational study, through analysis of medical records of patients admitted in the wards of clinical medicine and surgery from the HU-UFS, in the period from January to December 2015, submitted to LP. The information collected fed a specific questionnaire developed by the authors, intended for research of socio-demographic data and clinical profile.

**Results** the frequency of request for LP was of 3.5%, with the majority of applications was performed by clinical medicine (71.2%), while the surgical clinic was responsible for 28.8%. The main reason for the request of LP was the presence of depressive symptoms (49.1%). There was a predominance of females (52.5%) and the mean age was 45.9 ± 14.6 years.

**Conclusions** The frequency of request for LP was very low, suggesting a difficulty in the early detection of mental disorders by physicians. This finding points to an underreporting of cases, since the prevalence of depressive symptoms in hospitalized patients is over 50% in this institution.

**Disclosure of interest** The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.01.585>

### EV0256

#### Psychosomatic process in patients with dissociative mental disorders

M. Axenov<sup>1,\*</sup>, O. Perchatkina<sup>2</sup>, A. Kostin<sup>1</sup>, V. Nikitina<sup>3</sup>

<sup>1</sup> Mental health research institute–Tomsk national research medical center- Russian academy of sciences, borderline states department, Tomsk, Russia

<sup>2</sup> Mental health research institute–Tomsk national research medical center- Russian academy of sciences, department of coordination of scientific investigations, Tomsk, Russia

<sup>3</sup> Mental health research institute–Tomsk national research medical center- Russian academy of sciences, department of psychoneuroimmunology and neurobiology, Tomsk, Russia

\* Corresponding author.