

Observations of convection in A-type stars

Barry Smalley

Astrophysics Group, School of Chemistry & Physics, Keele University, Staffordshire ST5 5BG, United Kingdom

Abstract. Convection and turbulence in stellar atmospheres have a significant effect on the emergent flux from A-type stars. The recent theoretical advancements in convection modelling have proven to be a challenge to the observers to obtain measurements with sufficient precision and accuracy to permit discrimination between the various predictions.

A discussion of the current observational techniques used to evaluate the various convection theories is presented. These include filter photometry, spectrophotometry, hydrogen lines, and metal lines. The results from these techniques are given, along with the successes and limitations.

Keywords. Convection, turbulence, techniques: photometric

1. Introduction

The gross properties of a star, such as broad-band colours and flux distributions, are significantly affected by the effects of convection in stars later than mid A-type. Consequently, our modelling of convection in stellar atmosphere models can significantly alter our interpretation of observed phenomena.

Convection in stellar atmospheres is usually based on mixing-length theory (MLT) of Böhm-Vitense (1958). In their discussion of the Kurucz (1979) ATLAS6 models Relyea & Kurucz (1978) found discrepancies between theoretical and observed *wby* colours which might be the result of an inappropriate treatment of convection within the models. Subsequently, several attempts have been undertaken to improve the situation. Lester *et al.* (1982), for example, introduced “horizontally averaged opacity” and a “variable mixing length” which improved the match with observed *wby* colours, but did not remove all the discrepancies. The ATLAS9 models (Kurucz 1993) introduced an “approximate overshooting” which has not been without its critics (see Castelli *et al.* 1997 for details).

Canuto & Mazzitelli (1991, 1992) proposed a turbulent model of convection to overcome one of the most basic short-comings of MLT, namely that a single convective element (or “bubble” or “eddy”) responsible for the transport of all the energy due to convection. This new model accounts for eddies of various sizes that interact with each other. The CM convection model was implemented in the ATLAS9 code by Kupka (1996).

Theoretical studies of convection in A-type stars have suggested that convection is inefficient (Freytag *et al.* 1996), and thus has only a very small influence on atmospheric structure (Heiter *et al.* 2002). However, these effects are nonetheless still important and in need of observational confirmation. For the early A-type stars (hotter than ~ 8500 K) convective flux is negligible. It is only from mid A-type and later when convection becomes important, as an extensive convection zone develops in the photosphere and below (e.g. Weiss & Kupka 1999).

Convection in A-type stars poses a challenge to both theorists and observers. In this review I will concentrate on the observational effects of convection and what can be deduced from various types of observations.

2. Colours

Photometric indices are a fast and efficient method for determining approximate atmospheric parameters of stars. For the commonly-used Strömgren *uvby* system a vast body of observational data exists (e.g., Hauck & Mermilliod 1998) which can be used to estimate parameters using calibrated model grids (e.g., Moon & Dworetzky 1985, Lester *et al.* 1986, Smalley & Dworetzky 1995). Conversely, knowing atmospheric parameters from other methods, allows observed colours to be compared to model predictions. This method has been used to compare various treatments of stellar convection.




Figure 1. The variation of m_0 index with $b-y$ showing the sensitivity to microturbulence (ξ_t), mixing-length (l/H) and “approximate overshooting”. At each temperature the model with $\log g = 4.0$, $\xi_t = 2 \text{ km s}^{-1}$ and $l/H = 0.5$ is denoted by a square. The arrows indicate the effect of varying ξ_t , l/H and including overshooting (for $l/H = 1.25$). The Philip & Egret (1980) main-sequence and the Smalley & Kupka (1997) CM relationships are included for reference. Note that the squares do not lie on the CM relationship due to differing $\log g$ values.

Smalley & Kupka (1997) compared the predicted *uvby* colours for the CM model with that from the standard Kurucz (1993) MLT models with and without “approximate overshooting”. Comparison against fundamental T_{eff} and $\log g$ stars revealed that the CM models gave better agreement than MLT without overshooting. Models with overshooting were clearly discrepant. This result was further supported by stars with T_{eff} values obtained from the Infrared Flux Method (IRFM) and $\log g$ values from stellar evolutionary models. However, some discrepancies still remained, including a “bump” around 6500 K in the $\log g$ values obtained for the Hyades and continued problems with the Strömgren m_0 index. Similar conclusions were found by Schmidt (1999) using Geneva photometry.

The m_0 index is sensitive to metallicity and microturbulence, but also to convection efficiency as discussed by Relyea & Kurucz (1978) and Smalley & Kupka (1997 Fig. 6). Inefficient convection (CM and MLT $l/H \sim 0.5$) clearly works in the domain of the A stars down to $b-y \sim 0.20$ (approx 7000 K, F0). For cooler stars, convection becomes more efficient and substantive within the atmosphere and higher values of mixing-length and lower microturbulent velocities would be required to fit the observed m_0 indices of the main sequence (see Fig. 1). The 2d numerical radiation hydrodynamics calculations of Ludwig *et al.* (1999) are useful in this regard. They indicate a rise in mixing-length

from $l/H \sim 1.3$ at 7000 K to $l/H = 1.6$ for the Sun (5777 K), while a much lower $l/H \sim 0.5$ was found for models at 8000 K (Freytag 1995). This is in agreement with that implied by m_0 index. However, around 6000 K there still remains a significant discrepancy, which could only be reduced by invoking the approximate overshooting option (Fig. 1). Thus, none of the convection models used in classical model atmospheres allows for the reproduction of the m_0 index, unless two parameters, i.e., l/H and the amount of “approximate overshooting”, are varied over the H-R Diagram.

3. Fluxes

The observed stellar flux distribution is influenced by the effects of convection on the atmospheric structure of the star. As we have seen with photometric colours, these effects have a clearly observable signature. Hence, high precision stellar flux measurements will provide significant and useful information on convection.




Figure 2. Fluxes for CM models and MLT models, with $l/H = 0.5$ and 1.25 , compared to that for a model with zero convection. At 8000 K the CM and MLT $l/H = 0.5$ models give essentially the same fluxes as zero convection. Note that the fluxes for MLT $l/H = 1.25$ with and without overshooting are almost identical. At 7000 K the differences are more noticeable, especially in the region $4000 \sim 5000 \text{ \AA}$, and the effect of overshooting is now considerable.

Lester *et al.* (1982) presented a study of convective model stellar atmospheres using a modified mixing-length theory. They found small, systematic differences in the optical fluxes. Their figures demonstrate that convection can have a measurable effect on stellar fluxes.

Figure 2 shows the effects of changing mixing length from 0, through 0.5 to 1.25 on the emergent flux for solar-composition models with 8000 K and 7000 K ($\log g = 4.0$ and




Figure 3. The effects of convection on the predicted shape of Balmer profiles for models with $T_{\text{eff}} = 7000$ K, $\log g = 4.0$, $[M/H] = 0.0$ and $\xi_t = 2$ km s $^{-1}$. H α is unaffected by the values of l/H , but sensitive to “approximate overshooting”, while H β is sensitive to both.

$\xi_t = 2$ km s $^{-1}$). At 8000 K both the CM and MLT $l/H = 0.5$ models give essentially the same fluxes as zero convection. At 7000 K, however, the differences are more noticeable, with the effect of overshooting being considerable.

In the ultraviolet the effects are even more significant, but comparison with observations would be complicated by the significant amount of metal line blanketing in this region. In contrast, in the infrared the effects are generally less compared to the optical. Hence, a combination of optical and infrared fluxes should provide a good basis for fixing the effective temperature of the star as well as testing the convective nature of the stellar atmosphere. In fact, the IRFM (Blackwell & Shallis 1977, Blackwell & Lynas-Gray 1994) is a very useful method for determining nearly model-independent T_{eff} for stars and has proved invaluable in our quest for accurate atmospheric parameters.

Unfortunately, very little high-precision stellar spectrophotometry exists. This situation will be rectified, once the ASTRA Spectrophotometer (Adelman *et al.* 2003, 2005) begins operation. This will allow spectrophotometry to be added to our observational diagnostic toolkit.

4. Balmer profiles

The temperature sensitivity of Balmer lines makes them an excellent diagnostic tool for late A-type stars and cooler (Gardiner 2000). However, as emphasised by van’t Veer & Mégessier (1996), the H α and H β profiles behave differently due to convection: H α is significantly less sensitive to mixing-length than H β . Both profiles are affected by the presence of overshooting, with H β being more influenced than H α (see Fig. 3). Since H α is formed higher in the atmosphere than H β , Balmer lines profiles are a very good depth probe of stellar atmospheres. Naturally, Balmer profiles are also affected by microturbulence, metallicity and, for the hotter stars, surface gravity (Heiter *et al.* 2002).

In their comparison of Balmer line profiles, Gardiner *et al.* (1999) found that both CM and MLT without overshooting gave satisfactory agreement with fundamental stars. Overshooting was again found to be discrepant. Intriguingly, they found that while $l/H = 0.5$ was generally preferred, the region between 6000 K and 7000 K required a higher value ($l/H \geq 1.25$). This corresponds to the “bump” region found for Hyades stars using




Figure 4. Variation of microturbulence with effective temperature. Based on results of Gray (2001) for stars with $\log g > 4.0$. Note the apparent relatively abrupt change in behaviour around 6600 K

wavy. However, this is not supported by either Fuhrmann *et al.* (1993) or van't Veer & Mégessier (1996).

Gardiner *et al.* (1999) also found evidence for significant disagreement between all treatments of convection for stars with T_{eff} around 8000 ~ 9000 K. Smalley *et al.* (2002) reviewed this region using binary systems with known $\log g$ values and their revised fundamental T_{eff} values of the component stars. They found that the discrepancy found by Gardiner *et al.* (1999) was no longer as evident. However, this region was relatively devoid of stars with fundamental values of both T_{eff} and $\log g$. Further fundamental stars are clearly required in this region.

One potential difficulty with Balmer profiles is their great width, which can pose problems during the extraction and rectification of observations. Locating the 'true' continuum is non-trivial and a small error can lead to significant errors in any temperature obtained from fitting the profile (Smith & van't Veer 1988).

5. Microturbulence

Microturbulence (long used as a free parameter in abundance analyses) does appear to vary with effective temperature. Chaffee (1970) found that microturbulent velocity (ξ_t) rose from 2 km s^{-1} for early A-type stars up to 4 km s^{-1} for late A-type stars, and then back to 2 km s^{-1} for mid F-type stars and back up to 4 km s^{-1} for solar-type stars. He also found that microturbulence correlated weakly with the Strömrgren m_0 index. Recently, Gray (2001) found that ξ_t varied from around 3 km s^{-1} for mid-A type down to around 1 km s^{-1} for solar-type stars, confirming the results of Couptry & Burkhart (1992) who found a variation from around 0 km s^{-1} for late B-type, up to around 3 km s^{-1} for mid A-type type, down to around 2 km s^{-1} for early F-type stars. Nissen (1981) gave a regression fit to both T_{eff} and $\log g$ for near-main-sequence solar-composition stars, from 2 km s^{-1} for F-type down to 1.5 km s^{-1} for solar-type stars. Fig. 4 shows the variation of ξ_t with T_{eff} for near main-sequence stars ($\log g > 4.0$) based on the results given by Gray (2001).

Velocity fields are present in stellar atmospheres which can be measured using line bisectors (Dravins 1987, Gray 1992). Compared to solar-type stars, the line bisectors are reversed, indicating small rising columns of hot gas and larger cooler downdrafts in A-type stars (Landstreet 1998). It is these motions that are thought to be responsible, at




Figure 5. Schematic summary of typical values of mixing-length and microturbulence found in the literature for various spectral types.

least in part, for the existence of microturbulence. In fact, 3-D numerical simulations of solar granulation can account for observed line profiles without the need for any microturbulence (Asplund *et al.* 2000). Similar results have been found for Procyon (Gray 1985, Allende Prieto 2002), which is also a star with well-known physical parameters (e.g., Kervella *et al.* 2004). Unfortunately, current numerical simulations for A-stars predict bisectors in the opposite direction to that observed (Freytag & Steffen 2005).

6. Böhm-Vitense gap

Böhm-Vitense (1970) found a scarcity of stars around 7000 K which were attributed to the abrupt onset of convection. Observationally, as discussed above, this region corresponds to the *uvby* “bump” which is a mismatch between the $\log g$ from evolutionary models and model atmosphere calculations, an apparent change in behaviour of microturbulence around 6600K, and the suggestions that Balmer profiles require a higher value for mixing-length than other regions. In addition, there is a dip in chromospheric activity in this region (Böhm-Vitense 1995). D’Antona *et al.* (2002) discussed the Böhm-Vitense gaps and the role of turbulent convection, concluding that a gap in the region just below T_{eff} of 7000 K is a T_{eff} rather than just a colour gap. This is an interesting area of the H-R Diagram where the convective core shrinks to become radiative, while the envelope and the atmosphere change from radiative to convective (de Bruijne *et al.* 2001). Chromospheric activity indicators are another useful diagnostic, with another transition regime around 8300 K (Simon *et al.* 2002). All in all, this is an observationally challenging region of the H-R Diagram, since there are both atmospheric and internal effects which we need to differentiate between.

7. Conclusions

Fig. 5 summarizes the typical results of the various observational tests. While there are some contradictions and anomalies the Figure does illustrate the current state of our observational understanding of stellar convection. For stars hotter than A0 there is no convection or significant microturbulence. For the early A-type stars there is essentially no convection within the atmosphere, since the temperature gradient is radiative, but there are velocity fields as indicated by the modest microturbulence values. Velocity fields increase as we go through mid to late A-type stars, and inefficient convection is required within the atmosphere. Once convection becomes efficient (F-type and later) the value of microturbulence is found to drop, while the mixing-length is required to increase (although some evidence points against this, such as the Balmer-line results discussed earlier).

Overshooting is still an issue to be resolved, since there are clearly velocity fields above the convection zone. While the “approximate overshooting” of Kurucz appears to have been discounted by observational tests, there is clearly the need for some sort of overshooting to be incorporated within model atmosphere calculations.

The effects of convection on the stellar atmospheric structure can be successfully probed using a variety of observational diagnostics. The combination of photometric colours and Balmer-line profiles has given us a valuable insight into the nature of convection in A-type stars. High quality observations that are currently available and those that will be in the near future, will enable further refinements in our theoretical models of convection and turbulence in stellar atmospheres.

Acknowledgements

The author gratefully acknowledges the support of PPARC, the Royal Society and the IAU. This research has made use of model atmospheres calculated with ATLAS9 at the Department of Astronomy of the University of Vienna, Austria, available from <http://ams.astro.univie.ac.at/nemo/>.

References

- Adelman S.J., Gulliver A.F., Smalley B., Pazder J.S., Younger P.F., Boyd L. & Epan D., 2003 In: Adelman S.J., Kupka F., Weiss W.W. (eds.) *Model Atmospheres and Spectrum Synthesis*. A.S.P. Conf. Proc. 44, Poster E55.
- Adelman S.J., Gulliver A.F., Smalley B., Pazder J.S., Younger P.F., Boyd L. & Epan D., 2005, *These Proceedings*, JP2
- Allende Prieto C., Asplund M., López, R.J.G., Lambert D.L., 2002, *ApJ* 567, 544
- Asplund M., Nordlund Å., Trampedach R., Allende Prieto C., Stein R.F., 2000, *A&A* 359, 729
- Blackwell D.E., Lynas-Gray A.E., 1994, *A&A* 282, 899
- Blackwell D.E., Shallis M.J., 1977, *MNRAS* 180, 177
- Böhm-Vitense E., 1958, *ZfA* 46, 108
- Böhm-Vitense E., 1970, *A&A* 8, 283
- Böhm-Vitense E., 1995, *A&A* 297, L25
- Canuto V.M., Mazzitelli I., 1991, *ApJ* 370, 295
- Canuto V.M., Mazzitelli I., 1992, *ApJ* 389, 724
- Canuto V.M., Goldman I., Mazzitelli I., 1996, *ApJ* 473, 550
- Castelli F., Gratton R.G., Kurucz R.L., 1997, *A&A* 318, 841
- Chaffee F.H., 1970, *A&A* 4, 291
- Coupry M.F., Burkhardt C., 1992, *A&AS* 95, 41
- D’Antona F., Montalbán J., Kupka F., Heiter U., 2002, *ApJ* 564, L93
- de Bruijne J.H.J., Hoogerwerf R., de Zeeuw P.T., 2001, *A&A* 367, 111
- Dravins D., 1987, *A&A* 172, 200
- Freytag B., 1995, *Ph.D. Thesis*, Universität Kiel
- Freytag B., Ludwig H.-G., Steffen M., 1996, *A&A* 313, 497
- Freytag B., Steffen, M. 2005, *These Proceedings*, 139
- Fuhrmann K., Axer M., Gehren T., 1993, *A&A* 271, 451
- Gardiner R.B., 2000, *Ph.D. Thesis*, University of Keele
- Gardiner R.B., Kupka F., Smalley B., 1999, *A&A* 347, 876
- Gray D.F., 1985, *ApJ* 255, 200
- Gray D.F., 1992, *The observation and analysis of stellar photospheres*, (2nd ed.), Cambridge University Press
- Gray R.O., Graham P.W., Hoyt S.R., 2001, *AJ* 121, 2159
- Hauck B., Mermilliod M., 1998, *A&AS* 129, 431
- Heiter U., Kupka F., van’t Veer-Menneret C., Barban C., Weiss W.W., Goupil M.-J., Schmidt W., Katz D., Garrido R., 2002, *A&A* 392, 619
- Kervella P., Thévenin F., Morel P., Berthomieu G., Bordé P., Provost J., 2004, *A&A* 413, 251

- Kupka F., 1996. In: Adelman S.J., Kupka F., Weiss W.W. (eds.) *Model Atmospheres and Spectrum Synthesis. ASP. Conf. Proc.* 44, p. 356
- Kurucz R.L., 1979, *ApJS* 40, 1
- Kurucz R.L., 1993, Kurucz CD-ROM 13: ATLAS9, SAO, Cambridge, USA
- Landstreet J.D., 1998, *A&A* 338, 1041
- Lester J.B., Gray R.O., Kurucz R.L., 1986, *ApJS* 61, 509
- Lester J.B., Lane M.C., Kurucz R.L., 1982, *ApJ* 260, 272
- Ludwig H.-G., Freytag B., Steffen M., 1999, *A&A* 346, 111
- Moon T.T., Dworetsky M.M., 1985, *MNRAS* 217, 305
- Nissen P.E., 1981, *A&A* 97, 145
- Philip A.G.D., Egret D., 1980, *A&AS* 40, 199
- Relyea L.J., Kurucz R.L., 1978, *ApJS* 37, 45
- Schmidt W., 1999, *Master's Thesis*, Johannes Kepler University Linz, Austria
- Simon T., Ayres T.R., Redfield S., Linsky J.L., 2002, *ApJ* 579, 800
- Smalley B., Dworetsky M.M., 1995, *A&A* 293, 446
- Smalley B., Kupka F., 1997, *A&A* 328, 439
- Smalley B., Gardiner R.B., Kupka F., Bessell M.S., 2002, *A&A* 395, 601
- Smith K.C., van't Veer C., 1988 in *Elemental Abundance Analyses*, Adelman S.J., Lanz T. (eds.), Institut d'Astronomie de l'Université de Lausanne, p. 133
- van't Veer C., Megessier C., 1996, *A&A* 309, 879
- Weiss W.W., Kupka F., 1999 In: Giménez Á., Guinan E.F., Montesinos B. (eds.), *Theory and Tests of Convection in Stellar Structure, ASP Conf. Proc.* 173, p. 21

Discussion

GREVESSE: Turbulence in the Sun not only shows up in the asymmetries of the lines through the shapes of the bisectors but also in a systematic but small blue shift of the centre of the line which is related to the depth of formation of the line.

KUPKA: In stellar evolution, “alpha” is tuned to match the radius of the present Sun. Now it may be calibrated by numerical simulations. But although such an MLT calibrated model matches the right entropy jump, its temperature gradient is likely to be wrong (and it will actually continue to fail in helioseismological tests), as will be the radiation field, and the colours computed from it. Thus, just tuning the alpha value with simulations will not help. You can only get one quantity at and perhaps near the calibration point. It does not prevent us from finding a better model, and the alpha to match a particular quantity may be whatever in a particular case.

SMALLEY: Indeed, from a purely observational viewpoint we could use a negative mixing-length if that fitted the observational data better!

TRAMPEDACH: Overshooting is not contentious, but it does transport a negative convective flux. Kurucz' s ‘overshoot’ is a flux-smoothing scheme, that results in a positive convective flux in the stable layers adjacent to a convection zone.

KUPKA ON TRAMPEDACH'S REMARK: On the overshooting: Yes, I agree that is what is found in numerical simulations, in refined non-local models, and also in geophysical measurements of convection zones in geophysical convection zones. But they all usually also find a small zone where $\nabla - \nabla_{\text{ad}} < 0$ while $F_c > 0$ which is what has been smoothed out by Kurucz. For 1D models it was just a fitting parameter and having a negative flux instead does not resolve the problems one finds for these models when comparing them to observations. In this sense one still has the problem of how to carry over these findings to single models and reproduce the observations without further tuning.