

## STRICTLY LOCALIZABLE MEASURES

P. JIMÉNEZ GUERRA AND B. RODRÍGUEZ-SALINAS

### Introduction

In this paper it is proved that every locally strictly localizable Radon measure of type  $(\mathcal{H})$ , is strictly localizable, from where it follows immediately the existence of lifting for these measures.

R. Ryan states in [9] that a complete measure has a lifting if and only if it is strictly localizable. The existence of lifting for the Lebesgue measure in  $\mathbf{R}^n$  has been proved by von Neumann [4] and for general  $\sigma$ -finite measures by D. Maharam [3]. A. and C. Ionescu Tulcea [2] have proved the existence of lifting for positive Radon measures in locally compact spaces, and L. Schwartz [10] has solved the problem for locally finite Radon measures (of type  $(\mathcal{H})$ ) in arbitrary topological Hausdorff spaces.

B. Rodríguez-Salinas and P. Jiménez Guerra [7] and [8] have proved that every locally  $\sigma$ -finite Radon measure of type  $(\mathcal{H})$  is strictly localizable, result which is an immediate consequence of the Maharam's theorem and of the theorem 2 in this paper (see Corollary 3).

Proposition 4 allows to extend, for locally strictly localizable Radon measures of type  $(\mathcal{H})$ , many results which are known for finite Radon measures of type  $(\mathcal{H})$ .

The results concerning the existence of different types of liftings for locally  $\sigma$ -finite Radon measures of type  $(\mathcal{H})$ , that were obtained by Rodríguez-Salinas in [6], can be easily extended for locally strictly localizable Radon measures of type  $(\mathcal{H})$ , using Theorem 2 and Proposition 4 of this work.

### Notations and fundamentals

We will denote by  $E$  an arbitrary topological space (Hausdorff or not) and by  $\mathcal{H}$  a class of closed subsets of  $E$ . If  $\mu$  is a Radon measure of

---

Received July 16, 1979.

type  $(\mathcal{H})$  on  $E$  and  $A \subset E$  we will denote by  $\mu_A$  the Radon measure of type  $(\mathcal{H}_A)$  on  $A$ , induced by the measure  $\mu$  (see Theorem 78 of [7]).

By  $\mu$ -compact set and Radon measure of type  $(\mathcal{H})$  we will understand the same as in [5].

A Radon measure  $\mu$  of type  $(\mathcal{H})$  on a topological space  $E$  is strictly localizable (Definition 8, p. 16 and 17 of [2]) if and only if there exists a family  $\mathcal{C}$  of  $\mu$ -measurable disjoint subsets of  $E$ , with positive and finite measure which verify one of the two following equivalent conditions:

$M_1$ .  $\sup \{\tilde{K} : K \in \mathcal{C}\} = E$  (where  $\tilde{K}$  is the equivalence class of the set  $K$  with respect to the equivalence relation:

$$A \equiv B \Leftrightarrow \mu'(A \triangle B) = 0,$$

being  $A$  and  $B$   $\mu'$ -measurable subsets of  $E$ ).

$M_2$ . For every set  $A \subset E$  with  $\mu'(A) < +\infty$ , there is a countable subset  $\mathcal{C}_A \subset \mathcal{C}$  such that  $A - \bigcup_{K \in \mathcal{C}_A} K$  is  $\mu'$ -negligible.

From now on we will say that  $\mathcal{C}$  is a family of strict localizability for  $\mu$  and we will denote by  $\bar{\mathcal{C}}$  the set  $\bigcup_{K \in \mathcal{C}} K$ .

**LEMMA 1.** *If  $\mu$  is a locally strictly localizable Radon measure of type  $(\mathcal{H})$  on  $E$ ,  $G$  is an open subset of  $E$  such that  $\mu(E - G) > 0$  and  $\mathcal{C}$  is a family of strict localizability for  $\mu_G$ , then there exists an open subset  $G'$  of  $E$  and a family  $\mathcal{C}'$  of strict localizability for  $\mu_{G'}$ , such that  $\mathcal{C} \subset \mathcal{C}'$  and  $G$  is strictly contained in  $G'$ .*

*Proof.* We have that  $\mu(E - G) > 0$ , then there exists a set  $H \in \mathcal{H}$  of measure  $\mu(H) > 0$ , such that  $H \subset E - G$ . Since  $H$  is  $\mu$ -compact,  $\mu$  is locally strictly localizable and  $\mu(H) > 0$ , it is easily deduced the existence of an open subset  $U$  of  $E$  such that  $\mu_U$  is strictly localizable and  $\mu(U \cap H) > 0$ . Evidently,  $G$  is strictly contained in  $G' = G \cup U$ .

Let  $\mathcal{D}$  be a family of strict localizability for  $\mu_U$ . For every subset  $\mathcal{S} \subset \mathcal{D}$  we set

$$\mathcal{S}' = \{K - G : K \in \mathcal{S}\}$$

and

$$\mathcal{S}'' = \{K' \in \mathcal{S}' : \mu(K') > 0\}.$$

We will prove now that  $\mathcal{C}^* = \mathcal{C} \cup \mathcal{D}''$  is a family of strict localizability for  $\mu_{G'}$  for which it is enough to verify that  $\mathcal{C}^*$  satisfies  $M_2$ .

If  $A \subset G'$  and  $\mu_{G'}(A) < +\infty$  then  $\mu_G(A \cap G)$  and  $\mu_U(A \cap U)$  are finite

and there exist two countable subsets  $\mathcal{C}_A \subset \mathcal{C}$  and  $\mathcal{D}_A \subset \mathcal{D}$  such that

$$\mu_G(A \cap G - \bar{\mathcal{C}}_A) = 0$$

and

$$\mu_U(A \cap U - \bar{\mathcal{D}}_A) = 0.$$

So,  $\mathcal{C}_A^* = \mathcal{C}_A \cup \mathcal{D}'_A$  is a countable subfamily of  $\mathcal{C}^*$  which verifies:

$$\begin{aligned} \mu_{G'}(A - \bar{\mathcal{C}}_A^*) &\leq \mu_{G'}(A \cap G - \bar{\mathcal{C}}_A^*) + \mu_{G'}[A \cap (U - G) - \bar{\mathcal{C}}_A^*] \\ &\leq \mu_{G'}(A \cap G - \bar{\mathcal{C}}_A) + \mu_U[A \cap (U - G) - \bar{\mathcal{D}}'_A] \\ &\leq \mu_{G'}(A \cap G - \bar{\mathcal{C}}_A) + \mu_U[A \cap (U - G) - \bar{\mathcal{D}}_A] \\ &\leq \mu_{G'}(A \cap G - \bar{\mathcal{C}}_A) + \mu_U(A \cap U - \bar{\mathcal{D}}_A) \\ &= 0 \end{aligned}$$

and, consequently,  $\mathcal{C}^*$  verifies  $M_2$  and the lemma is proved because  $\mathcal{C} \subset \mathcal{C}^*$  by construction.

It should be notice that it follows from  $M_2$  that for every  $H \in \mathcal{H}$  there exists a family  $\mathcal{S}_A \subset \mathcal{D}_A$  such that

$$\mu(A \cap U \cap H - \bar{\mathcal{S}}_A) = 0$$

and

$$\begin{aligned} \mu_U[A \cap (U - G) \cap H \cap \bar{\mathcal{S}}_A] &= \mu_U[A \cap (U - G) \cap H \cap \bar{\mathcal{S}}'_A] \\ &= \sum_{K \in \mathcal{S}'_A} \mu_U[A \cap (U - G) \cap H \cap K] \\ &= \sum_{K \in \mathcal{S}''_A} \mu_U[A \cap (U - G) \cap H \cap K] \\ &= \mu_U[A \cap (U - G) \cap H \cap \bar{\mathcal{S}}''_A], \end{aligned}$$

therefore the inequality

$$\mu_U[(A \cap (U - G) - \bar{\mathcal{D}}''_A) \cap H] \leq \mu_U[(A \cap (U - G) - \bar{\mathcal{D}}_A) \cap H]$$

holds, and it follows from Theorem 74.2 of [7] that

$$\mu_U[A \cap (U - G) - \bar{\mathcal{D}}''_A] \leq \mu_U[A \cap (U - G) - \bar{\mathcal{D}}_A].$$

**THEOREM 2.** *Every locally strictly localizable Radon measure of type  $(\mathcal{H})$  on  $E$ , is strictly localizable.*

*Proof.* Let  $\mu$  be a locally strictly localizable Radon measure of type  $(\mathcal{H})$  on  $E$  and let us consider the set  $\mathcal{A}$  of all pairs  $(G, \mathcal{C})$  where  $G$  is an open subset of  $E$ , such that  $\mu_G$  is strictly localizable and  $\mathcal{C}$  is a family

of strict localizability for  $\mu_G$ . We consider in  $\mathcal{A}$  the following order:

$$(G_1, \mathcal{C}_1) \leq (G_2, \mathcal{C}_2) \Leftrightarrow G_1 \subset G_2 \quad \text{and} \quad \mathcal{C}_1 \subset \mathcal{C}_2.$$

We will see that if  $\{(G_i, \mathcal{C}_i)\}_{i \in I}$  is a chain in  $(\mathcal{A}, \leq)$  then  $\mathcal{C} = \bigcup_{i \in I} \mathcal{C}_i$  is a family of strict localizability for  $\mu_G$ , being  $G = \bigcup_{i \in I} G_i$ , and therefore  $(\mathcal{A}, \leq)$  is inductive.

If  $A \subset G$  and  $\mu_G(A) < +\infty$  then  $A$  is  $\mu_G$ -compact and there is a countable subset  $I'$  of  $I$  such that

$$\mu_G(A - \bigcup_{i \in I'} G_i) = 0.$$

For every  $i \in I'$  we have that  $\mu_{G_i}(A \cap G_i) < +\infty$  and there exists a countable subfamily  $\mathcal{C}_i^*$  of  $\mathcal{C}_i$  such that

$$\mu_{G_i}(A \cap G_i - \bar{\mathcal{C}}_i^*) = 0$$

holds. Consequently  $\mathcal{C}^* = \bigcup_{i \in I'} \mathcal{C}_i^*$  is a countable subset of  $\mathcal{C}$  such that

$$\begin{aligned} \mu_G(A - \bar{\mathcal{C}}^*) &= \mu_G[(A \cap \bigcup_{i \in I'} G_i) - \bar{\mathcal{C}}^*] \\ &\leq \sum_{i \in I'} \mu_{G_i}(A \cap G_i - \bar{\mathcal{C}}_i^*) \\ &= 0 \end{aligned}$$

and  $M_2$  holds. Therefore  $\mathcal{C}$  is a family of strict localizability for  $\mu_G$  and  $(G, \mathcal{C}) \in \mathcal{A}$ .

From Zorn's axiom it is deduced the existence of a maximal element  $(G, \mathcal{C}) \in \mathcal{A}$  and it follows from Lemma 1 that  $E - G$  is  $\mu$ -negligible.

**COROLLARY 3.** *Every locally  $\sigma$ -finite Radon measure of type  $(\mathcal{H})$  on  $E$  is strictly localizable.*

*Proof.* It is an immediate consequence of Theorem 2, because every  $\sigma$ -finite measure is strictly localizable.

**PROPOSITION 4.** *Let  $\mu$  be a Radon measure of type  $(\mathcal{H})$  on  $E$  and  $\mathcal{C}$  a family of strict localizability for  $\mu$ , then we have:*

4.1. *If  $A \subset E$  is such that  $A \cap K$  is  $\mu$ -negligible for all  $K \in \mathcal{C}$ , then  $A$  is  $\mu$ -negligible.*

4.2. *If  $A \subset E$  is such that  $A \cap K$  is  $\mu_K$ -measurable for all  $K \in \mathcal{C}$ , then  $A$  is  $\mu$ -measurable.*

*Proof.* 4.1. For every  $H \in \mathcal{H}$  there exists a countable subclass  $\mathcal{C}_H$  of  $\mathcal{C}$  such that  $\mu(H - \bar{\mathcal{C}}_H) = 0$  and

$$\begin{aligned} \mu(A \cap \bar{\mathcal{C}} \cap H) &\leq \sum_{K \in \mathcal{C}_H} \mu(A \cap K \cap H) \\ &= 0 \end{aligned}$$

holds. Therefore it follows from Theorem 74.2 of [7] that

$$\begin{aligned} \mu(A \cap \bar{\mathcal{C}}) &= \sup \{ \mu(A \cap \bar{\mathcal{C}} \cap H) : H \in \mathcal{H} \} \\ &= 0 \end{aligned}$$

and  $\mu(A) = 0$ .

4.2. For every  $H \in \mathcal{H}$  there exists a countable subclass  $\mathcal{C}_H$  of  $\mathcal{C}$  such that  $\mu(H - \bar{\mathcal{C}}_H) = 0$ . Consequently,

$$\begin{aligned} \mu(H) &= \mu(H \cap \bar{\mathcal{C}}_H) \\ &= \sum_{K \in \mathcal{C}_H} \mu(H \cap K) \\ &= \sum_{K \in \mathcal{C}_H} [\mu(H \cap K \cap A) + \mu((H - A) \cap K)] \\ &= \mu(H \cap A) + \mu(H - A) \end{aligned}$$

and it follows from Theorem 75.2 of [7] that  $A$  is  $\mu$ -measurable.

*Remark 5.* If  $\mu$  is a Radon measure of type  $(\mathcal{H})$  and  $\mathcal{C}$  is a family of strict localizability for  $\mu$ , then there exists a family  $\mathcal{C}'$ , of strict localizability for  $\mu$ , such that  $\mathcal{C}' \subset \mathcal{H}$  and every  $K' \in \mathcal{C}'$  is contained in some  $K \in \mathcal{C}$ ,

### REFERENCES

- [ 1 ] Chatterji, S. D., Desintegration of measures and liftings, Proc. of the Symposium on vector and operator valued measures and applications, Acad. Press, New York (1973), 69–83.
- [ 2 ] Ionescu Tulcea, A. and C., Topics in the theory of lifting, Springer, Berlin, 1969.
- [ 3 ] Maharam, D., On a theorem of von Neuman, Proc. Amer. Math. Soc., **9** (1958), 187–994.
- [ 4 ] Neumann von, J., Algebraische Repräsentanten der Funktionen bis auf eine Menge von Masse Null, J. Crelle, **165** (1931), 109–115.
- [ 5 ] Rodríguez-Salinas, B., Teoría de la medida sobre los espacios topológicos no localmente compactos, Rev. Mat. Hispano-Amer., (4), **32** (1973), 257–274.
- [ 6 ] —,  $\mu$ -espacios de Suslin y Lusin. Propiedad del lifting fuerte, Rev. R. Acad. Ci. Madrid, **72** (1978), 541–557.
- [ 7 ] Rodríguez-Salinas, B. and P. Jiménez Guerra, Medidas de Radon de tipo  $(\mathcal{H})$  en espacios topológicos arbitrarios, Mem. R. Acad. Ci. Madrid, t. X, 1979.
- [ 8 ] —, Espacios de Radon de tipo  $(\mathcal{H})$ , Rev. R. Acad. Ci. Madrid, **69** (1975), 761–774.
- [ 9 ] Ryan, R., Representative sets and direct sums, Proc. Amer. Math. Soc., **15** (1964), 387–390.

- [10] Schwartz, L., Radon measures on arbitrary topological spaces and cylindrical measures, Oxford University Press, 1973.
- [11] Segal, I. E., Equivalences of measure spaces, Amer. J. Math., **73** (1951), 275–313.

*Dpto. de Teoría de Funciones  
Facultad de C. Matemáticas  
Universidad Complutense  
Madrid-3 (SPAIN)*