The Journal of Laryngology and Otology

A. LOGAN TURNER AND J. S. FRASER

Contents

DISPLACEMENT OF THE TREATMENT OF AT	ANTRO-NASAL WAI	LL IN THE	W. S. SYME
T. DIRECT DEMOVAL		LLOMATA-	
THROAT AND EAR DEF	ECTS FROM THE STAT	NDPOINT OF	ELEANOR LOWRY
A NOTE ON ADENOIDEC	TOMY	1.	F. PEARCE STURM
THE CONTROL OF IMME	DIATE BLEEDING IN TENOIDECTOMY .	THE OPERA-	JAMES KERR LOVE
How DO THE LABSTRINT	H FISTULA SYMPTOM	s Arise? .	S. H. MYGIND
CLINICAL RECORD:— FACIAL PARALYSIS	Associated with E.	AR DISEASE	T. A. MACGIBBON
SOCIETIES' PROCEEDINGS ROYAL SOCIETY OF BRITISH MEDICAL AND OTOLOGY.	:— MEDICINE—SECTION ASSOCIATION, BRADE	of Laryngo	ology. N OF LARYNGOLOGY
Abstracts:— EAR.	PHARYNX.	LAF	tynx.
REVIEWS OF BOOKS.	LETTER TO THE E	DITORS.	GENERAL NOTES.

Edinburgh: Oliver and Boyd London: 33 Paternoster Row

Monthly, 4/- net

Annual Subscription, 40/- net

"We have experimentally proved that Listerine is a powerful antiseptic, preventing the development of bacteria and decomposition of vegetable infusions."—British Medical Journal, May 3, 1890.

LISTERINE

A safe, non-poisonous, unirritating antiseptic solution

- LISTERINE is particularly useful in the treatment of abnormal conditions of the mucosa, and admirably suited for a wash, gargle, or douche, in catarrhal conditions of the nose and throat.
- LISTERINE in proper dilution may be freely and continuously used without prejudicial effect, either by injection or spray, in all the natural cavities of the body.
- LISTERINE embodies a two-fold antiseptic effect, in that after the evaporation of its volatile constituents—thyme, eucalyptus, mentha, gaultheria, and ethyl alcohol—a film of boracic and benzoic acids remains upon the surface to which Listerine has been applied, affording more prolonged antiseptic protection.
- LISTERINE is a trustworthy surgical dressing; it has no injurious effect upon the tissues in which the healing process is going on.
- LISTERINE in teaspoonful doses will often afford relief in fermentative dyspepsia, and is largely prescribed, with excellent results, in the various forms of diarrheea, occurring in infants and adults.
- LISTERINE literature, including special pamphlets upon Respiratory Diseases, may be had, by physicians, upon application to

LAMBERT PHARMACAL COMPANY

2101 Locust Street

St Louis, Mo., U.S.A.

British Agents: S. Maw, Son & Sons, 7 Aldersgate St., London, E.C.

The Journal of Laryngology and Otology

(Founded in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY

A. LOGAN TURNER AND J. S. FRASER

ASSISTED BY

DOUGLAS GUTHRIE AND IRWIN MOORE

- 1. Original Articles are accepted on the condition that they have not been published elsewhere.
- 2. Manuscripts should be typewritten, on one side only of the paper, and well spaced.
- 3. Galley proofs and engraver's proofs of illustrations are sent to the author. Corrections, which should be kept to a minimum, must be clearly marked and no additional matter should be added.
- 4. The order for reprints should be sent when returning galley proofs, and for this purpose special forms are supplied, on which the price of the reprints is stated.
- 5. Authors of Original Communications on Oto-Laryngology in other Journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.
- **6.** Editorial Communications should be addressed to "EDITORS, *Journal of Laryngology*, c/o Messrs OLIVER AND BOYD, Tweeddale Court, Edinburgh."
- 7. The Annual Subscription is Forty Shillings, post free, and is payable in advance.
- 8. Single copies of back numbers, both of the present and the previous series, are on sale at Four Shillings each.
- 9. An Annual Subscription can be commenced with any number of the Journal, and the preceding numbers of the current year can be purchased at Four Shillings each.

UNITED STATES OF AMERICA


Orders for this Journal, for single parts or yearly subscriptions, can be sent either through a local bookseller, or to Messrs G. E. STECHERT & Co., 31-33 East 10th Street, New York.

GRANDS PRIX:

Paris, 1900. Brussels, 1910. Buenos Aires, 1910.




Gold Medal, Allahabad, 1910.



A NEW TRACHEOTOMY TUBE

(Pilot permits breathing during introduction)
By Dr A. CUBLEY

Vide Journal of Laryngology and Otology, March 1922.



Needle for Suturing the Pillars of the Fauces, and Forceps for Carrying the Suture.

By Mr THOS. GUTHRIE, F.R.C.S.

Vide Journal of Laryngology and Otology, April 1920.

Manufactured only by

DOWN BROS., LTD.

Surgical Instrument Makers

21 and 23 ST THOMAS'S STREET, LONDON, S.E. 1

Telegrams
"DOWN, LONDON."

(OPPOSITE GUY'S HOSPITAL)

Telephone

HOP 4400 (4 lines)

JUST PUBLISHED. Demy 8vo, 424 pp. With 222 Text Illustrations and 12 Plates, of which 8 are in Colour. 20s. net, Postage 9d.

DISEASES OF THE NOSE, THROAT, AND EAR

FOR PRACTITIONERS AND STUDENTS

Edited by

A. LOGAN TURNER, M.D., F.R.C.S. (Ed.)

Senior Lecturer on Diseases of the Ear, Nose, and Throat, University of Edinburgh With the collaboration of

J. S. FRASER, M.B., F.R.C.S. (Ed.)

J. D. LITHGOW, M.B., F.R.C.S. (Ed.). G. EWART MARTIN, M.B., F.R.C.S. (Ed.).

W. T. GARDINER, M.C., M.B., F.R.C.S. (Ed.). Lecturers on Diseases of the Ear, Nose, and Throat, University of Edinburgh

and DOUGLAS GUTHRIE, M.D., F.R.C.S. (Ed.)

Lecturer on Diseases of the Ear, Nose, and Throat, School of Medicine of the Royal Colleges, Edinburgh

FOR a number of years, the volume on Diseases of the Throat, Nose, and Ear, written by the late Dr W. G. Porter, supplied a much-felt want, and the fact that it passed through three editions in a comparatively short space of time, was sufficient testimony of its popularity.

Seven years have now elapsed since Dr Porter fell in the service of his country, and the present volume, dedicated to his memory, is produced as the joint work of those engaged in the teaching and practice of the specialty in the Edinburgh Medical School.

While the writers have endeavoured to preserve the main object underlying Dr Porter's original book, the subject matter has been expanded along new lines, which may be justly regarded as increasing the usefulness of the volume. For this purpose, an illustrated description of the clinical anatomy of the various regions has been introduced. A further addition will be found in the detailed account given of many of the minor and major operations. A section on Endoscopy has also been added.

A special feature has been made of the illustrations and to the second of the subject was a special feature has been made of the illustrations and the second of the subject was a special feature has been made of the illustrations and the second of the subject was a special feature has been made of the illustrations and the second of the subject was a special feature has been made of the illustrations and the second of the subject was a second of the subject was a special feature has been made of the subject was a subject to the second of the subject was a subject to the second of the subject was a subject to the subject was a subject to the second of the subject was a subject to the subject wa

A special feature has been made of the illustrations, and no trouble has been spared to make these at once informative and exhaustive. It has been found necessary to almost treble them in number. Student and Practitioner alike should find it a practical, comprehensive, and up-to-

BRISTOL: JOHN WRIGHT & SONS LTD. LONDON: SIMPKIN & CO. LTD.

Central London Throat, Nose, and Ear Hospital

GRAY'S INN ROAD

(Near King's Cross Stations) DAILY CLINICS.

OVER 11,000 NEW PATIENTS LAST YEAR. 50,000 ATTENDANCES. IN-PATIENT DEPARTMENT, 43 BEDS.

CLINICAL ASSISTANTS

Are appointed from among those gentlemen who have enrolled themselves for attendance on the practice of the Hospital.

ATTENDANCE OF STAFF

				OP. CLINIC.	IP. OPERATIONS.
Dr Abercrombie	••		••	Thursday, 2 P.M.	Wednesday, 2 P.M.
Dr Wylie			••	Wednesday, 2 P.M.	Thursday, 2.30 p.m.
Dr Dan M'Kenzie				Monday, 2 P.M.	Tuesday, 2.30 P.M.
Mr Harold Kisch				Friday, 2 P.M.	Monday, 2 p.m.
Mr Nicol Rankin				Tuesday, 5 P.M.	•••
Mr Archer Rylani	o			Saturday, 2 P.M.	••
Mr Watkyn-Thomas	3	• •		Thursday, 2 P.M.	••
Mr Gill-Carey		••		Wednesday, 2 P.M.	••
Mr Armour Brown		••		Monday, 2 P.m.	••

FEES

For attendance on the practice of the Hospital-

. 5 guineas | Six months . . . 8 guineas Three months Perpetual ticket . . . 10 guineas

Further particulars on application to

HAROLD KISCH, F.R.C.S., Dean.

RICHARD KERSHAW, Secretary-Superintendent.

CONTENTS

										PAGE
DISPLACEMENT ATROPHIC R									OF .	481
THE DIRECT REI				AL PAP	ILLOMA	та—A S	SIMPLE •	TECH	NIC.	485
THROAT AND EA SCHOOL CHI										488
A NOTE ON ADE						• •				491
THE CONTROL O	r In	MEDIATI	E BLE	EDING I	N THE	,	•	F Ton	ISIL-	• • •
ADENOIDECT	OMY	. James	Kerr	Love, N	A.D.	•		•		493
How DO THE L							S. H	[. Myg	gind,	0
M.D., Copeni	_	n .	•	•	•	•	•	•	•	498
CLINICAL RECOR	D—									
Facial Paraly	sis A	Associate	d with	Ear Di	sease.	T. A. M	acGib	bon, M	I.D.,	
Christch	urch,	, N.Z.		•	•	•				502
SOCIETIES' PROC	EED:	INGS								
Royal Society	of :	Medicine-	—Sect	ion of L	aryngol	logy				503
British Med	ical	Associat	ion. I	Bradford	-Secti	on of 1	Larvne	ology	and	
Otology	•									515
ABSTRACTS—										
Ear .										523
Pharynx	•	•	•	•	•	•	•	•	•	525
Larynx .	·					·				527
REVIEWS OF BOO)KS						_			529
LETTER TO THE		mone.	•	•	•	•	·	-	•	•
		TOKS	•	•	•	•	٠	•	•	534
GENERAL NOTES		•	•	•						535

HEMOPLASTIN

(HÆMOSTATIC SERUM)

Remarkably efficacious in all forms of hæmorrhage

HEMOPLASTIN is a sterile fluid derived from normal serum of horses and cattle. It is composed principally of prothrombin and thrombokinase, together with a third substance known as anti-antithrombin. These constituents are in physiologically balanced solution ready for immediate use. It retains its properties for about three years.

HEMOPLASTIN is indicated in the treatment of all types of hæmorrhage, particularly those depending upon imperfect coagulation of the blood. It has proved of great value in pulmonary hæmorrhage, and in hæmorrhage incidental to surgical procedures, tonsillectomy, etc. Its administration to hæmophiliac subjects about to undergo operations is a valuable prophylactic measure.

HEMOPLASTIN may be administered intravenously or subcutaneously. The dose is 1 to 2 c.c. every 4 to 6 hours until the hæmorrhage is controlled. The prophylactic dose is 2 c.c. one hour before the operation. Hemoplastin may also be applied on sterile gauze to bleeding surfaces.

Hemoplastin (P., D. & Co.) is supplied in bulbs containing 2 c.c.

Parke, Davis & Co., 50-54 Beak St., London, W.1