

INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, <http://www.editorialmanager.com/psm/>. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

1. S.I. units should be used throughout in text, figures and tables.
2. Authors should spell out in full any abbreviations used in their manuscripts.
3. Foreign quotations and phrases should be followed by a translation.
4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* **286**, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* **32**, 173–179.

Cleckley HJ (1941). *The Mask of Sanity*, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank.org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq_tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. **Line artwork:** Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; **Combination artwork (line/tone):** Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; **Black and white halftone artwork:** Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; **Colour halftone artwork:** Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. *Tables* Tables should be numbered consecutively in the text in Arabic numerals and each typed on a separate sheet after the References section. Titles should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

REVIEW ARTICLE

Childhood bullying and the association with psychosis in non-clinical and clinical samples: a review and meta-analysis

van Dam DS, van der Ven E, Velthorst E, Selten JP, Morgan C & de Haan L 2463

ORIGINAL ARTICLES

Childhood trauma and auditory verbal hallucinations

Daalman K, Diederens KJM, Derks EM, van Lutterveld R, Kahn RS & Sommer IEC 2475

Prospective study of cannabis use in adolescents at clinical high risk for psychosis: impact on conversion to psychosis and functional outcome

Auther AM, McLaughlin D, Carrión RE, Nagachandran P, Correll CU & Cornblatt BA 2485

Evidence that the wider social environment moderates the association between familial liability and psychosis spectrum outcome

Binbay T, Drukker M, Alptekin K, Elbi H, Aksu Tanik F, Özkınay F, Onay H, Zağlı N & van Os J 2499

Interbirth spacing and offspring mental health outcomes

Riordan DV, Morris C, Hattie J & Stark C 2511

Longitudinal follow-up of cavum septum pellucidum and adhesio interthalamica alterations in first-episode psychosis: a population-based MRI study

Trzesniak C, Schaufelberger MS, Duran FLS, Santos LC, Rosa PGP, McGuire PK, Murray RM, Scazufca M, Menezes PR, Hallak JEC, Crippa JAS & Busatto GF 2523

Is there change in intelligence quotient in chronically ill schizophrenia patients? A longitudinal study in twins discordant for schizophrenia

Hedman AM, van Haren NEM, van Baal GCM, Brans RGH, Hijman R, Kahn RS & Hulshoff Pol HE 2535

Source memory errors in schizophrenia, hallucinations and negative symptoms: a synthesis of research findings

Brébion G, Ohlsen RI, Bressan RA & David AS 2543

The expression of positive and negative schizotypy in daily life: an experience sampling study

Kwapil TR, Brown LH, Silvia PJ, Myin-Germeys I & Barrantes-Vidal N 2555

Negative emotion impairs working memory in pediatric patients with bipolar disorder type I

Schenkel LS, Passarotti AM, Sweeney JA & Pavuluri MN 2567

Association between pubertal development and depressive symptoms in girls from a UK cohort

Joinson C, Heron J, Araya R, Paus T, Croudace T, Rubin C, Marcus M & Lewis G 2579

A comparison of the clinical characteristics of women with recurrent major depression with and without suicidal symptomatology

Bi B, Xiao X, Zhang H, Gao J, Tao M, Niu H, Wang Y, Wang Q, Chen C, Sun N, Li K, Fu J, Gan Z, Sang W, Zhang G, Yang L, Tian T, Li Q, Yang Q, Sun L, Li Ying, Rong H, Guan C, Zhao X, Ye D, Zhang Y, Ma Z, Li H, He K, Chen J, Cai Y, Zhou C, Luo Y, Wang S, Gao S, Liu J, Guo L, Guan J, Kang Z, Di D, Li Yajuan, Shi S, Li Yihan, Chen Y, Flint J, Kendler K & Liu Y 2591

Validation of the Long-term Difficulties Inventory (LDI) and the List of Threatening Experiences (LTE) as measures of stress in epidemiological population-based cohort studies

Rosmalen JGM, Bos EH & de Jonge P 2599

Short-term SSRI treatment normalises amygdala hyperactivity in depressed patients

Godlewska BR, Norbury R, Selvaraj S, Cowen PJ & Harmer CJ 2609

Changes in severity of depressive symptoms and mortality: the Italian Longitudinal Study on Aging

Scafato E, Galluzzo L, Ghirini S, Gandin C, Rossi A, Solfrizzi V, Panza F, Di Carlo A, Maggi S & Farchi G, for the ILSA Working Group 2619

Personality disorders, common mental disorders and receipt of disability benefits: evidence from the British National Survey of Psychiatric Morbidity

Knudsen AK, Skogen JC, Harvey SB, Stewart R, Hotopf M & Moran P 2631

Generalized anxiety and C-reactive protein levels: a prospective, longitudinal analysis

Copeland WE, Shanahan L, Worthman C, Angold A & Costello EJ 2641

Pregnancy outcomes of women with and without a history of anorexia nervosa

Eagles JM, Lee AJ, Amalraj Raja E, Millar HR & Bhattacharya S 2651

A randomized clinical trial of cognitive behavioral therapy and interpersonal psychotherapy for panic disorder with agoraphobia

Vos SPF, Huibers MJH, Diels L & Arntz A 2661

Offending prior to first psychiatric contact: a population-based register study

Stevens H, Agerbo E, Dean K, Nordentoft M, Nielsen PR & Mortensen PB 2673

Correspondence

2685