

patients with TLE due to alien tissue lesions, but neglects other factors which appeared to play an equally prominent role in Taylor's series, viz., sinistrality and femininity (Taylor, 1975). While the discrepancies in prevalence of schizophrenia between alien tissue lesions and mesial temporal sclerotic lesions may be attributed to differences in the likelihood of misconnections following these two lesions, other explanations cannot be ignored. These include differences in the topography of the two types of lesions. Moreover, temporal lobe alien tissue lesions may be accompanied by similar lesions in other brain regions. The latter could account for the psychopathology. A more provocative explanation for the alien tissue/mesial sclerosis discrepancy – that mesial sclerosis arising from perinatal lesions *protects* against schizophrenia – may not be acceptable to proponents of the aetiological role of obstetric trauma in schizophrenia (Murray *et al*, 1988).

V. L. NIMGAONKAR

Western Psychiatric Institute and Clinic
3811 O'Hara Street
Pittsburgh, PA 15213, USA

References

- MURRAY, R. M., LEWIS, S. W. & REVELEY, A. M. (1985) Towards an aetiological classification of schizophrenia. *Lancet*, *i*, 1023–1026.
SMALL, J. G., SMALL, I. F. & HOYDEN, M. P. (1966) Further psychiatric investigations of patients with temporal and non-temporal lobe epilepsy. *American Journal of Psychiatry*, **123**, 303–310.
TAYLOR, D. C. (1975) Factors influencing the occurrence of schizophrenia-like psychosis in patients with temporal lobe epilepsy. *Psychological Medicine*, **5**, 249–254.
VASQUEZ, J. (1952) Epilepsies temporales et manifestations mentales. *Union Medicale du Canada*, **81**, 1062–1068.

Unreliable urine samples

SIR: I was very interested to read the results of Gossop *et al*'s follow-up of opiate addicts after treatment (*Journal*, March 1989, **154**, 348–353). I am, however, a little unhappy about their use of urine specimens to support claims of abstinence, as in my experience such testing is not sufficiently reliable for such inferences to be drawn. It is difficult, without undue strain to the doctor-patient relationship, to be sure of the origins of a proffered sample. I feel that urine drug screening is often a test of uncertain accuracy, performed on a sample of dubious antecedents, provided by a population noted for the deviousness of their behaviour. It has a place in the assessment of drug use, but added little to Dr Gossop *et al*'s study.

STEVE BROWN

Queen's Medical Centre
University Hospital
Nottingham NG7 2UH

Munchausen's syndrome by proxy

SIR: Munchausen's syndrome by proxy has been called the "hinterland of child abuse" (Meadow, 1977) and there have been numerous, and sometimes quite bizarre, case descriptions. The parents are known, by using fabrication and manipulation, to cause their children to receive unnecessary and potentially harmful investigations and treatment. The presenting problems can vary widely and include neurological symptoms, haematuria and bacteriuria, recurrent diarrhoea and bloody stools, vomiting, dehydration, drowsiness, and urticaria. The management of this disorder can present quite a challenge to health professionals (Nicol & Eccles, 1985; Rosen *et al*, 1983). I report a case with an unusual presentation of self-induced vomiting, anorexia, and weight loss in a 3-year-old girl.

Case report: A 3-year-old girl was admitted to the paediatric ward via her general practitioner, after having been ill for 5 days with an upper respiratory tract infection and perhaps a mild chest infection, for which she had been treated with amoxycillin. This would not in itself have merited admission, but the girl's mother complained that the girl was refusing to eat or drink, and was making herself vomit by "sticking her finger down her throat", hence becoming dehydrated. The girl had been admitted to a hospital in Leeds nine months before this admission and had presented with "vomiting and food refusal". Her mother was very concerned about the child's condition, and repeatedly asked the doctors to set up a drip as she felt her child was "very dehydrated".

Child psychiatric opinion was sought, and it soon became clear that the child's mother was suffering from bulimia nervosa. She gave clear descriptions of bingeing and self-induced vomiting since she was a teenager. Her life had been unsettled and her relationships short-lived. Her child's father left before the child was born, and the mother then lived with another man with whom she had another child. He also left, after much conflict at home, and took their two-year-old girl with him. Now the mother is seeing another man and is pregnant again.

She described her own childhood as "terrible", as her own mother died from "not eating" and she and her sister were brought up from an early age by her maternal grandmother, who was very strict and would beat the children "until they stopped crying". The mother left home as soon as she could and tried to pass examinations in nursing but was unsuccessful. She worked as an auxiliary nurse before having her first baby. She has had numerous admissions into various hospitals for short spells with "non-specific" problems, leaving the child in voluntary care or with friends. Her last admission was for suspected ectopic pregnancy after she collapsed on the paediatric ward while visiting her daughter, who had been readmitted with self-induced vomiting. The mother was discharged after a short period of observation.

We encouraged the mother to attend for individual psychotherapy, with a view to helping her cope with