

FC63 *Neurosciences, psychopharmacology and biological psychiatry*

Symposium : CLINICAL PSYCHIATRIC RESEARCH IN FRANCE :
WITH FFP ("Fédération Française de Psychiatrie"), A NEW SITUATION

3. Psydoc-fr, the French psychiatry on the Internet.
Dr. M. Grohens, Mrs M. Thurin, Dr M. Botbol

Displaying the different services of Psydoc.fr

- "Pour la recherche", electronic edition
- National Scholarly Societies and their activities
- Reviews and their contents ; current bibliographic database.
- Status of the ongoing researches
- Communications of the Research Days and FFP symposia
- Biomedical and Psychiatric Internet addresses
- Information, Events and Seminars
- Forum.

FC64 *Neurosciences, psychopharmacology and biological psychiatry*

4. Research, publications, vocabulary, documentation.

Dr C. Veil

Taking the following into consideration : Abundance and variety of the French scientific publications, of the pool of hospital and university libraries, and also the cutting-off of sites and documentation networks, the FFP promulgated a working group associating researchers and documentalists in order to check off the resources, to identify all sorts of obstacles (institutional inflexibility, anomy of vocabularies, etc.) and to improve inter-group communication.

Presenting the ongoing actions and evaluating the present stage...

FC65 *Neurosciences, psychopharmacology and biological psychiatry*

Symposium : CLINICAL PSYCHIATRIC RESEARCH IN FRANCE
WITH FFP ("Fédération Française de Psychiatrie"), A NEW SITUATION

5. Ethics : from research to practice

Dr J. Garrabé

The significant attached to the observance of medical ethics is one of the most striking effects of research on psychiatric practice. In the following text, we intend to formulate a statement of the present french situation in this field, citing a few examples provided by the activities of the French Federation of Psychiatry. At the time when European texts regarding Bio-ethics are under discussion, this should afford an opportunity to compare the french situation to that pertaining in other countries

A symposium "Ethics and psyche" which took place at UNESCO on october the tenth allowed us to take stock of the international situation. Positions taken on ethical questions by the World Psychiatric Association should allow the psychiatrists to protect their identity from any danger that might stem from the findings of recent research.

FC66 *Neurosciences, psychopharmacology and biological psychiatry*

BRAIN α_2A -ADRENOCEPTORS AND ASSOCIATED REGULATORY PROTEINS IN SUICIDE

J.A. García-Sevilla, P.V. Escribá, R. La Harpe, C. Walzer, J. Guimón. Department of Psychiatry and Institute of Forensic Medicine, University of Geneva, Switzerland, and Lab. of Neuropharmacology, University of the Balearic Islands, E-07071 Palma de Mallorca, Spain.

The regulation of α_2A -adrenoceptors (α_2AR) and regulatory GTP-binding ($G\alpha_{i2}$) proteins and G protein-coupled receptor kinase (GRK2 or βARK) was assessed in brains of suicides. Frontal cortices were collected from 26 suicides (14 M and 12 F; age, 48 ± 3 years; PMD, 30 ± 4 h) and 15 controls (7 M and 8 F; age, 48 ± 5 years; PMD, 31 ± 4 h). Levels of α_2AR , $G\alpha_{i2}$ and GRK2 proteins were assessed by immunoblotting using specific antisera. In brain of suicides, the immunoreactive levels of α_2AR were increased compared with matched controls ($133\pm 4\%$, $n = 26$, $t = 8.02$, $p < 0.01$). The levels of $G\alpha_{i2}$ ($143\pm 8\%$, $n = 21$, $t = 5.64$, $p < 0.01$) and GRK2 ($121\pm 4\%$, $n = 24$, $t = 4.82$, $p < 0.01$) proteins also were increased in the same brain samples. Levels of α_2AR correlated weakly with those of $G\alpha_{i2}$ ($r = 0.35$, $n = 21$, $p = 0.12$), but showed a clear correlation with those of GRK2 ($r = 0.55$, $n = 24$, $p = 0.005$). The results indicate that in brains of suicides 1) the abundance of α_2AR is increased (33%), 2) the density of $G\alpha_{i2}$ proteins is also up-regulated (43%) suggesting an increase in signal transduction through α_2AR , and 3) the abundance of GRK2 is also increased (21%) probably because of the higher functioning of the $\alpha_2AR/G\alpha_{i2}$ complex.

Supported by FNRS grant 3100-043258.95, Switzerland (JG) and by DGICYT grant PB94-0002-Mod C, Spain (JAGS)