

The 13th European Nutrition Conference, FENS 2019, was held at the Dublin Convention Centre, 15–18 October 2019

The effect of a fibre extract from the red seaweed, *Palmaria palmata*, on lipid metabolism and inflammation in healthy adults

Zoe Irwin¹, Emeir M. McSorley², Mary M. Slevin², Lisa Rowan², Paul McMillen², Danielle McCullagh², Pamela J. Magee², Christopher I. R. Gill², Paul Cherry², William Crowe², Conall R. Strain³, Catherine Stanton³ and Philip J. Allsopp²

¹Nutrition Innovation Centre for Food and Health, Ulster University, Portstewart, United Kingdom,

²Nutrition Innovation Centre for Food and Health, Ulster University, Coleraine, United Kingdom and

³Teagasc Food Research Centre, Moorepark, Cork, Ireland

Abstract

Evidence from observational studies indicates that seaweed consumption may reduce the risk of non-communicable diseases such as cardiovascular disease, type two diabetes, and obesity. Accumulating evidence from in vitro and animal studies suggest seaweed have antihyperlipidemic, anti-inflammatory and antioxidant properties which may in part be attributed to the high content of soluble dietary fibre in seaweeds. The viscosity of seaweed fibres is suggested to mediate antihyperlipidemic effects via the alteration of lipid/bile acid absorption kinetics to decrease low-density lipoprotein cholesterol (LDL). Thus, there is a need to evaluate the efficacy of seaweed derived dietary fibre in the management of dyslipidemia. Therefore, the aim of this study was to determine the effect of a fibre rich extract from *Palmaria palmata* on the lipid profile as well as markers of inflammation and oxidative stress in healthy adults. A total of 60 healthy participants (30 male and 30 female) aged 20 to 58 years, were assigned to consume the *Palmaria palmata* fibre extract (5g/day), Synergy-1 and the placebo (maltodextrin) for a duration of 4 weeks with a minimum 4 week washout between each treatment in a double blind, randomised crossover study conducted over 5 months. Fasting concentrations of cholesterol, triglycerides and high-density lipoprotein cholesterol (HDL) were analysed and low-density lipoprotein cholesterol (LDL) and LDL: HDL ratio was calculated. C-reactive protein (CRP) and Ferric Reducing Ability of Plasma (FRAP) were analysed as markers of inflammation and oxidative stress, respectively. Supplementation for 4 weeks with *Palmaria palmata* resulted in favourable changes to lipid profiles with a reduced LDL:HDL ratio; however intention-to-treat univariate ANCOVA identified no significant difference between the treatment groups over time on any of the lipid profile markers. A non-significant increase in CRP and triglyceride concentration along with lower FRAP was also observed with *Palmaria palmata* supplementation. Evidence from this study suggests that *Palmaria palmata* may have effects on lipid metabolism and appears to mobilise triglycerides. More research is needed in individuals with dyslipidaemia to fully elucidate these effects.

Conflict of Interest

There is no conflict of interest.