

Review: Sensor techniques in ruminants: more than fitness trackers

C. H. Knight[†] 

University of Copenhagen, Dyrslægevej 100, Frederiksberg C, DK1870 København, Denmark

(Received 26 September 2019; Accepted 21 November 2019)

In this position paper, I shall summarise the current status of sensor technologies in ruminant livestock farming with emphasis on dairy cattle, outline the case for why I believe that sensor technologies could revolutionise global dairy farming in a positive way, describe the significant barriers that exist if that goal is to be achieved and highlight the benefits to animal wellbeing, profitability and sustainability that could result if the technologies are implemented to a significant extent. I shall not provide a comprehensive review of the sensor technology literature since that has been done before, but I intend to provide a sensible amount of background information and data that will allow the reader to obtain a picture not only of today's sensor usage but, more importantly, the possible future direction of dairy animal-oriented sensor technologies, and I shall substantiate my claims and conclusions with relevant literature.

Keywords: biosensors, dairy animals, precision livestock farming, health, welfare

Implications

Since this is a position paper, the implications are restricted to what might transpire if sensor technologies were to come into widespread use in the dairy industry. There is a potential for the health and welfare of dairy animals to improve as a result of better, more consistent and more frequent (even continuous) monitoring. This would impact directly and positively on farmer profitability, which could also benefit from a reduced labour requirement, and the industry as a whole could benefit from improved consumer perception as a consequence of better animal wellbeing.

Introduction: changes, capabilities and challenges in dairying

Some knowledge of the local and global dairy industries is needed if one is to appreciate the potential impact of 'smart technology'. In the latter years of the 20th century, UK dairy farms were small, cows were known individually and incomes were often marginal. Dairy products were, and still are, supermarket 'loss leaders', a key part of cheap food policies that proliferated across developed countries (Caraher and Coveney, 2004). We worked long hours and, besides attaching milking clusters, one of my regular jobs when our cow numbers increased to more than 100 was the late night walk through the herd looking for the increased activity of cows in heat that would be inseminated the next day.

Later, as a lactation scientist, I became actively involved in biological support to the engineering research that developed one of the first automatic milking systems (AMS) (Hillerton *et al.*, 1990), a technology that has come to be regarded as a dairy industry harbinger of what Daniel Beckmans dubbed Precision Livestock Farming (PLF) (Beckmans, 2008 and 2014; John *et al.*, 2016). What was the point of AMS? Almost certainly it was intended to reduce labour costs, although marketing of such systems has mainly focused on farmer lifestyle and cow choice. What was the chief capability that enabled AMS? In order to locate the teats for cluster attachment, the early robotic arms needed to know which cow they were dealing with so as to access a database of udder coordinates, and so radiofrequency identity (RFID: first patented by Hanton and Leach (1974) as an indwelling rumen bolus radiofrequency device) was the key element. Whether directly linked or not, the gradual uptake of AMS has been accompanied by significant increases in the size of dairy farms (Robbins *et al.*, 2016), a change that has been driven in large part by simple economics: as the number of cows increases, the farm's fixed overhead costs are diluted, especially if AMS allows the expansion to occur without extra labour. The dairy industry is a curious mix of entrepreneurship (adoption of robotics) and extreme conservatism; we were able to demonstrate convincingly that encouraging more frequent attendance at AMS could result in greater lactation persistency and hence enable extended lactations (Pettersson *et al.*, 2011), but the equally clear demonstration that longer lactations

[†] E-mail: chrisknight@breathescience.org

can be economically advantageous (De Vries, 2006) has never been accepted by farmers (or perhaps genetics companies!), who still aim to maximise output from AMS by increasing cow numbers, maximising the genetic potential of the individual cow and rebreeding to an annual schedule. These latter two objectives do not fit well together: high-yielding dairy cows suffer from 'infertility', that is to say, a reduced chance of rebreeding at around 60 days *postpartum*, regarded by many as a key benchmark (but see Dobson *et al.*, 2007 for a critical analysis). Technology has come to the fore in addressing this issue and the 'herd walk-through' is no longer needed. Cows that show increased activity indicative of estrous will be automatically flagged by a pedometer, collar, ear tag, rumen bolus or other device incorporating an accelerometer. Most importantly, the cost is not prohibitive, due in no small part to very much cheaper computing power (a modern 'tablet' contains processing capability that would have cost a theoretical \$10B in the 1960s: Michie *et al.*, 2020) and microcircuitry that ensures that sensors are small (rumen boluses are now available for sheep and goats, in addition to cattle: Castro-Costa *et al.*, 2015).

Larger herds, less intensive labour input, better reproductive management; these should certainly be regarded as important changes that have occurred in the last few decades driven by economics and assisted by technological capabilities. So, what of challenges? In the developed world, obesity is now commonly (but not yet scientifically) regarded as a greater health risk factor than smoking. This belief can actually be traced back almost to the millennium (Sturm and Wells, 2002), but the problem shows no signs of abating. Fitness tracking is part of an increasing trend to exercise more and live a better, healthier lifestyle, as recently reviewed by Böhm *et al.* (2019). Lifestyle choices extend to diet, where dairy products are often seen as obesogenic, hugely damaging to the environment and associated with poor animal welfare (see, e.g., Willett *et al.*, 2019). The first two of these arguments are totally refutable (see, e.g., Guo *et al.*, 2018 regarding obesity and for environmental sustainability, official Food and Agriculture Organization (FAO) data (FAO, 2010) showing that dairy production and distribution generate less than 3% of global greenhouse gas emissions). The third challenge is less easy to defend, since animal welfare is a highly subjective issue, but is an area in which sensors could have a very significant and positive impact. These 'negative' challenges are real and considerable, but dairying should also be seen as a positive part of another equally large societal challenge, our need to feed an expanding global population (Reynolds *et al.*, 2015). There are contrasting views in this respect. In the developed world, consumers are encouraged to reduce dairy consumption (Willett *et al.*, 2019), although the rationale for doing so is not particularly clear; in a nutritional context lower intakes appear to be justified on the basis of (perhaps) doing no harm, rather than doing good. In other parts of the world, the reality is rapid expansion of dairy production capacities; megadairies comprising many tens of thousands of cows are operating in Vietnam and under construction in Russia and China, for

instance (Vietnam Investment Review, 2018) whilst a major project co-funded by governments and a major multinational milk processor aims to revolutionise milk production in West Africa (CARE Denmark, 2015). Global consumer demands for dairy products are almost certainly set to expand considerably, FAO predicting a 22% increase globally in the next 10 years with a concentration in India and Pakistan (OECD, 2018). The challenges are clear, and it seems inevitable that technology will be instrumental in enabling sustainable and ethical production standards in very large herds and in areas of the world that have little tradition of organised dairy production.

State of the art in dairy animal sensors

At the time of writing (late 2019), the search term 'sensors and dairy cows' yields 6 400 000 Internet hits, and this headline statement, taken from Dairy Global (2017): 'Sensors used to detect oestrus, lameness, disease and calving are being touted as the next big thing in dairy production. It is not known, however, if these sensor systems actually improve the health and production of dairy herds'. Interestingly, replacing 'cows' by 'animals' in the search term reduced the number of hits (by 60%!) The take-home messages are clear:

- there is a lot of interest in sensors
- most of this relates to use in cows
- it is not yet easy to justify the investment

Farmers are persuaded that investing in sensors that deliver improved estrous detection is a price worth paying, an analysis borne out by economic modelling (Rutten *et al.*, 2014 and 2018). By contrast, the same group concluded that sensors for body condition score are not yet a good investment, but may be in the future as the technology improves (Rutten *et al.*, 2018). This leads to another take-home message:

- state of the art currently mainly comprises stand-alone sensors for single applications
- some of these (estrous detection in particular) are more mature than others

Perhaps it is not surprising, therefore, that estrous detection sensors dominate the market. In 2017, a colleague working for a manufacturer of sensors was able to identify 22 accelerometer-based commercially available sensor technologies for the neck, ear or leg, all but one of which professed to detect estrous (Thorup, 2017). A year earlier, our own review of the scientific literature identified 12 such products (Caja *et al.*, 2016). This should not be taken as an indication of the rate of expansion in the market! Perhaps our colleague did a better job than we did, but more likely this exposes a further take-home message:

- many of the commercially available technologies have not yet been independently tested under rigorous scientific conditions



Figure 1 (colour online) Overview of sensor technologies associated with dairy animals (example is dairy cow but others could apply equally well). The red zone and individual red dots show 'At Cow' sensors, chiefly accelerometer based but also in some cases including temperature, heart rate and pH analysis. The blue zone shows 'Near Cow' sensors such as video and sound analysis, climate analysis, feed analysis, GPS and interaction with the cloud (the latter two classified as 'near' on the basis of enabling real-time analysis). The green zone shows 'From Cow' sensors that monitor products coming from the cow (milk, hair, saliva, sweat, nasal secretion, breath, faeces, etc.). The black circles represent the main technologies that are commercially available and in widespread use. A cow is shown as example; many of the technologies are also applicable to other dairy ruminants. The figure is not exhaustive.

which takes us back to our earlier statement questioning whether sensors (yet) improve health and productivity. It is perhaps relevant that a recent meta-analysis of human activity trackers (sensors) concluded that almost all of a large research base (850/857 studies) were poorly controlled and 'No evidence was found for the effect of wearable activity trackers, on physical-activity-related outcomes' (Böhm *et al.*, 2019). Before considering whether these problems can be overcome such that sensors do have a future, it is important to have a fuller picture of the range of sensors that are available or conceived of.

Sensors fall into three broad categories, as shown in Figure 1. The first is often referred to as 'Wearable' sensors since these are found attached to the cow, but I prefer the term 'At Cow' which then also includes devices swallowed into the reticulorumen or inserted into the reproductive tract (red zone in Figure 1). By analogy, the second category is 'Near Cow' (blue zone in Figure 1) and the third is 'From Cow' (green zone in Figure 1). Near Cow includes all remote sensors that could watch, listen to, track, weigh, record or interrogate the cow or its immediate environment and includes elements that are geographically very remote (Global Positioning Satellite technology, GPS and cloud), but still 'near' in the sense of enabling real-time interaction. From Cow is the specific category of sensors that could collect and analyse data relating to products that have come from the cow, milk analysis being the prime example but biomarker analysis of body tissues/fluids (e.g., hair, saliva, sweat, nasal secretion, breath, faeces) also being included.

I include the potential for robotic collection of biomarker samples in the third category. To the best of my knowledge, this possibility has not yet been considered even at a research level, which leads us to consider which of these sensor technologies are not only commercially available but also in actual widespread use. These are represented by the black circles in Figure 1 and comprise numerous At Cow sensors (very largely accelerometers and/or temperature sensors) that in most cases communicate with cloud-based data management, together with online milk analysis. Automatic milking systems should also be mentioned as a non-sensor technology that has seen widespread adoption (Svennersten-Sjaunja and Pettersson, 2008). A comprehensive list of sensor technologies that were available in 2016 is available in tabular form in Caja *et al.* (2016). Other technologies are commercially available, such as image-based assessment of body condition (Krukowski, 2009) and udder disease conditions (Castro-Costa *et al.*, 2014), force plates for automated BW recording (Dickinson *et al.*, 2013) and limb disease conditions (Ghotoorlar *et al.*, 2012) and sensors for monitoring rumen pH and informing on rumen dysfunction (Sato *et al.*, 2012). I should add that this list is probably not comprehensive, the references refer to the basic technology rather than necessarily the commercial product and in some cases the scientific case for the technology or product is not established. Further consideration and comparison of a 'successful adoption' (estrous detection) and a 'failed adoption' (automated weighing) is given by Maltz (2020). The rumen pH bolus is an interesting case in point. The technology has been available for some years, as have commercial products.

However, the sensor is expensive and suffers from a short life due to a combination of the acidic conditions in the rumen and heavy power requirement. As a result, few (if any) farmers have invested directly in such sensors, which are used primarily for research (see companion paper by Dijkstra *et al.*) or by feed companies or veterinarians offering a service to farmers who have a herd-level nutritional problem. Many more technologies are in research development, such as image-based assessment of individual feeding behaviour (Bloch *et al.*, 2019), rumen function (Song *et al.*, 2019) and heart and respiratory rates (Beiderman *et al.*, 2014). Totally, novel sensor modalities under development include laser reflection-based LiDAR for body composition (Huang *et al.*, 2018) and micro-Doppler radar for heart and respiratory rate (Michie *et al.*, 2020).

Estrous detection

Technologies that require specific mention are RFID and tri-axial accelerometry since these are at the heart of estrous detection. Both have been comprehensively reviewed by Caja *et al.* (2016). Radiofrequency identity technologies include large high-frequency transponders (e.g., neck collars) that also enable collection and transmission of large data packages as well as the cheaper miniaturised low-frequency devices (ear tags, injectable devices or rumen boluses) that are purely for identity. Both types of transponder are 'passive' (i.e., power resides with the receiving element) and the two types are not inter-compatible. Radiofrequency identity technology is optional for most EU dairy cattle farmers, but compulsory for the larger national populations (>600 000 head) of small ruminants as the only way to trace and to build up credible animal inventories.

Tri-axial accelerometers are small, cheap and robust. In addition to estrous detection in cattle, they form the basis of the activity trackers that many of us carry or wear. Clearly, a device that measures changes in motion in three axes is not capable of detecting the main physiological indicators of estrous (pheromones) as a bull would. Furthermore, it is not activity *per se* that identifies the onset of estrous, rather, it is the change in activity of the individual animal from day to day. Trying to identify which of, say, 200 cows in a herd were in estrous on a specific day using only that day's data would give very unreliable results. This is a crucial observation that will underpin our later discussion of sensor technology use for wellbeing evaluation; technology is likely to be very much better at identifying change within a cow across time than variation between cows within a herd or between herds ('benchmarking') at a single point in time. Accelerometers may be located at various points on the animal (rear ankle, neck, ear, tailhead) as well as in the rumen. All are capable of detecting estrous and some can do more than that, especially if combined with a second sensor modality. Examples are a gyroscope that provides positional information to allow easier detection of lying/standing time, and temperature thermistors that enable

calving detection when expelled from the vagina or, as we shall see later, information on rumen function and general health as part of a bolus sensor. For a detailed review of estrous detection using sensors, see Mottram (2016).

Biomarkers

The third category of sensors identified in Figure 1 comprises biomarkers of one sort or another. In current commercial use, this relates exclusively to milk analysis. Milk can yield important information relating not just to downstream consumer properties (i.e., product quality) but also upstream animal status (Duplessis *et al.*, 2019). Systems are available for milk composition (relevant to metabolic status: Leitner *et al.*, 2012; Larsen *et al.*, 2016), somatic cell count and conductivity (relevant to udder disease status; Albrechtsen *et al.*, 2011) and, in one system, biomarkers of reproductive (Bruinjé *et al.*, 2019) and metabolic function. In contrast to this rather restricted commercial usage, the amount of research effort directed to biomarkers of one sort or another is enormous, and beyond the scope of this position paper. Fortunately, the field has been reviewed very recently by de Almeida *et al.* (2019) in relation to health and welfare and by Zachut *et al.* (2020) in relation to fertility and metabolic regulation. Biomarkers show great promise, but a word of caution is required. The physiological principle underpinning the use of milk conductivity for mastitis detection was first shown in the mid-70s (Linzell and Peaker, 1975), and it took 30 years or more for the principle to be put into commercial practice and some studies still question the validity of the approach (Khatun *et al.*, 2019).

Sensors for wellbeing

Several of the estrous detection manufacturers now also offer systems (usually add-on modules) that identify (or claim to identify) cows that are at risk of health or welfare problems of one sort or another: statements like 'your herd's health and well-being, at your fingertips' and 'effective animal health monitoring' are easy to find on commercial Internet sites. Principle amongst these are mastitis, lameness and metabolic disease detection, and it is probably no coincidence that these are the major health problems afflicting dairy cows. How realistic are the claims? From a scientific point of view, there has been little or no independent validation of which I am aware. Is that necessarily a problem? Consider the case of mastitis. Automatic milking systems is a technology that has been around for some time, and since there is no dairyman to detect udder abnormalities during the milking process, AMS requires a technological approach to mastitis detection. Sensor modalities in use include somatic cell counting (SCC), lactate dehydrogenase concentration and milk conductivity. We have just mentioned scientific concerns about some aspects of this (Khatun *et al.*, 2019) but is there any industry-level indication of significant problems? Frössling

et al. (2017) documented AMS uptake in Sweden between 2008 and 2011 and showed that AMS was associated with increased mastitis risk (elevated SCC), but that this effect decreased over time and was less important than other factors such as time of year. In Sweden as a whole, AMS use increased but recorded mastitis incidence decreased over the period, which could, of course, reflect either improved udder health or detection failure. In short, the situation is complicated but for this particular example, there is no real evidence of a growing problem. In relation to animal wellbeing as a whole, is this maintenance of *status quo* satisfactory? Not in my view! Two of the major factors that potentially decrease wellbeing are higher milk yields and larger farms. The first of these has been recognised for many years and is not straightforward (Knight *et al.*, 2004), but it is the second where technology can have real impact, and it should be to improve, rather than to avoid deterioration. The problem is simple. Economics drives the move towards bigger herds and more animals per dairyman (see discussion in Caja *et al.*, 2016), so surveillance becomes more difficult. The concept that technology could redress the balance by providing a precise diagnosis of a specific problem in a specific animal at a specific time is attractive, but almost certainly unrealistic. Mastitis detection systems are set to identify cows that *may* have a problem using algorithms that attempt to balance false positives and false negatives in an optimal way; the final decision that a cow does/does not need treatment is made by a dairyman. The crux is to develop systems that constantly monitor simple (but meaningful) wellbeing-related parameters, looking for change indicative of a suboptimal state ('constantly' will be qualified later and should not be confused with 'frequently'). Lying time is a measurable trait that is almost certainly related to lameness (Maselyne *et al.*, 2017) and is, again, almost certainly how at least one commercial sensor 'detects' lameness. The problem is that lying time is also affected by many other factors, so the same false-positive/negative issues apply, and it will probably be several years before the true efficacy of lameness detection systems can be established. Other simple indicators of wellbeing that can be measured using today's state of the art are feeding, rumination and drinking, and in the next section we will see how this is done and how it could be useful in the future.

Beyond state of the art

The premise of this position paper is that sensor technologies could revolutionise global dairy farming in a positive way, and Figure 2 summarises how that might be achieved. The first priority is to identify an optimistic but achievable target, here defined as Optimized Decision Support. The reasons for this target are twofold:

- the focus is on improving overall husbandry, rather than 'solving' specific disease problems
- it is clear that the technology supports the farmer, and not *vice versa*.




Figure 2 (colour online) Beyond state of the art: schematic of research and development routes to optimised sensor-based decision support for dairy animal husbandry. The red shaded routes represent the likely minimum effort required to achieve the objective. R&D = research and development; IOT: Internet of things.

The second priority is to recognise the need for coordinated, integrated and synergistic research and development (R&D) effort from a spectrum of essential disciplines that include a range of animal-related sciences together with an equally broad range of technology-related sciences. This was a major consideration in a recently completed COST Action in technology for dairy animal health and welfare, FA1308 DairyCare (DairyCare, 2019). Sadly, such integration does not always happen: animal scientists met recently at the Ghent Conference of the European Association of Animal Production to discuss (amongst other things) PLF, at exactly the same time that engineers were meeting in Cork to discuss the same issues in the European Conference on PLF.

There are a number of potential R&D routes from current state of the art towards the target which coalesce into two common elements, a need for data integration (considered below) and then a business model for implementation (considered later). Is there a need for new sensors to extend the state of the art? Is there a need for more of the sensors that we already have? I take the view that neither are essential, but both will happen and both are likely to speed progress. On the issue of new sensors, there is a need to work backwards from a target to identify how technology can achieve that target, rather than forwards from a technical possibility to a now redefined target that may not be really what is needed. The example of rumen bolus sensors is relevant. A major manufacturer of rumen pH boluses is currently testing the market for additional biochemical functionalities (ammonium and nitrate) to be added to their sensor. Whilst these may have relevance to researchers investigating nitrogen cycling, the relevance to farmers is difficult to see and the sensor is likely to suffer the same limited lifetime issues as the existing pH sensor. Those with knowledge of physiological rumen function will know that rumen muscular contractions of three distinct types mix digesta, regurgitate it for rumination and remove methane by eructation. It is

our belief that a pressure sensor could record these different contractions and therefore give farmer-relevant information on animal wellbeing (target: impaired rumen function is a 'first stop' for veterinary diagnosis) as well as carbon footprint (target: evidence-based low carbon footprint has potential economic value), and with funding from the Hannah Dairy Research Foundation and as a collaboration between Spanish animal scientists and Scottish engineers, we hope to develop such a sensor (Castro-Costa *et al.*, 2019).

Adding more and more of the same sensors to the global dairy sector takes us into the realms of 'big data' and 'the Internet of Things, IoT' (reviewed by Michie *et al.*, 2020). Internet of Things is the concept that 'things' (people or animals but also pieces of equipment, vehicles, sensors, etc.) can be connected through local communication wireless networks and the Internet (cloud communication). If large amounts of data are linked together in this way, it potentially becomes something referred to as big data and may accrue increased value. Do current generation dairy animal sensors produce big data? Emphatically no! The size of big data is generally accepted to be in the zetabyte range (zeta is 10^{21} so a zetabyte is a kilobyte raised to the seventh power) approximating to data from 6bn smartphones. Additional criteria apply; the data should have variety, be obtained in real time or approaching it and be fit for purpose (the four Vs: Volume, Variety, Velocity and Veracity). Accelerometer sensors, for example, can potentially collect very large amounts of data, but the amount that they communicate (transmit to a receiver and thereafter the IoT) is constrained by battery power. To get around this problem, the data are processed locally by the sensor device using algorithms that identify summary features relevant to the trait of interest, and it is this data that is then sent onwards. So the sensor may be sensing constantly, but for practical reasons the measurements are packaged into summaries covering discrete windows of time, corresponding to observation frequency. A typical estrous detection sensor will function effectively sending just 1 or 2 kilobytes of data daily! (Michie *et al.*, 2020). In short, big data is a misconception, but on the other hand all of the commercially available sensors are IoT enabled, and in one specific case the manufacturer is heavily targeting this aspect and offers a product that is built around integrating data from very many customers globally into an artificial intelligence (AI) system (Connecterra, 2019).

In current practice, although manufacturers claim to offer 'complete solutions', no one sensor system individually offers everything that could be achieved using a full combination of all systems operating together, so for maximum benefit the farmer's only course of action is to invest in more than one system, shown as 'Multiple Sensors' in Figure 2. Furthermore, almost without exception, the different technologies operate 'stand alone' and will not communicate with each other, limiting the usefulness of the IoT. However, there may be circumstances in which multiple sensors do achieve a specific objective. Some smartwatch devices produced for golfers employ an accelerometer to detect shots, which works well for drives and fairway shots, but not for putts. Although the

device could probably be made more sensitive to overcome this problem, this would introduce another problem, namely false positives. The solution adopted by the manufacturer is to supply an additional accelerometer that attaches directly to the putter, detects the shot and communicates it to the smartwatch. I am not aware of a similar approach being taken by dairy sensor manufacturers, but it could be done.

We have previously mentioned the 'Combined Sensor' in relation to, for instance, temperature and activity in a single rumen bolus. This approach will doubtless be repeated, and I would suggest that a single rumen bolus could potentially include sensors detecting activity, temperature, pressure and heartbeat.

Data mining is an increasingly familiar concept in many scientific fields and can be defined simplistically as 'the process of discovering patterns in large data sets'. The 'cocktail party effect' is a good example of what is happening: in a crowded and noisy room your attention is defocused from the background 'babble' (which your brain filters out) and focused onto a specific word or words, such as your own name. Another example is the processing of collected sensor data that is done by the device before transmission, where the raw activity data (for an accelerometer) are filtered to identify a 'signature' left by the step that the animal has just taken (or the golf shot just made). Steps proved to be relatively easy to find, and data mining algorithms have now advanced to the point where signatures for eating and ruminating can also be identified from several of the commercial accelerometer-based sensors (Michie *et al.*, 2020). Another interesting example is rumen temperature. It quickly became obvious that temperature would fall every time that the cow drank, and this signature was crystal clear even to the naked eye. So, wellbeing indicators including feeding, rumination and drinking are all already available.

Beyond data mining employing a 'static' algorithm, AI or Machine Learning constantly moves the goalposts, in other words, as more and more data arrive the algorithm automatically adapts or 'learns'. Cognitive computing is another term for this phenomenon, the goal being to simulate human thought processes in a computerised model. Artificial intelligence has arrived for dairy husbandry management! The firm marketing it is Dutch and their approach is built around existing sensor technology coming from many, many cows around the globe but with the additional input of the farmer adding a simple assessment of what the data mean to him, based on prompts from the system. As the database grows, the system will progressively get better and better at interpreting the data that sensors are generating. This is a long-term prospect; in the shorter term variation in how the farmers assess the data will likely limit gains made in the final interpretation.

Data integration is the next essential stage of the process, and is an area in which a lot could be achieved quickly, were it not for commercial sensitivities that restrict data sharing. All of the commercial systems do already integrate and compare data, in particular that coming from the same individual animal on consecutive days. The principle is simple; the

this value by appropriate interaction with the other players identified in the figure, and the potential returns will be reflected in the contract price agreed with the farmer. The further advantage of this sort of approach is that it overcomes the problem that technological approaches might only be affordable, or apply to, the largest farms. In this scenario, a cluster of say, 20 farms of 250 head all instrumented by the same service provider would in effect be equivalent to a single farm of 5000. The nature of the service provider is flexible. It could simply be a commercial company, but equally it could be a farmer cooperative or a national agency such as a breed improvement service.

Conclusions

The estimated global value of animal biosensor technologies is expected to exceed \$20 billion by 2020 (Neethirajan *et al.*, 2017) and a large share of that will be in dairy, so it is true to say that sensor systems for dairy husbandry have come a long way in a relatively short time. Since the only market to have been rigorously exploited to date is estrous detection, it is equally true to say that they are still in their infancy in comparison to what could become possible in the future. There is no reason why sensors systems could not find widespread use for health and welfare monitoring, and to do that not just in dairy cows but in other dairy species as well. Technologies capable of measuring relevant parameters such as eating and drinking already exist, and the ability to extract relevant information from complex datasets is growing all the time. The need for technologies that will assist husbandry is growing as dairy farms expand in size, and the potential for wellbeing data to add value to the primary product is there to be exploited. Success will require coordinated effort from a spectrum of biologists, engineers and business experts in areas of R&D and marketing that are identifiable. There is a potential for sensor systems to benefit not just the farmer and their animals, but also other players in the dairy foods chain as well as consumers. Indeed, it is probably not an overstatement to say that sensor-based husbandry support could be a key element of the dairy sector's vital future contribution to global food security.

Acknowledgements

This article is based upon work from COST Action FA1308 DairyCare, supported by COST (European Cooperation in Science and Technology, www.cost.eu). COST is a funding agency for research and innovation networks. COST Actions help connect research initiatives across Europe and enable scientists to grow their ideas by sharing them with their peers. This boosts their research, career and innovation.  C. H. Knight 0000-0001-9361-2249

Declaration of interest

The author does not receive any commercial or other financial or in-kind support related to the subject matter of this article and therefore declares no conflict of interest.

Ethics statement

Not applicable. No original data or experimentation is presented in this article.

Software and data repository resources

Not applicable. No original data or experimentation is presented in this article.

References

- Albrechtsen M, Duse AT, Bennedsgaard TW and Klaas IC 2011. Use of in-line measurements of somatic cell count to evaluate treatment efficacy of subclinical bovine *Staphylococcus aureus* mastitis. In *Udder health and communication* (ed. H Hogeveen and TJGM Lam), pp. 309–315. Wageningen Academic Pub, Wageningen, The Netherlands.
- Beiderman Y, Halachmi I and Zalevsky Z 2014. A novel approach for remote monitoring of heart beat rate, respiratory rate and chewing activity in cows. In *Proceedings of the First DairyCare Conference*, 22 August 2014. Abstract 2.6 DairyCare COST Action, Copenhagen. ISBN 978-0-9930176-0-5.
- Berckmans D 2008. Precision livestock farming (PLF). *Computers and Electronics in Agriculture* 62, 1–1.
- Berckmans D 2014. Precision livestock farming technologies for welfare management in intensive livestock systems. *Reviews of Science and Technology* 33, 189–196.
- Bloch V, Levit H and Halachmi I 2019. Assessing the potential of photogrammetry to monitor feed intake of dairy cows. *Journal of Dairy Research* 86, 34–39.
- Böhm B, Karwiese SD, Böhm H and Oberhoffer R 2019. Effects of mobile health including wearable activity trackers to increase physical activity outcomes among healthy children and adolescents: systematic review. *JMIR Mhealth and Uhealth* 7, e8298.
- Bruinjé TC, Colazo MG, Ribeiro ES, Gobikrushanth M and Ambrose DJ 2019. Using in-line milk progesterone data to characterize parameters of luteal activity and their association with fertility in Holstein cows. *Journal of Dairy Science* 102, 780–798.
- Caja G, Castro-Costa A and Knight CH 2016. Engineering to support wellbeing of dairy animals. *Journal of Dairy Research* 83, 136–147.
- Caraher M and Coveney J 2004. Public health nutrition and food policy. *Public Health Nutrition* 7, 591–598.
- CARE Denmark 2015. The milky way to development. Retrieved on 22 September 2019 from <https://care.dk/focuscountries/niger-eng/milky-way-development/>
- Castro-Costa A, Caja G, Michie C, Andonovic I and Knight CH 2019. Building and testing a rumen function sensor. *Proceedings of the Second Next Generation Dairying Workshop*, Edinburgh. Retrieved on 22 September 2011 from <https://www.journalofdairyresearch.org/next-generation-dairying.html>
- Castro-Costa A, Caja G, Salama AA, Rovai M, Flores C and Aguiló J 2014. Thermographic variation of the udder of dairy ewes in early lactation and following an *Escherichia coli* endotoxin intramammary challenge in late lactation. *Journal of Dairy Science* 97, 1377–1387.
- Castro-Costa A, Salama AA, Moll X, Aguiló J and Caja G 2015. Using wireless rumen sensors for evaluating the effects of diet and ambient temperature in non-lactating dairy goats. *Journal of Dairy Science* 98, 4646–4658.
- Connecterra 2019. Innovation through intelligence: an AI for the agricultural industry. Retrieved on 10 October 2019 from <https://www.connecterra.io/>
- Dairy Global 2017. Sensor value and viability for dairy cows. Retrieved on 22 September 2019 from <https://www.dairyglobal.net/Smart-farming/Articles/2017/8/Sensor-value-and-viability-for-dairy-cows-166802E/>
- DairyCare 2019. COST Action FA1308 DairyCare: technology for improved dairy animal husbandry. Retrieved on 15 September 2019 from <https://www.journalofdairyresearch.org/dairy-care-on-jdr.html>
- de Almeida AM, Zachut M, Hernandez-Castellano LE, Šperanda M, Gabai G and Mobasher A 2019. Biomarkers of fitness and welfare in dairy animals: healthy living. *Journal of Dairy Research* 86, 379–387.

- De Vries A 2006 Economic value of pregnancy in dairy cattle. *Journal of Dairy Science* 89, 3876–3885.
- Dickinson RA, Morton JM, Beggs DS, Anderson GA, Pyman MF, Mansell PD and Blackwood CB 2013. An automated walk-over weighing system as a tool for measuring liveweight change in lactating dairy cows. *Journal of Dairy Science* 96, 4477–4486.
- Dobson H, Smith RF, Royal MD, Knight CH and Sheldon IM 2007. The high producing dairy cow and its reproductive performance. *Reproduction in Domestic Animals* 42 (suppl. 2), 17–23.
- Duplessis M, Santschi DE, Plante S, Bergeron C, Lefebvre DM, Durocher J and Cue RI 2019. Milk β -hydroxybutyrate concentration measured by Fourier-transform infrared and flow-injection analyses from samples taken at different times relative to milking. *Journal of Dairy Research* 86, 208–210.
- FAO 2010. Greenhouse gas emissions from the dairy sector: a life cycle assessment. Retrieved on 15 September 2019 from <http://www.fao.org/3/k7930e/k7930e00.pdf>
- Frössling J, Ohlson A and Hallén-Sandgren C 2017. Incidence and duration of increased somatic cell count in Swedish dairy cows and associations with milking system type. *Journal of Dairy Science* 100, 7368–7378.
- Ghotoorlar SM, Ghamsari SM, Nowrouzian I, Ghotoorlar SM and Ghidary SS 2012. Lameness scoring system for dairy cows using force plates and artificial intelligence. *Veterinary Record* 170, 126.
- Guo J, Dougkas A, Elwood PC and Givens DI 2018. Dairy foods and body mass index over 10-year: evidence from the caerphilly prospective cohort study. *Nutrients* 10, E1515.
- Hanton JP and Leach HA 1974. Electronic livestock identification system. US Patent 4.262.632.
- Hillerton JE, Knight CH, Turvey A, Wheatley SD and Wilde CJ 1990. Milk yield and mammary function in dairy cows milked four times daily. *Journal of Dairy Research* 57, 285–294.
- Huang L, Li S, Zhu A, Fan X, Zhang C and Wang H 2018. Non-contact body measurement for Qinchuan cattle with LiDAR sensor. *Sensors (Basel)* 18. pii: E3014. doi: [10.3390/s18093014](https://doi.org/10.3390/s18093014).
- John AJ, Clark CE, Freeman MJ, Kerrisk KL, Garcia SC and Halachmi I 2016. Review: milking robot utilization, a successful precision livestock farming evolution. *Animal* 10, 1484–1492.
- Khatun M, Bruckmaier RM, Thomson PC, House J and Garcia SC 2019. Suitability of somatic cell count, electrical conductivity, and lactate dehydrogenase activity in foremilk before versus after alveolar milk ejection for mastitis detection. *Journal of Dairy Science* 102, 9200–9212.
- Knight CH 2020. Blueprint for action in the development of technology for improved dairy animal husbandry. *Journal of Dairy Research* 87, In press.
- Knight CH, Alamer MA, Sorensen A, Nevison IM, Flint DJ and Vernon RG 2004. Metabolic safety-margins do not differ between cows of high and low genetic merit for milk production. *Journal of Dairy Research* 71, 141–153.
- Krukowski M 2009. Automatic determination of body condition score of dairy cows from 3D images. MSc thesis, Royal Institute of Technology, School of Computer Science and Communication, Stockholm, Sweden.
- Larsen T, Alstrup L and Weisbjerg MR 2016. Minor milk constituents are affected by protein concentration and forage digestibility in the feed ration. *Journal of Dairy Research* 83, 12–19.
- Leitner G, Merin U, Lemberskiy-Kuzin L, Bezman D and Katz G 2012. Real-time visual/near-infrared analysis of milk-clotting parameters for industrial applications. *Animal* 6, 1170–1177.
- Linzell JL and Peaker M 1975. Efficacy of the measurement of the electrical conductivity of milk for the detection of subclinical mastitis in cows: detection of infected cows at a single visit. *British Veterinary Journal* 131, 447–461.
- Maltz E 2020. Individual dairy cow management: achievements, obstacles and prospects. *Journal of Dairy Research* 87, In press.
- Maselyne J, Pastell M, Thomsen PT, Thorup VM, Hänninen L, Vangeyte J, Van Nuffel A and Munksgaard L 2017. Daily lying time, motion index and step frequency in dairy cows change throughout lactation. *Research in Veterinary Science* 110, 1–3.
- Michie C, Andonovic I, Davison C, Tachtatzis C, Hamilton A, Gilroy M and Johnsson N 2020. The Internet-of-things enhancing animal welfare and farm operational efficiency. *Journal of Dairy Research* 87, In press.
- Mottram T 2016. Animal board invited review: precision livestock farming for dairy cows with a focus on oestrus detection. *Animal* 10, 1575–1584.
- Neethirajan S, Huang S-T, Tuteja SK and Kelton D 2017. Recent advancement in biosensors technology for animal and livestock health management. *Biosensors and Bioelectronics* 98, 398–407.
- OECD 2018. OECD-FAO agricultural outlook 2018–2027. Retrieved on 22 September 2019 from https://www.oecd-ilibrary.org/agriculture-and-food/oecd-fao-agricultural-outlook-2018-2027/dairy-and-dairy-products_agr_outlook-2018-10-en
- Pettersson G, Svennersten-Sjaunja K and Knight, CH 2011. Relationships between milking frequency, lactation persistency and milk yield in Swedish Red heifers and cows milked in a voluntary attendance automatic milking system. *Journal of Dairy Research* 78, 1–6.
- Reynolds LP, Wulster-Radcliffe MC, Aaron DK and Davis TA 2015. Importance of animals in agricultural sustainability and food security. *Journal of Nutrition* 145, 1377–1379.
- Robbins JA, von Keyserlingk MA, Fraser D and Weary DM 2016. Invited review: farm size and animal welfare. *Journal of Animal Science* 94, 5439–5455.
- Rutten CJ, Steeneveld W, Inchaisri C and Hogeveen H 2014. An ex ante analysis on the use of activity meters for automated estrus detection: to invest or not to invest? *Journal of Dairy Science* 97, 6869–6887.
- Rutten CJ, Steeneveld W, Oude Lansink AGJM and Hogeveen H 2018. Delaying investments in sensor technology: the rationality of dairy farmers' investment decisions illustrated within the framework of real options theory. *Journal of Dairy Science* 101, 7650–7660.
- Sato S, Mizuguchi H, Ito K, Ikuta K, Kimura A and Okada K. 2012. Technical note: development and testing of a radio transmission pH measurement system for continuous monitoring of ruminal pH in cows. *Preventative Veterinary Medicine* 103, 274–279.
- Song X, van der Tol PPJ, Groot Koerkamp PWG and Bokkers EAM 2019. Hot topic: automated assessment of reticulo-ruminal motility in dairy cows using 3-dimensional vision. *Journal of Dairy Science* 102, 9076–9081.
- Sturm R and Wells KB 2002. The health risks of obesity. Worse than smoking, drinking or poverty. RAND Research Briefs. Retrieved on 15 September 2019 from https://www.rand.org/pubs/research_briefs/RB4549.html
- Svennersten-Sjaunja KM and Pettersson G 2008. Pros and cons of automatic milking in Europe. *Journal of Animal Science* 86 (13 suppl.), 37–46.
- Thorup VM 2017. Precision livestock farming for dairy: how sensors can help. Retrieved on 10 October 2019 from <https://www.futurefarming.com/theatre/>
- Vietnam Investment Review 2018. TH Group kicks off giant dairy complex in Russia. Retrieved on 22 September 2019 from <https://www.vir.com.vn/th-group-kicks-off-giant-dairy-complex-in-russia-62508.html>
- Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, Garnett T, Tilman D, DeClerck F, Wood A, Jonell M, Clark M, Gordon LJ, Fanzo J, Hawkes C, Zurayk R, Rivera JA, De Vries W, Sibanda LM, Afshin A, Chaudhary A, Herrero M, Agustina R, Branca F, Lartey A, Fan S, Crona B, Fox E, Bignet V, Troell M, Lindahl T, Singh S, Cornell SE, Reddy KS, Narain S, Nishtar S and Murray CJL 2019. Food in the Anthropocene: the EAT-Lancet Commission on healthy diets from sustainable food systems. *Lancet* 393, 447–492.
- Zachut M, Speranda M, de Almeida A, Gabai G, Mobasher A and Hernandez-Castellano LE 2020. Biomarkers of fitness and welfare in dairy cattle: healthy productivity. *Journal of Dairy Research* 87, In press.