

NORMAL CURVATURE OF MINIMAL SUBMANIFOLDS IN A SPHERE

by SHARIEF DESHMUKH

(Received 23 June, 1995)

1. Introduction. Simons [5] has proved a pinching theorem for compact minimal submanifolds in a unit sphere, which led to an intrinsic rigidity result. Sakaki [4] improved this result of Simons for arbitrary codimension and has proved that if the scalar curvature S of the minimal submanifold M^n of S^{n+p} satisfies

$$\frac{n(n-1)(2n^2+n-8)}{2(n^2+n-3)} \leq S$$

then either M^n is totally geodesic or $S = 2/3$ in which case $n = 2$ and M^2 is the Veronese surface in a totally geodesic 4-sphere. This result of Sakaki was further improved by Shen [6] but only for dimension $n = 3$, where it is shown that if $S > 4$, then M^3 is totally geodesic (cf. Theorem 3, p. 791).

Let M^n be a compact minimal submanifold of the unit sphere S^{n+p} with normal bundle ν . We denote by R^\perp the curvature tensor field corresponding to the normal connection ∇^\perp in the normal bundle ν of M^n , and define $K^\perp: M \rightarrow R$ by

$$K^\perp = \sum_{i,j,\alpha,\beta} [R^\perp(e_i, e_j, N_\alpha, N_\beta)]^2,$$

where $\{e_1, \dots, e_n\}$ is a local orthonormal frame on M^n and $\{N_1, \dots, N_p\}$ is a local field of orthonormal normals. We call the function K^\perp the normal curvature of the minimal submanifold M^n . In this paper we prove the following result.

THEOREM. *Let M^n be a compact minimal submanifold of S^{n+p} . If the normal curvature K^\perp , the scalar curvature S and the square of the length of the second fundamental form σ of M^n satisfy*

$$K^\perp \leq \sigma, \quad S > (n-1)^2,$$

then M^n is totally geodesic.

This theorem can be considered as a partial generalization of the result of Shen [6, Theorem 3]. However, it will be an interesting question whether the condition $K^\perp \leq \sigma$ is redundant and Shen's result can be extended beyond dimension 3.

2. Preliminaries. Let M be a minimal submanifold of the unit sphere S^{n+p} , with normal bundle ν . Then the second fundamental form h of M^n satisfies

$$(\nabla h)(X, Y, Z) = (\nabla h)(Y, Z, X) = (\nabla h)(Z, X, Y), \quad X, Y, Z \in \mathcal{X}(M), \quad (2.1)$$

where $\mathcal{X}(M)$ is the Lie algebra of smooth vector fields on M and $(\nabla h)(X, Y, Z)$ is defined by

$$(\nabla h)(X, Y, Z) = \nabla_X^\perp h(Y, Z) - h(\nabla_X Y, Z) - h(Y, \nabla_X Z),$$

where ∇^\perp is the connection defined in ν and ∇ is the induced Riemannian connection

with respect to the induced Riemannian metric g on M^n . The second covariant derivative $(\nabla^2 h)(X, Y, Z, W)$ of the second fundamental form is given by

$$(\nabla^2 h)(X, Y, Z, W) = \nabla_X^\perp(\nabla h)(Y, Z, W) - (\nabla h)(\nabla_X Y, Z, W) \\ - (\nabla h)(Y, \nabla_X Z, W) - (\nabla h)(Y, Z, \nabla_X W), \quad X, Y, Z, W \in \mathcal{X}(M).$$

We have the following form of the Ricci identity

$$(\nabla^2 h)(X, Y, Z, W) - (\nabla^2 h)(Y, X, Z, W) = R^\perp(X, Y)h(Z, W) - h(R(X, Y)Z, W) \\ - h(Z, R(X, Y)W), \quad X, Y, Z, W \in \mathcal{X}(M), \quad (2.2)$$

where R^\perp and R are the curvature tensors of the connections ∇^\perp and ∇ respectively. Since M^n is a minimal submanifold for a local orthonormal frame $\{e_1, \dots, e_n\}$ of M^n we have

$$\sum_{i=1}^n (\nabla h)(X, e_i, e_i) = 0, \\ \sum_{i=1}^n (\nabla^2 h)(X, Y, e_i, e_i) = 0. \quad (2.3)$$

Using the Ricci tensor Ric , we define the symmetric operator R^* by

$$\text{Ric}(X, Y) = g(R^*(X), Y), \quad X, Y \in \mathcal{X}(M).$$

Then the Gauss equation gives

$$A_{h(Y,Z)}X = R(X, Y)Z + A_{h(X,Z)}Y - g(Y, Z)X + g(X, Z)Y, \quad (2.4)$$

$$R^*(X) = (n-1)X - \sum_{i=1}^n A_{h(e_i, X)}e_i, \quad X, Y, Z \in \mathcal{X}(M), \quad (2.5)$$

where $A_N, N \in \nu$, is the Weingarten map with respect to the normal N , satisfying $g(A_N X, Y) = g(h(X, Y), N)$. We define

$$\sigma = \sum_{ij} \|h(e_i, e_j)\|^2, \\ \|A_h\|^2 = \sum_{ij,k} \|A_{h(e_i, e_j)}e_k\|^2, \\ \|\nabla h\|^2 = \sum_{ij,k} \|(\nabla h)(e_i, e_j, e_k)\|^2. \quad (2.6)$$

Now we prove the following lemma.

LEMMA. Let M^n be a minimal submanifold of S^{n+p} , then for a local orthonormal frame $\{e_1, \dots, e_n\}$, we have

$$\sum_{ij,k} R(e_k, e_i; e_j, A_{h(e_i, e_j)}e_k) = -\sigma + \|A_h\|^2 + \frac{1}{2}K^\perp - \sum_{ij, \alpha, \beta} g(A_\alpha e_i, A_\beta e_j)^2,$$

where $A_\alpha \equiv A_{N_\alpha}$ and $\{N_1, \dots, N_p\}$ is a local field of orthonormal normals.

Proof. Using the Ricci equation

$$R^\perp(X, Y; N_1, N_2) = g([A_{N_1}, A_{N_2}](X), Y), \quad X, Y \in \mathcal{X}(M), N_1, N_2 \in \nu,$$

we get

$$\begin{aligned}
 K^\perp &= \sum_{i,j,\alpha,\beta} [R^\perp(e_i, e_j; N_\alpha, N_\beta)]^2 = \sum_{i,j,\alpha,\beta} [g(A_\alpha A_\beta e_i, e_j) - g(A_\beta A_\alpha e_i, e_j)]^2 \\
 &= 2 \sum_{i,j,\alpha,\beta} g(A_\alpha e_i, A_\beta e_j)^2 - 2 \sum_{i,j,\alpha,\beta} g(A_\alpha A_\beta e_i, e_j)g(A_\beta A_\alpha e_i, e_j), \tag{2.7}
 \end{aligned}$$

since $\sum_{i,j,\alpha,\beta} g(A_\alpha e_j, A_\beta e_i)^2 = \sum_{i,j,\alpha,\beta} g(A_\beta e_j, A_\alpha e_i)^2$ which follows from the symmetry of A_α and A_β . Next using the Gauss equation, we have

$$\begin{aligned}
 R(e_k, e_i; e_j, A_{h(e_i, e_j)} e_k) &= \delta_{ij}g(h(e_k, e_k), h(e_i, e_j)) - \delta_{kj}g(h(e_i, e_j), h(e_i, e_k)) \\
 &\quad + g(h(e_i, e_j), h(e_k, A_{h(e_i, e_j)} e_k)) - g(h(e_k, e_j), h(e_i, A_{h(e_i, e_j)} e_k)) \tag{2.8}
 \end{aligned}$$

since $A_{h(e_i, e_j)} e_k = \sum_\alpha g(A_\alpha e_i, e_j)A_\alpha e_k$, we obtain

$$\sum_{i,j,k} g(h(e_i, e_j), h(e_k, A_{h(e_i, e_j)} e_k)) = \sum_{i,j,k} g(A_{h(e_i, e_j)} e_k, A_{h(e_i, e_j)} e_k) = \|A_h\|^2 \tag{2.9}$$

and

$$\sum_{i,j,k} g(h(e_k, e_j), h(e_i, A_{h(e_i, e_j)} e_k)) = \sum_{i,j,\alpha,\beta} g(A_\alpha A_\beta e_i, e_j)g(A_\beta A_\alpha e_i, e_j). \tag{2.10}$$

Then using (2.7), (2.9) and (2.10) in (2.8) and using minimality of M^n we find

$$\sum_{i,j,k} R(e_k, e_i; e_j, A_{h(e_i, e_j)} e_k) = -\sigma + \|A_h\|^2 - \sum_{i,j,\alpha,\beta} g(A_\alpha e_i, A_\beta e_j)^2 + \frac{1}{2}K^\perp$$

which proves the lemma.

3. Proof of the theorem. Let M^n be a compact minimal submanifold of S^{n+p} satisfying the hypothesis of the theorem. Define $F: M \rightarrow R$ by $F = \frac{1}{2}\sigma$. Then it is straightforward to compute the Laplacian ΔF of the function F as

$$\Delta F = \sum_{i,j,k} g((\nabla^2 h)(e_k, e_k, e_i, e_j), h(e_i, e_j)) + \sum_{i,j,k} \|(\nabla h)(e_i, e_j, e_k)\|^2.$$

Using the Ricci identity (2.2) and equations (2.1) in above equation we arrive at

$$\begin{aligned}
 \Delta F &= \sum_{i,j,k} [R^\perp(e_k, e_i; h(e_k, e_j), h(e_i, e_j)) - R(e_k, e_i; e_k, A_{h(e_j, e_i)} e_j) \\
 &\quad - R(e_k, e_i; e_j, A_{h(e_i, e_j)} e_k)] + \|\nabla h\|^2.
 \end{aligned}$$

We employ (2.4) in the Ricci equation, to compute

$$\begin{aligned}
 R^\perp(e_k, e_i; h(e_k, e_j), h(e_i, e_j)) &= g(A_{h(e_i, e_j)} e_k, A_{h(e_i, e_j)} e_k) \\
 &\quad + R(e_i, e_k) e_j - \delta_{kj} e_i + \delta_{ij} e_k - g(A_{h(e_k, e_j)} e_k, A_{h(e_i, e_j)} e_i)
 \end{aligned}$$

or

$$\begin{aligned}
 \sum_{i,j,k} R^\perp(e_k, e_i; h(e_k, e_j), h(e_i, e_j)) &= \|A_h\|^2 - \sigma + \sum_{i,j,k} R(e_i, e_k; e_j, A_{h(e_i, e_j)} e_k) \\
 &\quad - g(A_{h(e_k, e_j)} e_k, A_{h(e_i, e_j)} e_i). \tag{3.2}
 \end{aligned}$$

Since (2.5) gives $R^*(e_j) = (n - 1)e_j - \sum_k A_{h(e_k, e_j)} e_k$, we have

$$\begin{aligned}
 \sum_{i,j,k} g(A_{h(e_k, e_j)} e_k, A_{h(e_i, e_j)} e_i) &= \sum_{i,j} g((n - 1)e_j - R^*(e_j), A_{h(e_i, e_j)} e_i) \\
 &= (n - 1)\sigma - \sum_{i,j} \text{Ric}(e_j, A_{h(e_i, e_j)} e_i)
 \end{aligned}$$

$$\begin{aligned}
&= (n-1)\sigma - \sum_{i,j,k} R(e_k, e_j, A_{h(e_i, e_j)} e_i, e_k) \\
&= (n-1)\sigma + \sum_{i,j,k} R(e_k, e_i, e_k, A_{h(e_i, e_j)} e_j).
\end{aligned} \tag{3.3}$$

Thus using (3.3) in (3.2), we have

$$\begin{aligned}
\sum_{i,j,k} R^\perp(e_k, e_i; h(e_k, e_j), h(e_i, e_j)) &= -n\sigma + \|A_h\|^2 + \sum_{i,j,k} [R(e_i, e_k; e_j, A_{h(e_i, e_j)} e_k) \\
&\quad - R(e_k, e_i, e_k, A_{h(e_i, e_j)} e_j)].
\end{aligned} \tag{3.4}$$

Using (3.4) in (3.1), we obtain

$$\begin{aligned}
\Delta F &= -n\sigma + \|A_h\|^2 - 2 \sum_{i,j,k} [R(e_k, e_i; e_j, A_{h(e_i, e_j)} e_k) \\
&\quad - R(e_k, e_i, e_k; A_{h(e_i, e_j)} e_j)] + \|\nabla h\|^2.
\end{aligned} \tag{3.5}$$

Also, we have

$$\begin{aligned}
\sum_{i,j,k} R(e_k, e_i; e_k, A_{h(e_i, e_j)} e_j) &= -\sum_{i,j} \text{Ric}(e_i, A_{h(e_i, e_j)} e_j) \\
&= -\sum_{i,j} g(R^* e_i, A_{h(e_i, e_j)} e_j) \\
&= -\sum_{i,j,\alpha} g(R^* e_i, A_\alpha e_j) g(A_\alpha e_i, e_j) \\
&= -\sum_{i,j,\alpha} g(R^* A_\alpha e_j, e_i) g(A_\alpha e_j, e_i) \\
&= -\sum_{j,\alpha} g(R^* A_\alpha e_j, A_\alpha e_j) \\
&= -\sum_{j,\alpha} \text{Ric}(A_\alpha e_j, A_\alpha e_j) \\
&= -\sum_{j,\alpha} (n-1)g(A_\alpha e_j, A_\alpha e_j) + \sum_{i,j,\alpha} \|h(e_i, A_\alpha e_j)\|^2 \\
&= -(n-1)\sigma + \sum_{i,j,\alpha,\beta} g(A_\beta e_i, A_\alpha e_j)^2.
\end{aligned} \tag{3.6}$$

Using (3.6) and the lemma in Section 2 in (3.5), we obtain

$$\Delta F = (n-1)\sigma - \|A_h\|^2 + (\sigma - K^\perp) + \|\nabla h\|^2. \tag{3.7}$$

Now using the facts that

$$\begin{aligned}
\|A_h\|^2 &= \sum_{i,j,k} \|A_{h(e_i, e_j)} e_k\|^2 = \sum_{i,j,k,\alpha} g(A_\alpha e_i, e_j)^2 \|A_\alpha e_k\|^2 \\
&= \sum_{i,j,\alpha} g(A_\alpha e_i, e_j)^2 \|A_\alpha\|^2 = \sum_\alpha \|A_\alpha\|^2 \|A_\alpha\|^2 = \sum_\alpha \|A_\alpha\|^4
\end{aligned}$$

and $\sigma = \sum_\alpha \|A_\alpha\|^2$, in (3.7) and integrating it over M^n we obtain

$$\int_M \left\{ \sum_\alpha [(n-1) - \|A_\alpha\|^2] \|A_\alpha\|^2 + (\sigma - K^\perp) + \|\nabla h\|^2 \right\} dv = 0. \tag{3.8}$$

From the hypothesis of the theorem $S > (n - 1)^2$, it follows that

$$n(n - 1) - \sum_{\alpha} \|A_{\alpha}\|^2 > (n - 1)^2,$$

that is, $\sum_{\alpha} \|A_{\alpha}\|^2 < (n - 1)$, consequently $\|A_{\alpha}\|^2 < (n - 1)$, and that $K^{\perp} \leq \sigma$. Thus in order for (3.8) to hold we must have $\|A_{\alpha}\| = 0$, that is M^n is totally geodesic.

REFERENCES

1. J. L. Barbosa and M. do Carmo, Stability of minimal surfaces and eigenvalues of the Laplacian, *Math. Z.* **173** (1980), 13–28.
2. S.-Y. Cheng, Eigenvalue comparison theorems and its geometric applications, *Math. Z.* **143** (1975), 289–297.
3. S. S. Chern, M. do Carmo and S. Kobayashi, Minimal submanifolds of a sphere with second fundamental form of constant length, *Functional analysis and related fields* (Springer, 1970), 59–75.
4. M. Sakaki, Remarks on the rigidity and stability of minimal submanifolds, *Proc. Amer. Math. Soc.* **106** (1989), 793–795.
5. J. Simons, Minimal varieties in riemannian manifolds, *Ann. of Math. (2)* **88** (1968), 62–105.
6. Y. B. Shen, Curvature pinching for the three-dimensional minimal submanifolds in a sphere, *Proc. Amer. Math. Soc.* **115** (1992), 791–795.

DEPARTMENT OF MATHEMATICS
 COLLEGE OF SCIENCE
 KING SAUD UNIVERSITY
 P.O. BOX 2455
 RIYADH-11451
 SAUDI ARABIA