

First Report of *Sphenothallus* Hall (Cnidaria, Medusozoa) from the Mesozoic Erathem (Upper Triassic, Slovenia)

Heyo Van Iten,^{1,2*} Rok Gašparič,^{3,4} Tomaž Hitij,^{5,6} Tea Kolar-Jurkovšek,⁷ and Bogdan Jurkovšek⁸

¹Department of Geology, Hanover College, Hanover, Indiana 47243, USA <vaniten@hanover.edu>

²Department of Invertebrate Paleontology, Cincinnati Museum Center, 1301 Western Avenue, Cincinnati, Ohio 45203, USA

³Oertijmuseum, Bosscheweg 80, 5293 WB Boxtel, the Netherlands <rok.gasparic@gmail.com>

⁴Institute for Palaeobiology and Evolution, Novi trg 59, 1241 Kamnik, Slovenia

⁵University of Ljubljana, Faculty of Medicine, Vrazov trg 2, 1000 Ljubljana, Slovenia <tomazhitij@gmail.com>

⁶Institute for Palaeobiology and Evolution, Novi trg 59, 1241 Kamnik, Slovenia

⁷Geological Survey of Slovenia, Dimičeva ulica 14, 1000 Ljubljana, Slovenia <tea.kolar-jurkovsek@geo-zs.si>

⁸Kamnica 27, 1262 Dol pri Ljubljani, Slovenia <geolog.bj@gmail.com>

Introduction

Sphenothallus Hall, 1847, one of the most widely distributed and longest ranging genera in the fossil record, has been documented from all systems of the Paleozoic Erathem except the Permian (Table 1), although it has been stated (e.g., Choi, 1990; Bolton, 1994; Fatka et al., 2012) that the genus also occurs in that system. At present the first appearance of this epibenthic, polypoid medusozoan cnidarian lies in Cambrian Stage 3, while the previously known youngest occurrences are in the Pennsylvanian System. *Sphenothallus* has been found in numerous formations on all continents except Australia and Antarctica. It occurs in a variety of marine facies ranging from shallow near-shore to deep offshore and has even been found in strata of coastal lacustrine origin, probably as an allochthonous element (Lerner and Lucas, 2011). Many of the rock units known to contain *Sphenothallus* also contain conulariids (Table 1), an extinct group of marine scyphozoans that may have been closely related to *Sphenothallus* (Van Iten et al., 1992, 1996). Van Iten et al. (1992) interpreted *Sphenothallus* as a medusozoan cnidarian of uncertain class-level affinities, but later Dzik et al. (2017) documented internal peridermal structures that may be homologous to similar features in the periderm of coronate scyphozoans (see for example illustrations in Van Iten, 1992, and Van Iten et al., 1996).

The present article describes multiple, very well-preserved specimens of *Sphenothallus carniolica* (Kolar-Jurkovšek and Jurkovšek, 1997) from limestone strata of early Late Triassic (late Carnian) age in the Julian Alps of northwest Slovenia. Some of these specimens were originally described under the name *Valvasoria carniolica* Kolar-Jurkovšek and Jurkovšek, 1997, which was interpreted as a tubiculous worm of possible nematode or sipunculid affinities (see also Hitij et al., 2019). Thus, this is the first report of *Sphenothallus* from the Mesozoic Erathem. Furthermore, *Sphenothallus* is now established as a long-ranging medusozoan cnidarian, which, together with conulariids, survived the End Permian Mass Extinction Event (MacLeod, 2013).

Geological setting

The studied *Sphenothallus* specimens were collected from thin, gray, laminated lime mudstones in the Kozja dnina Member of the Martuljek Limestone (early Late Triassic, late Carnian) in the northeast Julian Alps (Vrata Valley) of northwest Slovenia (Fig. 1). The late Carnian (Tuvalian) age of the section that yielded the specimens is based on conodont assemblages (*Quadralella polygnathiformis* Zone; Kolar-Jurkovšek, 1991). The section at Kozja dnina is ~80 m thick and represents a deep water paleoenvironment (Bitner et al., 2010). The strata were deposited in an interplatform basin, where anoxic conditions and rapid sedimentation enabled exceptional preservation of both invertebrate and vertebrate fossils (Celarc and Kolar-Jurkovšek, 2008). In addition to *Sphenothallus*, the Kozja dnina limestones also contain bivalves, brachiopods, echinoids, crinoids, asteroids, ammonites, belemnites, scleractinian corals, shrimp, lobsters, thylacocephalans, and fishes (Hitij et al., 2019; Gašparič et al., 2020).

Material and methods

The present study is based on direct examination of 31 specimens of *Sphenothallus carniolica* (Kolar-Jurkovšek and Jurkovšek, 1997) from the Kozja dnina Member of the Martuljek Limestone. All specimens occur on exposed bedding planes, their preserved periderm appearing light to dark brown on the light gray weathered limestone, in some cases with a blueish hue resulting from vivianitization. Specimens were examined and photographed using reflected light and scanning electron microscopy (secondary electron mode). Material photographed under reflected light was whitened with ammonium chloride sublimate. Photographs were taken with a millimetric scale bar using a Nikon Z 6II digital camera equipped with a NIKKOR Z MC 105mm f/2.8 VR S lens. Photographs were edited in Photoshop CS6, and figures containing light photographs were assembled in CorelDRAW X8. Scanning electron microscopy was conducted using a JEOL JSM-6490LV. Finally, the elemental composition of the periderm of one of the specimens (T-1287) was determined using an Oxford INCA Energy 350 EDS under the following operating conditions: chamber vacuum

*Corresponding Author.

Table 1. Selected documented occurrences of *Sphenothallus* Hall, 1847, in the Phanerozoic rock record. Asterisk next to a formation or member name indicates that the unit also yields conulariids.

System	Stage(s) or Series	Formation(s)	Locality(ies)	Reference(s)
Cambrian	Stage 3	Guojiaba, Xiannudong	China (Shaanxi Province)	Li et al., 2004
	Stage 3	Niutitang	China (Hunan and Guizhou provinces)	Peng et al., 2005; Muscente and Xiao, 2015
	Stage 3	Balkasandstein	Denmark	Troppenz, 2020
	Stages 3–4	Shipai, Shuijingtuo	China (Hubei Province)	Muscente and Xiao, 2015; Chang et al., 2018
	Stage 4	Kaili	China (Anhui Province)	Zhao et al., 1999; Zhu et al., 2000; Peng et al., 2005
	Stage 4	Harkless	USA (Nevada)	Skovsted and Holmer, 2006
	Wuliuan	Stephen	Canada (British Columbia)	Van Iten et al., 2002
	Wuliuan	Jince	Czech Republic	Fatka et al., 2012
	Wuliuan	Buchava Skryje Mbr.	Czech Republic	Fatka and Kraft, 2013
	Ordovician	Tremadocian	Dumugol	Korea
Tremadocian–Floian		Fezouata Shale*	Morocco	Van Iten et al., 2016a
Floian		Tonggao *	China (Hubei Province)	Van Iten et al., 2013
Floian		Fenxiang*	China (Hubei Province)	Baliński and Sun, 2015; Dzik et al., 2017
Floian–Dapingian		Klabava*	Czech Republic	Bruthansová and Van Iten, 2020
Sandbian		Zahořany*	Czech Republic	Bruthansová and Van Iten, 2020
Sandbian		Viivikonna*	Estonia	Vinn and Kirsimäe, 2015
Sandbian–Katian		Dillsboro*	USA (Indiana)	Bodenbender et al., 1989
Katian		Lindsay Collingwood Mbr.*	Canada (Ontario)	Bolton, 1994
Katian		Utica Shale*	Canada (Quebec)	Bolton, 1994
Silurian	Katian	Maquoketa Elgin* and Brainard* mbrs.	USA (Iowa and Minnesota)	Van Iten et al., 1996, 2005
	Rhuddanian	Lung-Machi	China (Guizhou Province)	Yi et al., 2003
	Telychian	Brandon Bridge	USA (Wisconsin)	Miller et al., 2022
Devonian	Llandovery	Cape Schuchert	Greenland	Peel, 2021
	Emsian	Hunsrück Slate*	Germany	Fauchald et al., 1986; Südkamp, 2017
Mississippian	(multiple)	(multiple)	USA (multiple states)	Mason and Yochelson, 1985
	Pragian–Emsian	Ponta Grossa Shale Jaguariaíva Mbr.*	Brazil (Paraná State)	Van Iten et al., 2019
	Famennian	Chagrín Shale*	USA (Ohio)	Feldmann et al., 1986; Neal and Hannibal, 2000
Pennsylvanian	Serpukhovian	Heath Bear Gulch Mbr.*	USA (Montana)	Van Iten et al., 1992
	Serpukhovian	Gurovo Dashkovko Mbr.*	Russia (Moscow Basin)	Van Iten et al., 2022
Triassic	(multiple)	(multiple)	USA (multiple states)	Mason and Yochelson, 1985
	Kazimovian	Atrasado Tinajas Mbr.	USA (New Mexico)	Lerner and Lucas, 2005, 2011
	(multiple)	(multiple)	USA (multiple states)	Mason and Yochelson, 1985
Triassic	(multiple)	(multiple)	Belgium, France, Germany, Netherlands	Schmidt and Teichmüller, 1956, 1958
	Carnian	Martuljek Limestone Kozja dnina Mbr.	Slovenia	This paper

20 Pa, accelerating voltage 20 kV, spot size 48 µm, working distance 10 mm, and analysis time 60 seconds.

Repositories and institutional abbreviations.—BJ = Paleontological collection of Jurkovšek, Dol pri Ljubljani, Slovenia. T = Paleontological collection of Tomaž Hitij and Jure Žalohar, Godič, Slovenia. All collections are registered at the Slovenian Museum of Natural History in Ljubljana.

Systematic paleontology

- Phylum Cnidaria Verrill, 1865
- Subphylum Medusozoa Peterson, 1979
- Class, Order, Family uncertain
- Genus *Sphenothallus* Hall, 1847

Type species.—*Sphenothallus angustifolius* Hall, 1847, originally described from the Upper Ordovician of eastern New York State, USA.

Sphenothallus carniolica (Kolar-Jurkovšek and Jurkovšek, 1997)
 Figures 3–5

1997 *Valvasoria carniolica* Kolar-Jurkovšek and Jurkovšek, p. 1, pl. 1, figs. 1–4.

2019 *Valvasoria carniolica*; Hitij et al., p. 21, figs. 16, 17.

Holotype.—BJ1286 (originally designated as the holotype of *Valvasoria carniolica* Kolar-Jurkovšek and Jurkovšek, 1997; currently deposited in the collection of Bogdan Jurkovšek).

Paratypes.—BJ1287, BJ1419 (originally designated as paratypes of *Valvasoria carniolica*; currently deposited in the collection of Bogdan Jurkovšek).

Diagnosis.—Peridermal tube very gently tapered (angle of expansion mostly <2°), non-branching, lacking regular transverse annulations.

Occurrence.—Thin, gray, laminated lime mudstones in the Kozja dnina Member of the Martuljek Limestone (early Late Triassic, late Carnian, Tuvalian; *Paragondolella polygnathiformis* conodont Zone) in the Vrata Valley in the northeast Julian Alps, northwest Slovenia. The geographic coordinates of the fossil locality are Lat. 46°24'08.1"N, Long. 13°50'46.2"E.




Figure 1. Simplified geographical map of northwest Slovenia. The location of Kozja dnina in the Vrata Valley is marked by a star (figure adapted from Bitner et al., 2010).

Description.—Partial to nearly complete, compressed specimens lying parallel to bedding and ranging up to 129.5 mm in length and 5.5 mm in width. Specimens consist of one or more portions of the slender, originally sub-elliptical (transversely) main tube or of nearly the entire periderm, including the sub-conical attachment disc, although without the basal membrane. Attachment disc measures up to ~3.1 mm in diameter. Apertural margin not preserved; terminal schott (apical wall) absent. Main tube very gently tapered (mostly $<2^\circ$) in the plane of the well-developed pair of oppositely situated, longitudinal thickenings, variably curved parallel to bedding, in some cases with the degree of curvature increasing toward the apical end. Relatively thin periderm between the longitudinal thickenings missing or, where present, exhibiting coarse, irregular, transverse to oblique wrinkles. Regular transverse annulations absent. Longitudinal

thickenings terminate or thin near the apertural end of the main tube. Skeletal material phosphatic, very finely lamellar, light to dark brown or dark bluish gray, with exfoliated lamellae in some specimens exhibiting possible “plywood [micro]structure” (Vinn and Mironenko, 2021). Some specimens exhibit minute, shallow, sub-circular to sub-elliptical pores or pits ranging from ~2–4 μm in diameter.

Additional material.—BJ2505 (currently repositied in the collection of Bogdan Jurkovšek); T-1256, T-1257, T-1270–T-1287 (currently repositied in the collection of Tomaž Hitij and Jure Žalohar).

Remarks.—The principal diagnostic feature of *Sphenothallus*, namely the pair of longitudinal thickenings situated at the end points of the major diameter of the subelliptical main tube, is




Figure 2. Lithostratigraphy of the Tuvalian section with platy limestones yielding *Sphenothallus carniolica* (Kolar-Jurkovšek and Jurkovšek, 1997) (arrow) at Kozja dnina (figure adapted from Celarc and Kolar-Jurkovšek, 2008). Norian stage subdivisions are abbreviated as: Lac. = Lacinian; Alaun. = Alaunian; Sevat. = Sevatian.

well developed in the Slovenian material (Figs. 3.1–3.3, 3.5, 4.1–4.3, 5.1–5.4, 5.8). As in *Sphenothallus* from Paleozoic formations, compression of the Slovenian specimens perpendicular to bedding has caused the longitudinal thickenings to form berm-like elevations rising above the thinner, deformed skeletal wall between them, which in some cases (Fig. 5.8) is now absent. Also well displayed in the Slovenian material is the characteristic, parallel lamellar microstructure (Figs. 4.8, 5.5, 5.6), which is commonly exfoliated. Lamellae in one specimen appear to exhibit “plywood [micro]structure” (Fig. 5.5, 5.6), discovered by Vinn and Mironenko (2021, fig. 2C, D) in material from the Upper Mississippian of central Russia. A somewhat similar microstructure is present in the medusozoan *Torella* Holm, 1893, from the upper Cambrian of Estonia (Vinn, 2006,

2022). The presence of calcium phosphate (apatite), which constitutes the bulk of most Paleozoic specimens, is corroborated both by the brown to blue-gray color of the specimens and by the results of EDS analysis (Fig. 6), which yielded strong spectral peaks for Ca and P.

Turning to other characters, the very low rate of taper of the main tube of the Slovenian specimens is similar to that of *S. angustifolius* from the Upper Ordovician of Ontario and Quebec, Canada (see for example Bolton, 1994, pls. 1.1–1.3) and *S. sica* Clarke, 1913, from the Lower Devonian Ponta Grossa Shale of Brazil (see for example Van Iten et al., 2019, fig. 4). The size of the attachment disc (Figs. 3.1, 3.2, 3.4, 3.5, 4.1–4.4, 4.6, 4.7, 5.1, 5.2), both absolute and in proportion to the size of the main tube, matches closely that of the attachment disc of *Sphenothallus* sp. from the Bear Gulch Member of the Upper Mississippian Heath Formation of Montana, western USA (see Van Iten et al., 1992, fig. 1). Interestingly, specimens preserving both the attachment disc and the long slender tube above it may be attached to hard biological substrates (see for example Van Iten et al., 2019, fig. 6), or, like the Slovenian specimens and the Bear Gulch material referred to in the preceding sentence, they may be unattached, suggesting that their original substrate may have been plant or other organic matter susceptible to rapid decay. Finally, smooth curvature of the main tube, with the degree of curvature commonly decreasing in the direction of the apertural end (Figs. 3.2, 4.1, 4.2, 4.6, 5.3, 5.4, 5.6), is a feature exhibited by many Paleozoic specimens (see for example Van Iten et al., 1996, pl. 2.1), as are thinning of the longitudinal thickenings in the vicinity of the aperture (Figs. 3.1, 3.3, 5.4) and the absence of well-defined annulation or other ornament (Fig. 4.5).

A noteworthy additional feature, previously documented in *Sphenothallus* cf. *S. angustifolius* from the Upper Mississippian of central Russia (Vinn and Mironenko, 2021, fig. 2f), is the presence in the main tube of some of the Slovenian specimens of microscopic, sub-circular to sub-elliptical pores or pits (Fig. 5.7), originally detected in specimen BJ1287 by Kolar-Jurkovšek and Jurkovšek (1997, figs. 2, 3). These shallow perforations are similar in size, shape, areal density, and pattern of distribution to the microscopic circular pores, or micropores, of conulariids (see Van Iten et al., 2005, 2006a, b, 2022, and references cited therein). Conulariid micropores have been compared with microscopic borings and bioclastrations, and they have been interpreted as primary anatomical features. If the latter hypothesis is true, then the presence of these features in conulariids and *Sphenothallus* may provide additional support for the hypothesis (e.g., Van Iten et al., 1996) that the skeletons of these two taxa are mutually homologous.

Concluding remarks

Sphenothallus Hall, 1847, an extinct genus of medusozoan cnidarians, is now known from the lower Upper Triassic of north-west Slovenia (Julian Alps). Moreover, the well-preserved specimens of *S. carniolica* from Slovenia are the only documented representatives of the genus from the Mesozoic Erathem. The present discovery extends the known age range of *Sphenothallus* by ca. 80 million years, from the Kazimovian to the late Carnian, and thus across the critical Permian-Triassic boundary.




Figure 3. *Sphenothallus carniolica* (Kolar-Jurkovšek and Jurkovšek, 1997) from the Late Triassic (late Carnian) Kozja dnina Member of the Martuljek Limestone of northwest Slovenia (Vrata Valley). (1–4) BJ1286, the holotype and most complete specimen: (1) side view of the entire specimen, which preserves both the apical holdfast (left end of the specimen) and the apertural region of the main tube (right end of the specimen); (2) detail of the area in box (A) in (1), showing the apical region with berm-like marginal thickenings; (3) detail of the area in box (B) in (1), showing the heavily crumpled, relatively thin apertural region; (4) detail of the area in box (C) in (2), highlighting the sub-conical apical holdfast; (5) BJ1419, paratype, a nearly complete specimen with partially preserved apical holdfast. Scale bars = 30 mm (1) and 10 mm (2–5).

Additionally, since *Sphenothallus* has not yet been documented from any part of the Permian System, it may be a Lazarus taxon (Jablonski, 1986). Nevertheless, we predict that the existence of *Sphenothallus* of Permian age, previously only asserted, will be

confirmed and that additional discoveries of Triassic and maybe even younger *Sphenothallus* will be made, based in part on the ability of *Sphenothallus* to survive the most severe crisis in the history of life during the Phanerozoic Eon. Finally, given the




Figure 4. *Sphenothallus carniolica* (Kolar-Jurkovšek and Jurkovšek, 1997) from the Late Triassic (late Carnian) Kozja dnina Member of the Martuljek Limestone of northwest Slovenia (Vrata Valley). (1) T-1277, mostly complete specimen preserving the apical holdfast; (2) T-1270, two nearly contiguous specimens, both preserving the upper portion of the apical holdfast (basal membrane missing); (3) detail of the area in box (A) in (1), highlighting the apical region; (4) detail of the area in box (C) in (2), showing the apical regions of the two nearly contiguous specimens; (5, 7) scanning electron photomicrographs of T-1277: (5) detail of the area in box (B) in (1), showing the non-annulated external surface of the main tube; (7) detail of the area in box (D) in (3), showing the apicalmost portion of the periderm (7); (6, 8) scanning electron photomicrographs of T-1270: (6) detail of the area in box (E) in (4), showing the apicalmost portions of the specimens; (8) detail of the area in box (F) in (6), showing the wavy laminated microstructure of one of the holdfasts in transverse cross section (8). Scale bars = 10 mm (1, 2), 1 mm (3, 4), 500 μ m (5, 6), 100 μ m (7), and 50 μ m (8).




Figure 5. *Sphenothallus carniolica* (Kolar-Jurkovšek and Jurkovšek, 1997) from the Late Triassic (late Carnian) Kozja dnina Member of the Martuljek Limestone of northwest Slovenia (Vrata Valley). (1) T-1273, two associated specimens, the shorter one of which preserves the apical holdfast; (2) detail of the box (A) in (1), highlighting the apical holdfast and the marginal thickenings, which are slightly elevated above the thinner periderm between them; (3) T-1257, portion of the main tube of a nearly complete specimen exhibiting pervasive irregular wrinkling of the thin periderm between the longitudinal thickenings; (4) T-1256, nearly complete specimen exhibiting pronounced curvature of the main tube in the apical region (left) and pervasive wrinkling of the thin periderm between the robust longitudinal thickenings (right); (5) detail (scanning electron micrograph) of the box (B) in (3), showing the absence of regular annulation of the main tube as well as areas (central part of the field of view) of possible “plywood [micro]structure” (Vinn and Mironenko, 2021); (6) detail (scanning electron micrograph) of the box (C) in (5), showing one of the areas of possible “plywood [micro]structure” (Vinn and Mironenko, 2021); (7) scanning electron micrograph of BJ1419, showing multiple, sub-circular micropores (arrows) on the surface of the periderm; (8) T-1271, assemblage of at least four specimens consisting entirely or predominantly of one or both longitudinal thickenings. Scale bars = 10 mm (1, 3, 4, 6, 8), 1 mm (2), 500 μ m (5), and 20 μ m (7).




Figure 6. EDS spectrum of specimen T-1287, corroborating the presence of calcium phosphate in the periderm.

presence of conulariids, which are commonly associated with *Sphenothallus*, in strata of terminal Ediacaran age (Van Iten et al., 2016b; Leme et al., 2022), we would not be surprised if the first appearance of *Sphenothallus* were extended downward into that system.

Acknowledgments

The authors are grateful to M. Miller (Geological Survey of Slovenia) for SEM photography and EDS analysis of *Sphenothallus* specimens and to the administration of the Triglav National Park for permission to perform paleontological research within its boundary. We also thank reviewers J.S. Peel (Department of Earth Sciences, Uppsala University, Sweden) and O. Vinn (Department of Geology, University of Tartu, Estonia) and associate editor R.J. Elias (Department of Earth Sciences, University of Manitoba, Winnipeg, Canada) for their constructive comments on and corrections of the original manuscript.

Declaration of competing interests

The authors declare none.

References

- Baliński, A., and Sun, Y., 2015, Fensiang biota: a new Early Ordovician shallow-water fauna with soft-part preservation from China: *Science Bulletin*, v. 60, p. 812–818.
- Bitner, M.A., Jurkovšek, B., and Kolar-Jurkovšek, T., 2010, New record of the inarticulate brachiopod genus *Discinisca* from the Upper Triassic (Carnian) of the Julian Alps, NW Slovenia: *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, v. 257, p. 367–372.
- Bodenbender, B.E., Wilson, M.A., and Palmer, T.J., 1989, Paleocology of *Sphenothallus* on an Upper Ordovician hardground: *Lethaia*, v. 22, p. 217–225.
- Bolton, T.E., 1994, *Sphenothallus angustifolius* Hall, 1847 from the lower Upper Ordovician of Ontario and Quebec: *Geological Survey of Canada Bulletin*, v. 479, p. 1–11.
- Bruthansová, J., and Van Iten, H., 2020, Invertebrate epibionts on Ordovician conulariids from the Prague Basin (Czech Republic, Bohemia): *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 558, 109963. <https://doi.org/10.1016/j.palaeo.2020.109963>.

- Celarc, B., and Kolar-Jurkovšek, T., 2008, The Carnian–Norian basin–platform system of the Martuljek Mountain Group (Julian Alps, Slovenia): Progradation of the Dachstein carbonate platform: *Geologica Carpathica*, v. 59, p. 211–224.
- Chang, S., Clausen, S., Zhang, L., Feng, Q.-L., Steiner, M., Bottjer, D.J., Zhang, Y., and Shi, M., 2018, New probable cnidarian fossils from the lower Cambrian of the Three Gorges area, South China, and their ecological implications: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 505, p. 150–166.
- Choi, D.K., 1990, *Sphenothallus* (“Vermes”) from the Tremadocian Dumugol Formation, Korea: *Journal of Paleontology*, v. 64, p. 403–408.
- Clarke, J.M., 1913, Fósseis Devonianos do Paraná: Rio de Janeiro, Monographias do Serviço Geológico e Mineralógico do Brasil, Volume I, 353 p.
- Dzik, J., Baliński, A., and Sun, Y., 2017, The origin of tetradial symmetry in cnidarians: *Lethaia*, v. 50, p. 306–321.
- Fatka, O., and Kraft, P., 2013, *Sphenothallus* Hall, 1847 from Cambrian of Skryje-Týřovice Basin (Barrandian area, Czech Republic): *Annales Societatis Geologorum Poloniae*, v. 83, p. 309–315.
- Fatka, O., Kraft, P., and Szabad, M., 2012, A first report of *Sphenothallus* Hall, 1847 in the Cambrian of Variscan Europe: *Comptes Rendus Palevol*, v. 11, p. 539–547.
- Fauchald, K.W., Stürmer, W., and Yochelson, E.L., 1986, *Sphenothallus* “Vermes” in the Early Devonian Hunsrück Slate, West Germany: *Paläontologische Zeitschrift*, v. 60, p. 57–64.
- Feldmann, R.M., Hannibal, J.T., and Babcock, L.E., 1986, Fossil worms from the Devonian of North America (*Sphenothallus*) and Burma (“Vermes”) previously identified as phyllocarid arthropods: *Journal of Paleontology*, v. 60, p. 341–346.
- Gašparič, R., Audo, D., Hitij, T., Jurkovšek, B., and Kolar-Jurkovšek, T., 2020, One of the oldest polychelidan lobsters from Upper Triassic (Carnian) beds at Kozja dnina locality in the Julian Alps, Slovenia: *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen*, v. 296, p. 107–117.
- Hall, J., 1847, *Palaeontology of New York, Volume 1, Containing Descriptions of the Organic Remains of the Lower Division of the New York System: Albany, Charles Van Benthuysen*, 338 p.
- Hitij, T., Gašparič, R., Založar, J., Jurkovšek, B., and Kolar-Jurkovšek, T., 2019, Paleontološko bogastvo Kozje dnine: *Acta Triglavensia*, v. 7, p. 5–42. [in Slovenian with English summary]
- Holm, G., 1893, Sveriges Kambrisk–Siluriska Hyolithidae och Conulariidae: *Sveriges Geologiska Undersökning, Afhandlingar och Uppsatser*, ser. C, no. 112, p. 1–172. [in Swedish with English summary]
- Jablonski, D., 1986, Background and mass extinctions: the alternation of macro-evolutionary regimes: *Science*, v. 231, p. 129–133.
- Kolar-Jurkovšek, T., 1991, Mikrofavna srednjega in zgornjega Triasa Slovenije in njen biostratigrafski pomen (Microfauna of Middle and Upper Triassic in Slovenia and its biostratigraphic significance): *Geologija*, v. 33, p. 21–170. [in Slovenian with English summary]
- Kolar-Jurkovšek, T., and Jurkovšek, B., 1997, *Valvasoria carniolica* n. gen. n. sp., a Triassic worm from Slovenia: *Geologia Croatica*, v. 50, p. 1–5.

- Leme, J.M., Van Iten, H., and Simões, M.G., 2022. A new conulariid (Cnidaria, Scyphozoa) from the terminal Ediacaran of Brazil: *Frontiers in Earth Science*, v. 10. <https://doi.org/10.3389/feart.2022.777746>.
- Lerner, A.J., and Lucas, S.G., 2005. First New Mexico record of *Sphenothallus* from the Upper Pennsylvanian (Missourian) Atrasado Formation of Socorro County, New Mexico: *New Mexico Geology*, v. 27, p. 52.
- Lerner, A.J., and Lucas, S.G., 2011. Allochthonous *Sphenothallus* (Cnidaria) from a lacustrine Lagerstätte, Carboniferous of New Mexico, USA, in Sullivan, R.M., Lucas, S.G., and Spielmann, J.A., eds., *Fossil Record 3: New Mexico Museum of Natural History and Science, Bulletin 53*, p. 86–89.
- Li, G.-X., Zhu, M.-Y., Van Iten, H., and Li, C.-W., 2004. Occurrence of the earliest known *Sphenothallus* Hall in the Lower Cambrian of southern Shaanxi Province, China: *Geobios*, v. 37, p. 229–237.
- MacLeod, N., 2013. *The Great Extinctions. What Causes Them and How They Shape Life*: Buffalo, New York, Firefly Books, 208 p.
- Mason, C., and Yochelson, E.L., 1985. Some tubular fossils (*Sphenothallus*: “Vermes”) from the middle and late Paleozoic of the United States: *Journal of Paleontology*, v. 59, p. 85–95.
- Miller, A.A., Huntley, J.W., Anderson, E.P., and Jacquet, S.M., 2022. Biotic interactions between conulariids and epibionts from the Silurian Waukesha Biota: *Palaios*, v. 37, p. 691–699.
- Muscente, A.D., and Xiao, S., 2015. New occurrences of *Sphenothallus* in the lower Cambrian of South China: implications for its affinities and taphonomic demineralization of small shelly fossils: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 437, p. 141–164.
- Neal, M.L., and Hannibal, J.T., 2000. Paleocological and taxonomic implications of *Sphenothallus* and *Sphenothallus*-like specimens from Ohio and areas adjacent to Ohio: *Journal of Paleontology*, v. 74, p. 369–380.
- Peel, J.S., 2021. Holdfasts of *Sphenothallus* (Cnidaria) from the early Silurian of western North Greenland (Laurentia): *GFF*, v. 143, p. 384–389.
- Peng, J., Babcock, L.E., Zhao, Y.-L., Wang, P.-L., and Yang, R.-J., 2005. Cambrian *Sphenothallus* from Guizhou Province, China: early sessile predators: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 220, p. 119–127.
- Peterson, K.W., 1979. Development of coloniality in Hydrozoa, in Larwood, G., and Rosen, B., eds., *Biology and Systematics of Colonial Organisms*: New York, Systematics Association, Special Volume 11, p. 105–139.
- Schmidt, W., and Teichmüller, M., 1956. Die Enträtselung eines bislang unbekannt Fossils im deutschen Oberkarbon, *Sphenothallus stubblefeldi* n. sp., und die Art seines Auftretens: *Geologische Jahrbuch*, v. 71, p. 243–298.
- Schmidt, W., and Teichmüller, M., 1958. Neue Funde von *Sphenothallus* auf dem westeuropäischen Festland, insbesondere in Belgien, und ergänzende Beobachtungen zur Gattung *Sphenothallus*: *Association pour l'Étude de la Paléontologie et de la Stratigraphie Houillères, Publication 33*, 34 p.
- Skovsted, C.B., and Holmer, L.E., 2006. The lower Cambrian brachiopod *Kyrshabaktella* and associated shelly fossils from the Harkless Formation, southern Nevada: *GFF*, v. 128, p. 327–337.
- Südkamp, W., 2017. *Life in the Devonian. Identification Book. Hunsrück Slate Fossils*: München, Germany, Verlag Dr. Friedrich Pfeil, 176 p.
- Tropenz, U.-M., 2020. Ein mutmaßlicher unterkambrischer *Sphenothallus* sp. (Cnidaria) von Nienhagen bei Warnemünde, Mecklenburg, Deutschland: *Mitteilungen der Naturforschenden Gesellschaft Mecklenburg*, v. 20, p. 27–31.
- Van Iten, H., 1992. Anatomy and phylogenetic significance of the corners and midlines of the conulariid test: *Palaeontology*, v. 35, p. 335–358.
- Van Iten, H., Cox, R.S., and Mapes, R.H., 1992. New data on the morphology of *Sphenothallus* Hall: implications for its affinities: *Lethaia*, v. 25, p. 135–144.
- Van Iten, H., Fitzke, J.A., and Cox, R.S., 1996. Problematical fossil cnidarians from the Upper Ordovician of the north-central USA: *Palaeontology*, v. 39, p. 1037–1064.
- Van Iten, H., Zhu, M.-Y., and Collins, D., 2002. First report of *Sphenothallus* Hall, 1847 in the Middle Cambrian. *Journal of Paleontology*, v. 76, p. 902–905.
- Van Iten, H., Vhylasová, Z., Zhu, M.-Y., and Zhuo, E.-J., 2005. Widespread occurrence of microscopic pores in conulariids: *Journal of Paleontology*, v. 79, p. 400–407.
- Van Iten, H., Lichtenwalter, M., Leme, J.M., and Simões, M.G., 2006a. Possible taphonomic bias in the preservation of phosphatic macroinvertebrates in the uppermost Maquoketa Formation (Upper Ordovician) of northeastern Iowa (north-central USA): *Journal of Taphonomy*, v. 4, p. 207–220.
- Van Iten, H., Leme, J.M., and Simões, M.G., 2006b. Additional observations on the gross morphology and microstructure of *Baccaconularia* Hughes, Gundersen et Weedon, 2000, a Cambrian (Furongian) conulariid from the north-central USA: *Palaeoworld*, v. 15, p. 294–306.
- Van Iten, H., Muir, L.A., Botting, J.P., Zhang, Y.-D., and Li, J.-P., 2013. Conulariids and *Sphenothallus* (Cnidaria, Medusozoa) from the Tonggao Formation (Lower Ordovician, China): *Bulletin of Geosciences*, v. 88, p. 713–722.
- Van Iten, H., Muir, L., Simões, M.G., Leme, J.M., Marques, A.C., and Yoder, N., 2016a. Palaeobiogeography, palaeoecology and evolution of Lower Ordovician conulariids and *Sphenothallus* (Medusozoa, Cnidaria), with emphasis on the Fezouata Shale of southeastern Morocco: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 460, p. 170–178.
- Van Iten, H., Leme, J.M., Pacheco, M.L.A.F., Simões, M.G., Fairchild, T.R., Rodrigues, F., Galante, D., Boggiani, P.C., and Marques, A.C., 2016b. Origin and early diversification of Phylum Cnidaria: key macrofossils from the Ediacaran System of North and South America, in Goffredo, S., and Dubinsky, Z., *The Cnidaria, Past, Present and Future*: Springer, Cham International Publishing, p. 31–40.
- Van Iten, H., Leme, J.M., Simões, M.G., and Cournoyer, M.E., 2019. Clonal colony in the Early Devonian cnidarian *Sphenothallus* from Brazil: *Acta Palaeontologica Polonica*, v. 64, p. 409–416.
- Van Iten, H., Mironenko, A.A., and Vinn, O., 2022. A new conulariid from the Upper Mississippian (early Serpukhovian) of central Russia (Moscow Basin): systematics, microstructure, and growth abnormalities: *PalZ*. <https://doi.org/10.1007/s12542-022-00636-4>.
- Verrill, A.E., 1865. Classification of polyps (extract condensed from Synopsis of the Polyps and Corals of the North Pacific Exploring Expedition under Commodore C. Ringgold and Captain John Rogers, U. S. N.): *Communications of the Essex Institute*, v. 4, p. 145–152.
- Vinn, O., 2006. Possible cnidarian affinities of *Torellella* (Hyolithelminthes, upper Cambrian, Estonia): *Paläontologische Zeitschrift*, v. 80, p. 384–389.
- Vinn, O., 2022. Phosphatic biomineralization in Scyphozoa (Cnidaria): a review: *Minerals*, v. 12, 1316. <https://doi.org/10.3390/min12101316>.
- Vinn, O., and Kirsimäe, K., 2015. Alleged cnidarian *Sphenothallus* in the Late Ordovician of Baltica, its mineral composition and microstructure: *Acta Palaeontologica Polonica*, v. 60, p. 1001–1008.
- Vinn, O., and Mironenko, A.A., 2021. Discovery of plywood structure in *Sphenothallus* from Gurovo Formation (Mississippian), central Russia: *Annales Societatis Geologorum Poloniae*, v. 91, p. 67–74.
- Yi, W., Hao, S.-G., Chen, X., Rong, J.-Y., Li, G.-X., Liu, J., and Xu, H., 2003. *Sphenothallus* from the Lower Silurian of China: *Journal of Paleontology*, v. 77, p. 583–588.
- Zhao, Y.-L., Yuan, M.-Y., Zhu, M.-Y., Guo, Q.-J., Zhuo, Z., Yang, A.-H., and Van Iten, H., 1999. The early Cambrian Taijiang Biota of Taijiang, Guizhou, PRC: *Acta Palaeontologica Sinica*, v. 38, p. 108–115.
- Zhu, M.-Y., Van Iten, H., Cox, R.S., Zhao, Y.-L., and Erdtmann, B.D., 2000. Occurrence of *Byronia* Mathew and *Sphenothallus* Hall in the Lower Cambrian of China: *PalZ*, v. 74, p. 227–238.

Accepted: 19 December 2022