

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.191>

EW0578

Outcomes in a group of 7–8-year-old children in a developmental-based intervention in autism spectrum disorder

M.T. Sindelar^{1,*}, N. Furland²

¹ Universidad Sudoeste, CS Humanas, Bahía Blanca, Argentina

² National center of research, Conicet, Biology, Bahía Blanca, Argentina

* Corresponding author.

Healthy parent-child interaction is essential for child development. Parents play a central role in the acquisition of social and communicative skills, both in typical and atypical children. Increased support for parent-mediated intervention in autism has been demonstrated. Developmental approaches for ASD are based in establishing strong interpersonal relationships through natural play to foster developmental capacities. This work reports outcomes from an intensive approach based on a developmental, individual difference, home-based intervention program with children with severe ASD. Parents were trained with DIR approach by a specialist. An individual intensive rehabilitation program was set up for each child and implemented at school, home and rehabilitation center. The intervention program also focused on semi-structured activities to promote problem solving, and meliorate sensory dysfunction. Standardized scales were administered pre- and post-intervention, with ABC). The Childhood Autism Rating Scale (CARS), the Social Communication Questionnaire (SCQ) and Vineland Adaptive Behaviours Scales.

Results Before initiating the intervention all children presented severe difficulties in communication, social interaction, lack of language, and gestures of communication, auto- and hetero-aggressive behaviors. After 6 months, children showed significant changes in mean scores for emotional functioning, communication, and daily living skills and diminution of aggressive behaviors. The present results provide strong support for the effectiveness of a developmental-based intervention, specifically in the domains of social skills behavior, social responsiveness, in a group of children with severe autism. These results highlight the positive effects of a rehabilitative approach that works in harmony with the family, school and professional team.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.192>

EW0579

Music as a helpful instrument in the treatment of children with Asd in their school inclusion program

M.T. Sindelar^{1,*}, C. Meini²

¹ Universidad Sudoeste, CS Humanas, Bahía Blanca, Argentina

² Università Vercelli, Lingua, Vercelli, Italy

* Corresponding author.

Since birth infants are active and communicative partners engaged in protoconversations with caretakers. Motherese, the simplified language adults spontaneously use with infants, has a musical structure. We believe that for developmental and evolutionistic reasons music is a preferential tool to favor communication and to promote group identity. We carried on a musical experience with a group of autistic (ASD) children aged 5 to 7 years. Each child participated at their school with 10 typically developed classmates and their teachers. Our ASD children love music and enjoy playing

and singing. With music, they overcome some communicative and social difficulties. Their bodily posture changed with music, facilitating joint attention and improvement of verbal language. When singing children learned linguistic skills, they ameliorated vowels' pronunciation and understood how a question and an answer differ in melodic contour. Taking into account the unique sensory motor profile of each ASD child, we proposed rhythmic music with high proprioceptive input (for under-reactive children) and smooth and calming music for avoidant and easily overwhelming children in order to ameliorate intentionality and enlarge circles of communication. A combination of semistructured and spontaneous activity is the main components of our approach, which has both therapeutic and educational impacts. In the musical group, all the ASDs appeared to be more attentive, motivated, better performing and able to teach their acquired skills to their peers. Typical peers interact more with children with ASD with music. We consider this very helpful in the inclusion of ASD children in a school setting.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.193>

EW0580

Impulsivity and current alcohol use in adolescents

N. Smaoui, N. Charfi*, M. Turki, M. Maâlej-Bouali, J. Ben Thabet, N. Zouari, L. Zouari, M. Maâlej

Hedi Chaker university hospital, department of psychiatry C, Hedi Chaker university hospital, Sfax, Tunisia

* Corresponding author.

Introduction Many studies have focused on the strong link between impulsivity and addictive behaviors, such as alcohol use disorders.

Objective Our study aimed to investigate the links between alcohol use and impulsivity in a sample of Tunisian pupils.

Methods This was a cross-sectional study conducted in May and June 2016. It enrolled 317 pupils from four colleges and schools in Sfax (Tunisia). The participants were asked to answer a self-administered questionnaire, after their consent. Alcohol use disorders identification test (AUDIT) was used to evaluate alcohol dependence. The Barratt Impulsiveness Scale (BIS), in its 11th version, was used to assess impulsivity trait. Its three second-order factors were attentional impulsiveness, motor impulsiveness, and non-planning impulsiveness.

Results The mean age was 16 years with a sex-ratio of 1.07. The school children reported having drunk alcohol at least once in 18.9% of cases and 41.66% of them still consume. According to AUDIT, 1.6% of alcohol users presented an alcohol misuse and 21.6% presented dependence. According to BIS, impulsivity prevalence (IP) in the whole sample was 27.8% while it was 40% among the alcoholic participants versus 24.9% in non-alcoholic participants. The average BIS score was more important in the alcoholic participants (69.25 versus 65.53; $P=0.011$). Alcohol consumption was correlated to attentional impulsiveness ($P=0.01$) and motor impulsiveness ($P=0.008$). Alcohol dependence was correlated to motor impulsiveness ($P=0.018$).

Conclusion These results show the importance of clinically evaluating and intervening on impulsive personality traits to better prevent addictive behavior such as alcoholism among adolescents.

Disclosure of interest The authors have not supplied their declaration of competing interest.

<http://dx.doi.org/10.1016/j.eurpsy.2017.02.194>

EW0581

Influence of peers drinking and parental drinking and attitudes on adolescent drinking