

CERTAIN VALUES OF COMPLETENESS AND
SATURATEDNESS OF A UNIFORM IDEAL
RULE OUT CERTAIN SIZES OF THE
UNDERLYING INDEX SET

BY
FRANTISEK FRANEK

ABSTRACT. Using the method of non-well-founded generic ultrapowers, we shall prove a generalization of a theorem of Taylor that certain values of completeness and saturatedness of a uniform ideal rule out certain sizes of the underlying index set.

1. **Introduction.** “There is no κ^+ -complete κ^+ -saturated ideal over κ^+ , κ an uncountable cardinal” is the straightforward generalization of the classical result of Ulam (see [2] or [6]) “there is no nontrivial σ -additive measure on \aleph_1 ”, proved by so-called Ulam matrices. The method of well-founded generic ultrapowers was first used by Solovay (see [4]) to prove that if “there exists a κ -complete κ -saturated ideal over κ ”, κ must be a large cardinal (badly Mahlo). Later they were extensively studied by Jech and Prikry (see [3]) in connection with precipitous ideals.

The method of non-well-founded generic ultrapowers was first used by Silver (see [5]).

Kunen observed (private communication) that using the method of well-founded generic ultrapower one can show that there is no \aleph_1 -complete \aleph_2 -saturated uniform ideal over a cardinal κ if $\aleph_\omega < \kappa < \aleph_{\omega_1}$.

Taylor (private communication) proved a generalization of this, namely “there is no \aleph_α -saturated λ^+ -complete uniform ideal over a cardinal κ if $\aleph_\lambda < \kappa < \aleph_{\lambda^+}$ and $\alpha < \lambda$ and λ is an infinite cardinal”, using some combinatorial results of Jech and Prikry. His proof is purely combinatorial.

Inspired by Kunen’s observation and using a technical insight into generic ultrapowers developed in [3], we shall prove a generalization of Taylor’s theorem with a significantly shorter proof.

2. **Definitions.** (For details, though for κ -complete ideals over κ rather than λ -complete ideals over κ , $\lambda \leq \kappa$, see [3]).

Let I be an ideal over a set S .

Then $I^+ = \{X \subset S : X \notin I\}$.

Received by the editors May 17, 1984 and, in final revised form, October 22, 1984.

AMS Subject Classification: Primary, 04A20; Secondary, 03C90, 03E40.

© Canadian Mathematical Society 1984.

$W \subset I^+$ is *I-disjoint* if $(\forall X, Y \in W)(X \cap Y \in I)$.

Let λ be a cardinal, I is λ -saturated if no $X \in [I^+]^\lambda$ is *I-disjoint*.

I is λ -complete if for any $\xi < \lambda$ and any $\{X_\alpha: \alpha \in \xi\} \subset I, U\{X_\alpha: \alpha \in \xi\} \in I$.

Let M be a transitive model of ZFC. Let $\kappa \in \text{Ord}^M$ and $\lambda \in \text{Card}^M$. $G \subset M$ is a *non-principal M - λ -complete M -ultrafilter over κ* if

- (1) $(\forall x \in G)(\forall y \in P(\kappa) \cap M)(y \supset x \Rightarrow y \in G)$;
- (2) $(\forall x \in P(\kappa) \cap M)(x \in G \text{ or } \kappa - x \in G)$;
- (3) $(\forall X \in [M]^{<\lambda} \cap M)(X \subset G \Rightarrow \cap X \in G)$;
- (4) $\cap G = \emptyset$.

If $f, g \in {}^*M \cap M$, then

$$f \in {}^*g \text{ iff } \{\alpha \in \kappa: f(\alpha) \in g(\alpha)\} \in G$$

$$f = {}^*g \text{ iff } \{\alpha \in \kappa: f(\alpha) = g(\alpha)\} \in G.$$

For every $f \in {}^*M \cap M$ let us choose (in V) a representative $[f]$ from the class $\{g \in {}^*M \cap M: g = {}^*f\}$, and form (in V) *generalized ultrapower* $\text{Ult}(M, G) = \{[f]: f \in {}^*M \cap M\}$.

Let $\text{ext}([f]) = \{[g] \in {}^*M \cap M: [g] \in {}^*[f]\}$.

For every $x \in M$ define $c_x \in {}^*M \cap M$ by $c_x(\alpha) = x$ for all $\alpha \in \kappa$. Then as usual j defined by $j(x) = [c_x]$ is an elementary embedding of M into $\text{Ult}(M, G)$ (it is often called *canonical embedding*) and (Loś theorem) $\text{Ult}(M, G) \models \phi([f_0], \dots, [f_n])$ iff $\{\alpha \in \kappa: M \models \phi(f_0(\alpha), \dots, f_n(\alpha))\} \in G$, for every formula $\phi(x_0, \dots, x_n)$ and every sequence $\langle [f_0], \dots, [f_n] \rangle \in \text{Ult}(M, G)$. In the case that \in^* is well-founded on the whole class $\text{Ult}(M, G)$, we identify $\text{Ult}(M, G)$ with its transitive collapse.

3. Preliminaries.

LEMMA 1. Let $M \subset V$ be a transitive model of ZFC. Let $G \in V$ be a non-principal M - λ -complete M -ultrafilter over κ , $\aleph_1^M \leq \lambda \leq \kappa$ cardinals in M . Let $j: M \rightarrow \text{Ult}(M, G)$ be the canonical embedding. Then

- (1) $|\alpha| \leq |\text{ext}(j(\alpha))|$ for all $\alpha \in \text{Ord}^M$;
- (2) $|\alpha| = |\text{ext}(j(\alpha))|$ (since $\text{ext}(j(\alpha)) = \{[c_\beta]: \beta \in \alpha\}$) for all $\alpha \in \lambda$;
- (3) $|\text{ext}(j(\aleph_\alpha^M))| \leq \aleph_\alpha^V$ for all $\alpha \in \lambda$;
- (4) $\{[c_\beta]: \beta \in \lambda\}$ is an initial segment of $\text{Ord}^{\text{Ult}(M, G)}$;
- (5) if G is uniform, i.e. $(\forall x \in G)(|x|^M = \kappa)$, then $|(\kappa^+)^M| \leq |\text{ext}(j(\kappa))|$.

(Note: the cardinalities are computed in V .)

PROOF. (1)–(4) follow from 2.2.2, 2.2.4, 2.2.5 and 2.3.1 in [3], when generalized from M - κ -complete M -ultrafilters over κ to M - λ -complete M -ultrafilters over κ , $\lambda \leq \kappa$.

(5) Choose, in M , a family $F \subset {}^*\kappa$ of size κ^+ of almost disjoint functions (such family always exists, see e.g. [2]). Since G is uniform, $f \neq g \in F \Rightarrow [f] \neq [g]$ as $\{\gamma \in \kappa: f(\gamma) = g(\gamma)\} \supset \beta$ for some $\beta \in \kappa$ and hence $\{\gamma \in \kappa: f(\gamma) \neq g(\gamma)\} \in G$. So $|\text{ext}(j(\kappa))| \geq |F|$. \square

NOTE. Let M be a transitive model of ZFC . Let, in M , I be an ideal over S . Let $X \subset^* Y$ mean $X - Y \in I$. One can view the poset $\langle I^+, \subset^* \rangle$ as a forcing notion. Then, if G is $\langle I^+, \subset^* \rangle$ -generic over M , we shall say that G is I -generic over M .

LEMMA 2. *Let M be a transitive model of ZFC . Let I be, in M , a λ -complete (uniform) ideal over a cardinal κ so that $\aleph_1^M \leq \lambda \leq \kappa$. Let G be I -generic over M . Then G is a non-principal M - λ -complete (uniform) M -ultrafilter over κ .*

PROOF. Easy. Left to the interested reader, or see [3]. \square

4. Main result.

THEOREM 3. *Let λ be an uncountable cardinal, $\alpha < \lambda$ and $\mu = \omega_0 \cdot \alpha$. Let $\aleph_\mu < \kappa < \aleph_\lambda$. Then there is no λ -complete \aleph_α -saturated uniform ideal over κ .*

PROOF. Assume that there are an M , a transitive model of ZFC , and I , a λ -complete \aleph_α^M -saturated uniform ideal over κ in M , and that $\alpha < \lambda$, λ is an uncountable cardinal and $\aleph_\mu^M < \kappa < \aleph_\lambda^M$ and $\mu = \omega_0 \cdot \alpha$. Let G be I -generic over M .

Since I is \aleph_α^M -saturated, \aleph_α^M is a cardinal in $M[G]$. Let $\aleph_\alpha^M = \aleph_\delta^{M[G]}$ for some $\delta \leq \alpha$. Let $\xi = \alpha - \delta$. Then $\alpha + \omega_0 \cdot \xi = \delta + \omega_0 \cdot \xi \leq \omega_0 \cdot \alpha = \mu$. Thus $\aleph_\gamma^M = \aleph_\gamma^{M[G]}$ for all $\gamma \geq \mu$. Let $\kappa = \aleph_\beta^M$ for some β . Then $\mu < \beta < \lambda$. By Lemma 1 (5) and (3) (since $\beta < \lambda$),

$$\aleph_{\beta+1}^{M[G]} = \aleph_{\beta+1}^M = |\aleph_{\beta+1}^M| \leq |\text{ext}(j(\aleph_\beta^M))| \leq \aleph_\beta^{M[G]},$$

a contradiction. \square

COROLLARY 4. *Taylor's theorem.*

PROOF. Let I be a ξ^+ -complete \aleph_α -saturated uniform ideal over a cardinal κ , $\alpha < \xi$, ξ an infinite cardinal and $\aleph_\xi < \kappa < \aleph_{\xi^+}$. Let $\lambda = \xi^+$. Let $\mu = \omega_0 \cdot \alpha$. Then $\aleph_\mu < \kappa < \aleph_\lambda$, λ is an uncountable cardinal and I is λ -complete, \aleph_α -saturated and uniform, which contradicts Theorem 3. \square

NOTE. (1) $\kappa \leq \aleph_\lambda$ is the best upper bound, for Foreman and Magidor (private communication) constructed a model with an \aleph_1 -complete \aleph_2 -saturated ideal over \aleph_{ω_1+1} .

(2) Theorem 3 gives a better lower estimate for κ than Taylor's theorem, and if ξ is weakly inaccessible, then Theorem 3 shows the non-existence of ξ^+ -complete \aleph_ξ -saturated uniform ideals over κ , $\aleph_{\omega_0 \cdot \xi} < \kappa < \aleph_{\xi^+}$, while Taylor's theorem deals only with \aleph_α -saturated ideals for $\alpha < \xi$.

REFERENCES

1. F. Franek, *Some results about saturated ideals and about isomorphisms of κ -trees*, Ph.D. thesis, University of Toronto, 1983.
2. T. Jech, *Set Theory*, Academic Press, 1978.
3. T. Jech and K. Prikry, *Ideals over uncountable sets: applications of almost disjoint functions and generic ultrapowers*, Mem. Amer. Math. Soc. **18** (1979), no. 214.

4. R. M. Solovay, *Real-valued measurable cardinals*, Axiomatic Set Theory (Proc. Sympos. Pure Math., Vol. XIII, Part I, Univ. California, Los Angeles, California, 1967), pp. 397–428.
5. J. Silver, *On the singular cardinals problem*, Proceedings of the International Congress of Mathematicians (Vancouver, BC, 1974), Vol. I, pp. 265–268.
6. S. Ulam, *Zur Masstheorie in der allgemeinen Mengenlehre*, Fund. Math. **16** (1930).

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF TORONTO
TORONTO, ONTARIO
CANADA M5S 1A1

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE
DARTMOUTH COLLEGE
HANOVER, NEW HAMPSHIRE
USA 03755