S276 Accepted posters

Restorative Justice by Proxy: A Forensic Psychiatry Case Study

Dr Suhanthini Farrell*

Greater Manchester Mental Health NHS Foundation Trust, Manchester, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.660

Aims. Restorative justice (RJ) is the process of bringing together offender and victim in the aftermath of an offence. Handled correctly, there can be substantial benefits for both parties. Communication is typically direct with an RJ facilitator from the criminal justice system, but the Crown Prosecution Service makes reference to the option of "shuttle mediation", with the RJ facilitator relaying information indirectly between parties.

The use of RJ in forensic psychiatry has been limited, with reasons including concern for the patient offender's vulnerability, impact on their mental state from the stress of facing their victim, and unpredictability of response. Adaptation of the "shuttle mediation" model with use of a trusted mental health professional may help to overcome many such obstacles.

Methods. Patient A committed his index offence of murder at age 22. During his mandatory life sentence, he developed schizophrenia and was transferred to a secure hospital. His illness proved treatment resistant and many years passed before he was discharged to the community on life licence following Parole Board release. Although generally stable, he was easily overwhelmed by stressful situations, in which he could struggle to articulate his thoughts.

His victim's wife and daughter enquired about the possibility of restorative justice. A's probation officer liaised with A's consultant psychiatrist. A was considered unsuitable for direct approach, but following discussion with an RJ facilitator, a proxy model was agreed upon. This was based on shuttle mediation but with the consultant psychiatrist acting as shuttle, or proxy, for A.

The aims of the family were elicited in advance. Capacitous consent to information sharing was obtained from A, with boundaries of what would be shared clearly agreed. Information A wished to share with the family was established, including verbatim quotes. **Results.** Following the RJ meeting, feedback from the victim's family and A was strongly positive. The victim's family reported a sense of peace and closure from the experience. A's long-standing anxiety, with feelings of guilt and self-blame reduced. Both parties expressed gratitude for the meeting and did not find the use of a proxy negated the benefits of the experience.

Conclusion. This case report demonstrates that a proxy model of restorative justice is feasible and can be successful for patient offenders with severe and enduring mental disorder. The potential benefits to patient and victim suggest that it could be considered more widely as an option in forensic psychiatry settings.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard BJPsych Open peer review process and should not be quoted as peer-reviewed by BJPsych Open in any subsequent publication.

Case Study of Traumatised Sub-Postmasters

Dr Paul Foster*

Psychiatric & Psychological Consultant Services, London, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.661

Aims. The scandal of sub-postmasters wrongly accused by the post office of offences relating to the faulty IT Horizon system is of ongoing media prominence.

Since May 2021 I have undertaken personal injury medico legal assessments at the request of a solicitor representing those falsely accused and convicted of offences by the post office. Their convictions had been quashed in the court of appeal.

Methods. I interviewed fourteen individuals, age range 35 to 70 years, five women and nine men. Ten had brought claims for Malicious Prosecution. The remaining four sub-postmasters were part of the historical financial shortfalls scheme set up for those who had not been prosecuted but were applying for compensation.

Of the ten convicted sub-postmasters, four spent periods in prison and the remaining six were given suspended or community sentences. Those in prison were often separated from young children by distance or withholding family members.

I assessed to what extent and in what way their mental health and that of their families had been adversely affected as a result of their experiences.

Assessments were undertaken remotely, including speaking with a family member.

ICD 10 diagnostic criteria were used.

Results. All of the cases revealed evidence of psychopathology at the time of the allegations or convictions and continued to varying degrees subsequently.

A diagnosis of PTSD was made in five cases, Adjustment Disorder in 2 cases, Dysthymia in one case and depressive illness in eight cases. In only four cases had the disorders resolved.

Four of the five cases of PTSD had evidence of a past psychiatric history prior to becoming sub-postmasters. Past history included depression, impulsive overdose, eating disorder, problems related to gambling and abuse of alcohol and cocaine.

Both sub-postmasters who had been accused of taking money due to faulty software and those who had been wrongly convicted had high rates of psychopathology.

These findings are consistent with the only other study of the psychological effects on sub-postmasters which found high rates of psychopathology in both accused and convicted individuals.

Existing diagnostic criteria were limited in capturing the suffering of individuals who had endured such complex trauma so a narrative description including the effects on family members was also used.

Conclusion. This study of the mental health of falsely accused sub-postmasters demonstrates a high degree of psychopathology which may require therapeutic intervention.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard BJPsych Open peer review process and should not be quoted as peer-reviewed by BJPsych Open in any subsequent publication.

Imaginator

Dr Emily Gardner-Bougaard^{1*}, Ms Athina Sevri² and Dr Martina Do Simplicio²

¹London, United Kingdom and ²Imperial, London, United Kingdom *Presenting author.

doi: 10.1192/bjo.2024.662

Aims. Self-harm affects around 20% of all young people in the UK. Treatment options for self-harm remain limited and those available are either non-specific or long and costly and may not suit all young people. There is an urgent need to develop new scalable interventions to address this gap.

Imaginator is a novel imagery-based intervention targeting self-harm initially developed for 16–25-year-olds. It is a blended digital intervention delivering Functional Imagery Training (FIT) via therapist sessions and a smartphone app. In this study