

# Differences in Neurocognitive Impairment Among HIV-Infected Latinos in the United States

María J. Marquine,<sup>1</sup> Anne Heaton,<sup>1</sup> Neco Johnson,<sup>2</sup> Monica Rivera-Mindt,<sup>3</sup> Mariana Cherner,<sup>1</sup> Cinnamon Bloss,<sup>1</sup> Todd Hulgan,<sup>4</sup> Anya Umlauf,<sup>1</sup> David J. Moore,<sup>1</sup> Pariya Fazeli,<sup>5</sup> Susan Morgello,<sup>6</sup> Donald Franklin Jr.,<sup>1</sup> Scott Letendre,<sup>7</sup> Ron Ellis,<sup>8</sup> Ann C. Collier,<sup>6</sup> Christina M. Marra,<sup>9</sup> David B. Clifford,<sup>9,10</sup> Benjamin B. Gelman,<sup>11</sup> Ned Sacktor,<sup>12</sup> David Simpson,<sup>6</sup> J. Allen McCutchan,<sup>7</sup> Igor Grant,<sup>1</sup> AND Robert K. Heaton<sup>1</sup>

<sup>1</sup>Department of Psychiatry, University of California, San Diego, California

<sup>2</sup>Advancing Diversity through Aging Research (ADAR) Program, San Diego State University, San Diego, California

<sup>3</sup>Department of Psychology, Fordham University, Bronx, New York

<sup>4</sup>Department of Medicine, Vanderbilt University, Nashville, Tennessee

<sup>5</sup>Department of Psychology, University of Alabama Birmingham, Birmingham, Alabama

<sup>6</sup>Department of Neurology, Mount Sinai Medical Center, Miami, Florida

<sup>7</sup>Department of Medicine, University of California, San Diego, California

<sup>8</sup>Department of Neurosciences, University of California, San Diego, California

<sup>9</sup>Department of Neurology, University of Washington, Seattle, Washington

<sup>10</sup>Department of Medicine, University of Washington, Seattle, Washington

<sup>11</sup>Department of Pathology, University of Texas Medical Branch, Galveston, Texas

<sup>12</sup>Department of Neurology, Johns Hopkins University, Baltimore, Maryland

(RECEIVED February 14, 2017; FINAL REVISION July 11, 2017; ACCEPTED July 14, 2017; FIRST PUBLISHED ONLINE September 6, 2017)

## Abstract

**Objectives:** Human immunodeficiency virus (HIV) disproportionately affects Hispanics/Latinos in the United States, yet little is known about neurocognitive impairment (NCI) in this group. We compared the rates of NCI in large well-characterized samples of HIV-infected (HIV+) Latinos and (non-Latino) Whites, and examined HIV-associated NCI among subgroups of Latinos. **Methods:** Participants included English-speaking HIV+ adults assessed at six U.S. medical centers (194 Latinos, 600 Whites). For overall group, age:  $M = 42.65$  years,  $SD = 8.93$ ; 86% male; education:  $M = 13.17$ ,  $SD = 2.73$ ; 54% had acquired immunodeficiency syndrome. NCI was assessed with a comprehensive test battery with normative corrections for age, education and gender. Covariates examined included HIV-disease characteristics, comorbidities, and genetic ancestry. **Results:** Compared with Whites, Latinos had higher rates of global NCI (42% vs. 54%), and domain NCI in executive function, learning, recall, working memory, and processing speed. Latinos also fared worse than Whites on current and historical HIV-disease characteristics, and nadir CD4 partially mediated ethnic differences in NCI. Yet, Latinos continued to have more global NCI [odds ratio (OR) = 1.59; 95% confidence interval (CI) = 1.13–2.23;  $p < .01$ ] after adjusting for significant covariates. Higher rates of global NCI were observed with Puerto Rican ( $n = 60$ ; 71%) versus Mexican ( $n = 79$ , 44%) origin/descent; this disparity persisted in models adjusting for significant covariates (OR = 2.40; CI = 1.11–5.29;  $p = .03$ ). **Conclusions:** HIV+ Latinos, especially of Puerto Rican (vs. Mexican) origin/descent had increased rates of NCI compared with Whites. Differences in rates of NCI were not completely explained by worse HIV-disease characteristics, neurocognitive comorbidities, or genetic ancestry. Future studies should explore culturally relevant psychosocial, biomedical, and genetic factors that might explain these disparities and inform the development of targeted interventions. (*JINS*, 2018, 24, 163–175)

**Keywords:** Hispanics, Human immunodeficiency virus, Culture, Cognitive function, Minority health, Health status disparities

## INTRODUCTION

Despite the availability of virally suppressive antiretroviral therapy (ART), human immunodeficiency virus (HIV)-

associated neurocognitive impairment (NCI) continues to be common and impactful (Heaton et al., 2011). NCI is observed in up to half of HIV-infected (HIV+) persons (Heaton et al., 2011), and is associated with worse every day functioning outcomes (e.g., medication nonadherence, driving problems, unemployment; Gorman, Foley, Ettenhofer, Hinkin, & van Gorp, 2009; Heaton et al., 2004; Thames, Arentoft, Rivera-Mindt, & Hinkin, 2013).

Correspondence and reprint requests to: María J. Marquine, Department of Psychiatry, UCSD School of Medicine, 9500 Gilman Drive, La Jolla, CA 92093-0603. E-mail: mmarquine@ucsd.edu

HIV is an especially significant public health concern for Hispanics/Latinos/as, who represent the largest ethnic/racial minority group in the United States (U.S. Census Bureau, 2015). Hispanics/Latinos/as are three-times more likely to have HIV infection compared with non-Hispanic Whites (Centers for Disease Control and Prevention, 2015), hereafter referred to as “Latinos” and “Whites,” respectively. Latinos tend to be disadvantaged at nearly every component of the HIV disease treatment cascade (Gardner, McLees, Steiner, del Rio, & Burman, 2011), including delayed HIV testing and delayed care engagement (Chen, Gallant, & Page, 2012; Dennis, Napravnik, Sena, & Eron, 2011; Shapiro et al., 1999; Sheehan, Trepka, & Dillon, 2015; Turner et al., 2000). Thus, not surprisingly, Latinos tend to present with worse HIV disease characteristics, including higher rates of acquired immunodeficiency syndrome (AIDS) and opportunistic infections, and lower CD4 cell counts than Whites (Swindells et al., 2002).

Latinos also might be at increased risk for HIV-associated NCI in the era of combination ART (Durvasula, Miller, Myers, & Wyatt, 2001; Heaton et al., 2015; Rivera-Mindt et al., 2008, 2014; Wojna et al., 2006). In a recent longitudinal study of HIV+ persons in the CNS HIV Anti-Retroviral Therapy Effects Research (CHARTER) study, Latino ethnicity was a significant, independent predictor of neurocognitive decline over an average of three years (Heaton et al., 2015). While the latter study was not focused on examining ethnic differences, and thus statistical models were not specifically constructed to address this question, a small number of studies have focused on investigating HIV-associated NCI among Latinos (Rivera-Mindt et al., 2008, 2014; Wojna et al., 2006). A study of 49 HIV+ women with nadir CD4 cell counts of  $<500$  cells/mm<sup>3</sup> living in Puerto Rico found elevated rates of NCI in this group compared with other HIV+ cohorts (global NCI = 78%; Wojna et al., 2006).

Two additional studies investigated NCI among HIV+ Latinos primarily of Caribbean (i.e., Puerto Rican/Dominican) origin/descent and living in the northeastern United States (i.e., New York). One of these studies included participants in the Manhattan HIV Brain Bank, with advanced HIV disease (51 Latino and 49 White; Rivera-Mindt et al., 2008). Findings indicated Latinos performed worse than Whites on tests of abstraction/executive function in a model adjusting for socioeconomic status, English reading level (i.e., Wide Range Achievement Test, WRAT-3), and CD4 cell counts. However, there were no significant ethnic group differences in similar adjusted models on global neurocognitive function, learning, and attention/working memory.

Another study in the northeastern United States (Rivera-Mindt et al., 2014) found worse global and domain (processing speed, learning, and memory) neurocognitive performance among older (50+ years) HIV+ Latinos ( $n = 22$ ) compared with older HIV+ Whites ( $n = 22$ ), but no significant differences in younger HIV+ Latinos ( $n = 62$ ) and Whites ( $n = 20$ ). Taken together, results from these studies indicate that at least some subgroups of Latinos might be at increased risk for

HIV-associated NCI. However, none of the studies to date has examined NCI among large groups of Latinos, which would allow for comparisons among subgroups of Latinos.

Latinos are highly heterogeneous, comprising multiple national origins, racial groups, patterns of immigration, and linguistic and educational backgrounds (Guarnaccia et al., 2007; Llorente, 2008). Latinos are also known to possess genetic backgrounds shaped by admixture from several source continental populations, with often the precise contribution from each source population unknown (Bryc, Durand, Macpherson, Reich, & Mountain, 2015; Conomos et al., 2016; Libiger & Schork, 2013). Recent findings from a large population-based study of Latino health, indicate differential neurocognitive performances among subgroups of middle-aged and older Latinos (González et al., 2015).

Furthermore, there are documented differences in the HIV care continuum among subgroups of Latinos based on their country of origin (Chen et al., 2012; Lee & Held, 2015; Sheehan et al., 2015), which might have important implications for both general health and mental health care use and outcomes, including NCI. Latinos born in Mexico have an increased risk for late HIV diagnosis (Sheehan et al., 2015), and HIV+ Latinos of Puerto Rican origin/descent are more likely to use services compared with other HIV+ Latino subgroups (Lee & Held, 2015; Sheehan et al., 2015). Yet, HIV+ Puerto Ricans are at increased risk of early mortality (Sheehan et al., 2015). They also have the highest rates of HIV transmission through injection drug use of all Latino subgroups (Chen et al., 2012), which is linked to worse retention in care, ART adherence, and viral suppression (Gray, Valverde, Tang, Siddiqi, & Hall, 2015). While these findings suggest that there are important differences among HIV+ Latino subgroups, we do not know whether these differences translate to differential rates of NCI.

There were two main purposes to the present study. First, we aimed to investigate differences in rates and correlates of NCI in HIV+ Latinos and Whites, and expand on prior studies by: (a) including a large cohort of participants, who were broadly representative of HIV+ persons receiving care at academic medical centers in the United States, and (b) by examining the role of HIV disease characteristics and comorbidities on ethnic differences in HIV-associated NCI. We hypothesized that HIV+ Latinos would show increased rates of NCI compared with HIV+ Whites.

The second aim was to investigate differences in rates and correlates of NCI among U.S. Latino subgroups based on self-reported country of origin/descent, that is, Puerto Rico and Mexico. These were the two Latino subgroups for whom we had considerable data available and they also represent the two largest subgroups of Latinos in the United States (Ennis, Rios-Vargas, & Albert, 2011). We refer to these subgroups as Puerto Rican and Mexican for simplicity, but Latinos were living in the continental United States at the time of study participation, and we did not have consistent data on country of birth or duration of residence in the continental United States. In addition to these two main aims, we were also interested in investigating the role of genetically

defined ancestry on ethnic group differences in NCI among a subgroup of participants with available genetic data. Better characterization of NCI among Latino subgroups is key for the development of targeted, culturally relevant interventions aimed at preventing and/or ameliorating NCI disparities in this large, vulnerable, heterogeneous, and often understudied segment of the U.S. population.

## METHODS

### Participants

The sample included 794 HIV+ adults (194 Latinos and 600 Whites) who were participants in the CHARTER study (Heaton et al., 2010, 2015), a longitudinal multisite cohort study of HIV+ persons. All participants were fluent in English, as ascertained by participant self-report and determined by study staff. Only baseline data were used for present analyses. Study visits took place between 2002 and 2010, with 98% of study visits occurring by 2007. Inclusion criteria were self-identifying as Latino or White, and having neurocognitive data.

### Materials and Procedures

Assessments were performed by trained staff, who were certified by the coordinating center (UC San Diego). All data were obtained in compliance with the Helsinki Declaration and Human Research Protections Programs at the study sites.

#### *Demographic characteristics*

Demographic information (age, years of education, gender, ethnicity/race) was obtained *via* self-report. Self-reported ethnicity was ascertained following NIH guidelines, which define “Hispanic/Latino” as a person of Mexican, Puerto Rican, Cuban, South or Central American, or other Spanish-speaking culture of origin regardless of race (Office of Management and Budget, 1997). Per this methodology, participants were asked whether they were of Spanish/Hispanic/Latino origin, and provided with the following response options: (1) No, not Spanish/Hispanic/Latino; (2) Yes, Mexican, Mexican American, Chicano; (3) Yes, Puerto Rican; (4) Yes, Cuban; and (5) Yes, Other.

#### *Neuromedical evaluation*

Participants underwent a comprehensive neuromedical evaluation that included assessment of medical history, structured medical and neurological examinations, and the collection of blood, cerebrospinal fluid (CSF), and urine samples, as described in Heaton and colleagues (2010). Central nervous system (CNS) penetration effectiveness (CPE) was included as an estimate of the degree of CNS penetration of current ART regimen (Letendre, 2011; Letendre et al., 2008). Neurocognitive comorbidity status was ascertained by pre-established international

criteria outlined in Antinori and colleagues (2007). In short, comorbidity status was based on self-reported history of HIV-related opportunistic CNS conditions and unrelated developmental, psychiatric, and neuromedical confounds, and categorized as incidental, contributing, or confounding.

Details about the procedure and its inter-rater reliability in the CHARTER study are available in Heaton and colleagues (2010). The Veterans Aging Cohort Study (VACS) Index score was computed as described previously (Justice et al., 2012, 2013). It is a composite marker of disease severity in HIV, which includes “traditional” HIV biomarkers (current CD4 count and HIV-1 plasma RNA), and non-HIV biomarkers [indicators of renal and liver function, anemia, and hepatitis C virus (HCV) co-infection]. HCV co-infection was ascertained based on HCV serostatus (in 83% of the sample) or self-report.

#### *Neurocognitive evaluation*

Participants completed a comprehensive neurocognitive battery assessing seven cognitive domains (see Heaton et al., 2010, for a list of tests by domain). Raw scores for each test were converted to T-scores adjusting for demographic characteristics (age, education, gender). T-scores were then converted to deficit scores ranging from 0 (no impairment) to 5 (severe impairment). Deficit scores for tests comprising each domain were averaged to compute domain deficit scores. Deficit scores from all tests were averaged to compute Global Deficit Scores that reflect the number and severity of deficits across the entire neurocognitive battery (Carey et al., 2004). Presence of domain and global NCIs were defined as domain deficit score  $>.5$  and global deficit score  $\geq .5$ , respectively. Two Processing Speed tests in the battery (i.e., Digit Symbol and Symbol Search from the Wechsler Adult Scale Intelligence Scale-Third Edition; WAIS-II; Wechsler, 1997) have Latino ethnicity-corrected T-scores available (Heaton, Taylor, & Manly, 2002), and T-scores on these tests were also examined separately. All other tests did not include adjustments for ethnicity.

#### *Psychiatric and substance use characteristics*

Current (past 30 days) and lifetime histories of major depressive and substance use disorders were evaluated using the computer-assisted Composite International Diagnostic Interview (Wittchen et al., 1991), which follows Diagnostic and Statistical Manual-Fourth Edition criteria (American Psychiatric Association, 1994). Substance use disorder was determined as meeting criteria for substance abuse or dependence for alcohol, marijuana, cocaine, hallucinogens, inhalants, methamphetamine, opioids, phencyclidine, sedatives, or other substances. History of injection drug use was ascertained by self-report. Current mood was assessed using the Beck Depression Inventory-II (BDI-II; Beck, Steer, & Brown, 1996), and recent substance use was identified *via* urine toxicology for various substances (i.e., marijuana, cocaine, methamphetamine, amphetamine,

barbiturates, benzodiazepines, and phencyclidine) on the day of testing.

### Genotyping and genetic ancestry

Genome-wide genotype information was available on a subset of participants (127 Latinos and 419 Whites) from whom genomic DNA was isolated from whole blood and subjected to genotyping using the Affymetrix Genome-Wide Human SNP Array 6.0 platform (Affymetrix, Inc., Santa Clara, CA). These data were subjected to rigorous quality control analyses with exclusion of variants with less than 95% genotyping efficiency or minor allele frequencies less than 1% in the study population.

To assess genetic ancestry based on genotype information for this subset of participants, the method developed by Libiger and Schork (2013) was used. Specifically, an ancestry-informative reference panel was constructed by bringing together genotype data from 2513 individuals of known ancestry from 63 populations around the world using several publically available sources (Cann et al., 2002; International HapMap 3 Consortium et al., 2010; Nelson et al., 2008; Xing et al., 2009). The reference panel was created in a stepwise manner to ensure that the included individuals were not admixed among six major continental populations (African, Central Asian, East Asian, European, Native American, and Oceanic) and that each continental population was represented by a reasonably large number of diverse individuals originating in the relevant continent. The assembled reference panel contained genotype information at 16,433 strand-unambiguous SNPs. These markers exhibited low linkage disequilibrium ( $r$ -squared less than 0.1 was observed between 99% of marker pairs), and allele frequency was higher than 1%.

To assess ancestry and admixture proportions a supervised clustering approach implemented in the ADMIXTURE software (Alexander, Novembre, & Lange, 2009) probabilistically assigned each participant to six clusters corresponding to the six major continental populations listed above. The genotype profiles of the six populations were defined by the individuals who made up the reference panel. Based on this approach, the degree of ancestry of each participant was determined, effectively quantifying the proportion of their genome that is likely to be derived from each of the six populations (African, Central Asian, East Asian, European, Native American, and Oceanic). Major genetic ancestry was defined as having a genetic ancestry proportion of  $\geq .5$  for a given region. Participants with a hybrid pattern of ancestries (all with frequencies  $< .5$ ) were assigned to “no single major ancestry.”

### Statistical Analyses

To investigate differences between Latinos and Whites, we conducted independent sample  $t$  tests and Chi-square tests on demographic, HIV disease, medical and psychiatric comorbidity, genetic ancestry, and global and domain NCI by

group. To examine whether differences in rates of NCI might be explained by HIV disease characteristics and comorbidities we ran a logistic regression model on global and domain NCI, including as predictors ethnic group and significant covariates. Covariates examined included HIV disease characteristics, and medical and psychiatric comorbidity (Table 1). To build the models, we entered covariates that were significantly different by ethnic group at  $p < .10$  into a stepwise backward regression models (minimum Akaike information criterion) on global NCI. Variables selected *via* this method were entered in final multivariable models on global and domain NCI. We also examined the impact of these covariates on ethnic differences in NCI *via* path analyses (for continuous mediators, see Preacher & Hayes, 2004; for categorical mediators, see Imai, Keele, & Tingley, 2010).

To investigate the impact of other HIV-disease characteristics relevant to persons on ART (detectable plasma and CSF RNA, and CPE), we ran a logistic regression model on global NCI within our subset of participants on ART, including significant covariates yielded by the overall model in addition to ART-relevant variables that differed significantly between groups. To examine predictors of global NCI by ethnic group, we first ran univariable analyses investigating the association of each of the HIV disease characteristics and psychiatric and medical comorbidities on global NCI separately by ethnic group. We then ran separate multivariable logistic regression models by ethnic group with predictors being all the variables that were associated with global NCI in univariable analyses at  $p < .10$ . To examine differences and associations among subgroups of Latinos, we conducted similar analyses comparing Puerto Ricans and Mexicans.

To examine whether factors possibly linked to genetic ancestry might explain differences in global NCI by ethnic group (and country of origin/descent) among HIV+ persons, we ran a logistic regression model on global NCI among our subset of participants with available genetic ancestry data, including self-reported ethnic (or country of origin) group and major genetic ancestry as predictors. We also ran separate univariable analyses examining the association of proportion of genetic ancestry from each region with global NCI by ethnic and country of origin group.

## RESULTS

### Cohort Characteristics by Ethnic Group

Table 1 shows characteristics of the study cohort by ethnic group. Compared with Whites, Latinos were slightly younger, had less formal education, and were more likely to be female. All participants who self-identified as White had a European major genetic ancestry. As expected, there was considerable variability in major genetic ancestry within the self-identified Latino group. Latinos were recruited primarily from sites in New York (NY), San Diego (CA), and

**Table 1.** Characteristics of the study cohort by ethnic group

Variable	Whites ( <i>n</i> = 600)	Latinos ( <i>n</i> = 194)	<i>p</i> <sup>a</sup>
<b>Demographics</b>			
Age, <i>M</i> ( <i>SD</i> )	43.07 (9.14)	41.33 (8.14)	.012
Education, <i>M</i> ( <i>SD</i> )	13.39 (2.69)	12.50 (2.77)	<.001
Gender, <i>n</i> (%) male	527 (88)	152 (78)	<.01
<b>Major genetic ancestry, <i>n</i> (%)<sup>b</sup></b>			
Europe	419 (100)	82 (65)	<.001
Africa	0 (0)	15 (12)	
American Indian	0 (0)	5 (4)	
East Asian	0 (0)	1 (1)	
No single major ancestry	0 (0)	24 (19)	
<b>Study site, <i>n</i> (%)</b>			
New York (NY)	61 (10)	61 (31)	<.001
San Diego (CA)	165 (28)	55 (28)	
Galveston (TX)	99 (17)	41 (21)	
Seattle (WA)	154 (26)	22 (11)	
St Louis (MO)	105 (18)	9 (5)	
Baltimore (MD)	16 (3)	6 (3)	
<b>HIV disease characteristics</b>			
AIDS, <i>n</i> (%)	310 (52)	115 (61)	.04
Estimated duration of infection (years), <i>M</i> ( <i>SD</i> )	9.44 (6.72)	9.99 (6.57)	.32
Current CD4, median (IQR)	471 (315, 647)	402 (258, 585)	<.01
Nadir CD4, median (IQR)	192 (60, 317)	150 (40, 281)	.04
On ART	406 (69)	140 (76)	.07
% detectable plasma <sup>c</sup>	157 (39)	74 (54)	<.01
% detectable CSF <sup>c</sup>	29 (9)	20 (18)	.02
CNS penetration effectiveness <sup>c</sup> , <i>M</i> ( <i>SD</i> )	7.53 (1.90)	7.66 (1.89)	.50
Months exposure to ART, median (IQR)	48.59 (6.67, 96.98)	59.33 (8.53, 97.05)	.79
VACS Index, median (IQR)	15 (7, 24)	16 (10, 27)	.08
<b>Comorbidity status, <i>n</i> (%)</b>			
Incidental	367 (61)	100 (53)	.11
Contributing	157 (26)	61 (32)	
Confounding	75 (13)	29 (15)	
<b>HCV co-infection, <i>n</i> (%)</b>			
HCV co-infection, <i>n</i> (%)	110 (18)	36 (19)	.90
<b>Psychiatric characteristics, <i>n</i> (%)</b>			
Current major depressive disorder	100 (17)	31 (16)	.86
Lifetime major depressive disorder	354 (59)	97 (51)	.04
Current any substance use disorder	25 (4)	7 (4)	.74
Lifetime any substance use disorder	433 (73)	128 (67)	.15
Lifetime alcohol use disorder	349 (58)	96 (50)	<.05
Lifetime cannabis use disorder	129 (22)	30 (16)	.07
Lifetime other substance use disorder	289 (48)	101 (53)	.28
Injection drug use ever	160 (27)	49 (25)	.70
Urine toxicology positive	76 (13)	31 (17)	.18
Beck Depression Inventory, <i>M</i> ( <i>SD</i> )	14.93 (11.19)	14.10 (9.88)	.33

Note. CSF = cerebrospinal fluid; CNS = central nervous system; IQR = interquartile range; HCV = hepatitis C virus; VACS = Veterans Aging Cohort Study.


<sup>a</sup>Based on independent sample t-tests and Chi-Square tests.

<sup>b</sup>Major genetic ancestry refers to having a genetic ancestry proportion of  $\geq .5$  for a given region (Africa, Europe, America, East Asia, Central Asia, and Oceania), with participants that have all frequencies  $< .5$  being assigned to "no single major ancestry"; Whites: *n* = 419; Latinos: *n* = 127.

<sup>c</sup>Among those on ART (antiretroviral therapy).

Galveston (TX). Latinos were more likely to have AIDS, had lower nadir and current CD4 cell counts, and among those on ART, were more likely to have detectable HIV RNA in plasma and CSF (failure of viral suppression on ART). There were no significant differences on other HIV disease

characteristics or overall comorbidity ratings, except for a trend for higher VACS Index scores (indicative of worse health) among Latinos. Regarding psychiatric characteristics, Latinos were somewhat less likely to have a lifetime history of major depressive and alcohol use disorders.



**Fig. 1.** Rates of global and domain neurocognitive impairment among HIV+ Latinos and Whites. \* $p < .01$ ; \*\* $p < .001$ .

### Neurocognitive Impairment by Ethnic Group

Figure 1 shows results from ethnic group comparisons on global and domain NCI. In unadjusted analyses, Latinos had higher rates of global NCI than Whites [odds ratio (OR) = 1.61; confidence interval (CI) = 1.16–2.23;  $p < .01$ ]. Within neurocognitive domains, Latinos had higher rates of NCI on executive function, working memory, learning and memory/delayed recall, and speed of information processing.

HIV disease characteristics and medical and psychiatric comorbidities, which were significantly different by ethnic group at  $p < .10$  (see Table 1), were entered into a backward stepwise regression model on global NCI along with ethnicity. This method selected current and nadir CD4, and the VACS Index as significant covariates. After including these variables, ethnic differences in global NCI remained significant (OR = 1.59; 95% CI = 1.13–2.23;  $p < .01$ ). Path analyses showed nadir CD4 (indirect effect = 0.049; 95% CI = 0.006–0.121) was the only significant partial mediator.

Multivariable models adjusting for significant covariates on neurocognitive domains, revealed that ethnic group differences were significant for executive function (OR = 1.54; 95% CI = 1.10–2.15;  $p = .01$ ), working memory (OR = 1.98; 95% CI = 1.39–2.81;  $p < .001$ ), learning (OR = 1.49; 95% CI = 1.07–2.09;  $p = .02$ ), and delayed recall (OR = 1.44; 95% CI = 1.03–2.03;  $p = .03$ ), but not for speed of information processing (OR = 1.40; 95% CI = 0.96–2.02;  $p = .08$ ). Within the subset of participants on ART, unadjusted models showed increased rates of NCI among Latinos than Whites (OR = 1.92; 95% CI = 1.30–2.85;  $p < .001$ ). A model including significant covariates identified by the overall model, in addition to detectable plasma and CSF RNA, again revealed increased rates of global NCI among Latinos than Whites (OR = 1.80; 95% CI = 1.14–2.85;  $p = .01$ ).

### Correlates of NCI by Ethnic Group

Among Latinos, results from univariable models showed that lower nadir CD4, being on ART, and increased comorbidity

status were all significant predictors of global NCI ( $p < .001$ ). Results from a multivariable model showed that being on ART (ON–OFF: OR = 2.30; 95% CI = 1.02–5.35;  $p = .04$ ) and comorbidity status (confounding-incidental: OR = 7.77; 95% CI = 2.86–25.29;  $p < .001$ ; contributing-incidental: OR = 3.21; 95% CI = 1.61–6.61;  $p < .001$ ) were both significant independent predictors of increased risk for global NCI among Latinos.

Among Whites, univariable analyses showed that AIDS, lower nadir CD4, more months of ART exposure, increased comorbidity status, and depressed mood (BDI scores) were significantly associated with global NCI ( $p < .01$ ). In a multivariable model, comorbidity status was the only significant predictor of increased risk of global NCI among Whites (confounding-incidental: OR = 10.45; 95% CI = 5.69–20.41;  $p < .001$ ; contributing-incidental: OR = 2.81; 95% CI = 1.89–4.19;  $p < .001$ ; confounding-contributing: OR = 3.72; 95% CI = 1.94–7.48;  $p < .001$ ).

### Differences Among Subgroups of Latinos by Country of Origin/Descent

There was considerable heterogeneity within our Latino group in regards to self-reported country of origin/descent (79 Mexican, 60 Puerto Rican, 6 Cuban, 18 South or Central American, 29 from another Spanish culture or origin, and 2 did not respond). Analyses comparing Puerto Ricans and Mexicans (Table 2) showed that Puerto Ricans were significantly older and more likely to be female. A majority of individuals in both groups had a European major genetic ancestry, but Puerto Ricans were somewhat more likely to have African as their major genetic ancestry, and Mexicans were more likely to have no single major genetic ancestry.

Most Puerto Ricans were tested in New York, and most Mexicans were tested in San Diego and Galveston (Table 2). Puerto Ricans had longer duration of infection and more months of exposure to ARTs. Among those on ARTs, Puerto Ricans also were more likely to have detectable HIV RNA in CSF (i.e., failure to suppress HIV replication in the CNS). Furthermore, Puerto Ricans had higher VACS Index scores and were more likely to have contributing and confounding comorbidity ratings (as compared with incidental) than Mexicans. Puerto Ricans were also more likely to test positive for illicit substances on the date of testing based on urine toxicology.

As Figure 2 shows, Puerto Ricans had significantly higher rates of global NCI than Mexicans, and this difference was primarily driven by the domains of learning and memory/delayed recall (with significant differences for both verbal and visual tests of learning/recall,  $ps < .05$ ).

HIV disease characteristics, and medical and psychiatric comorbidities which were significantly different by country of origin group at  $p < .10$  (see Table 2) were entered into a backward stepwise regression model on global NCI as predictors along with country of origin. This method selected ART and comorbidity status as covariates. Multivariable

**Table 2.** Characteristics of Latinos by country of origin/descent (Mexican and Puerto Rican)

	Mexican ( <i>n</i> = 79)	Puerto Rican ( <i>n</i> = 60)	<i>p</i> -Value <sup>a</sup>
<b>Demographics</b>			
Age, <i>M</i> ( <i>SD</i> )	40.15 (8.48)	43.87 (6.95)	<.01
Education, <i>M</i> ( <i>SD</i> )	12.27 (2.55)	11.98 (2.68)	.51
Gender, <i>n</i> (%) male	69 (87)	37 (62)	<.001
<b>Major genetic ancestry, <i>n</i> (%)<sup>b</sup></b>			
Europe	41 (66)	23 (70)	—
Africa	2 (3)	6 (18)	—
American Indian	3 (5)	0 (0)	—
East Asian	1 (2)	0 (0)	—
No single major ancestry	15 (24)	4 (12)	—
<b>Study site, <i>n</i> (%)</b>			
New York (NY)	2 (3)	47 (78)	<.001
San Diego (CA)	36 (46)	5 (8)	
Galveston (TX)	26 (33)	1 (2)	
Seattle (WA)	10 (13)	3 (5)	
St Louis (MO)	4 (5)	2 (3)	
Baltimore (MD)	1 (1)	2 (3)	
<b>HIV disease characteristics</b>			
AIDS, <i>n</i> (%)	47 (59)	38 (63)	.65
Estimated duration of infection (years), <i>M</i> ( <i>SD</i> )	8.56 (6.53)	12.57 (4.99)	<.001
Current CD4, median (IQR)	416 (274, 588)	394 (254, 506)	.40
Nadir CD4, median (IQR)	171 (47, 279)	145 (25, 258)	.36
On ART	55 (71)	48 (84)	.06
<i>n</i> (%) detectable plasma <sup>c</sup>	26 (48)	26 (54)	.54
<i>n</i> (%) detectable CSF <sup>c</sup>	6 (12)	11 (34)	.03
CNS penetration effectiveness <sup>c</sup> , <i>M</i> ( <i>SD</i> )	7.38 (1.57)	7.80 (2.69)	.34
Months exposure to ART, median (IQR)	50.71 (46.97)	73.50 (47.63)	<.01
VACS Index, median (IQR)	13 (7, 22)	22 (12, 37)	<.01
<b>Comorbidity status, <i>n</i> (%)</b>			
Incidental	48 (61)	22 (37)	—
Contributing	25 (32)	24 (40)	—
Confounding	6 (8)	14 (23)	—
<b>HCV co-infection, <i>n</i> (%)</b>			
14 (18)		16 (27)	.21
<b>Psychiatric characteristics, <i>n</i> (%)</b>			
Current major depressive disorder	13 (17)	10 (17)	.97
Lifetime major depressive disorder	42 (54)	29 (49)	.59
Current any substance use disorder	5 (6)	2 (3)	.42
Lifetime any substance use disorder	56 (72)	38 (64)	.33
Lifetime alcohol use disorder	46 (59)	26 (44)	.08
Lifetime cannabis use disorder	11 (14)	11 (19)	.47
Lifetime other substance use disorder	41 (53)	30 (51)	.84
Injection drug use ever	25 (32)	11 (18)	.07
Urine toxicology positive	5 (7)	16 (27)	<.01
Beck Depression Inventory, <i>M</i> ( <i>SD</i> )	14.00 (9.62)	13.93 (9.45)	.97

Note. HIV = human immunodeficiency virus; AIDS = acquired immunodeficiency syndrome; ART = antiretroviral therapy; IQR = interquartile range; CSF = cerebrospinal fluid; CNS = central nervous system; HCV = hepatitis C virus; VACS = Veterans Aging Cohort Study.


<sup>a</sup>Based on independent sample t-tests and Chi-Square or Fisher's exact tests.

<sup>b</sup>Major genetic ancestry refers to having a genetic ancestry proportion of  $\geq .5$  for a given region (Africa, Europe, America, East Asia, Central Asia, and Oceania), with participants that have all frequencies  $< .5$  being assigned to "no single major ancestry"; Mexican: *n* = 62; Puerto Rican: *n* = 33.

<sup>c</sup>Among those on ART (antiretroviral therapy).

models adjusting for these variables, continued to show a significant effect of country of origin on global NCI (Puerto Ricans-Mexicans: OR = 2.40; 95% CI = 1.11–5.29; *p* = .03),

as well as learning and memory/delayed recall (Puerto Ricans-Mexicans: Learning, OR = 3.09; 95% CI = 1.44–6.81; *p* < .01; Memory, OR = 2.22; 95% CI = 1.06–4.71; *p* = .03).



**Fig. 2.** Rates of global and domain neurocognitive impairment by Latino group (Mexican and Puerto Rican origin/descent). \* $p < .01$  \*\* $p < .001$ .

Path analyses on global NCI, showed that comorbidity status was the only significant partial mediator of differences associated with country of origin (indirect effect = 0.07; 95% CI = 0.02–0.15;  $p = .02$ ). Multivariable regression analyses on global NCI within the subset of participants on ART, adjusting for comorbidity status and detectable CSF RNA, also showed a significant effect of group (Puerto Ricans-Mexicans: OR = 3.10; 95% CI = 1.08–9.92;  $p = .04$ ).

Regarding predictors of NCI among Puerto Ricans, being on ART and increased comorbidity status were significantly associated with NCI in univariable analyses. In a multivariable model, being on ART (OR = 9.71; 95% CI = 1.78–71.09;  $p < .001$ ) and comorbidity status (confounding-incident: OR = 13.39; 95% CI = 1.86–290.95;  $p = .02$ ; contributing-incident: OR = 5.89; 95% CI = 1.35–35.34;  $p = .02$ ) were both significant independent predictors of NCI.

Among Mexicans, univariable analyses showed that lower nadir CD4 and increased comorbidity status were significant predictors of global NCI. In multivariable analyses, both lower nadir CD4 (OR/100 units = 0.75; 95% CI = 0.53–1.00;  $p = .05$ ), and comorbidity status were marginally associated with global NCI (confounding-incident: OR = 7.01; 95% CI = 1.00–141.7;  $p = .05$ ), but with a high ORs.

### Association of Genetic Ancestry and Self-Reported Ethnicity or Country of Origin/Descent to NCI

Stratified analyses by ethnic group showed there were no significant differences between the samples with and without genetic data on demographic variables (age, gender, education). Among participants with genetic ancestry data, a multivariable model in the overall sample on global NCI showed a significant effect of ethnicity (Latino-White: OR = 1.65; 95% CI = 1.02–2.66;  $p = .04$ ) and no significant effect of genetic ancestry ( $p = .45$ ). Separate univariable analyses by ethnic group showed proportions of European, African, or

American Indian genetic ancestry were not significantly associated with NCI in Latinos ( $ps = .59, .68, .46$ , respectively) or in Whites ( $ps = .47, .12, .85$ , respectively).

A multivariable analysis on global NCI among our subset of Mexicans and Puerto Ricans with available genetic ancestry data yielded a significant effect of country of origin/descent (Puerto Ricans-Mexicans: OR = 3.08; 95% CI = 1.23–8.27;  $p = .02$ ) and a significant overall effect of major genetic ancestry ( $p = .02$ ). Specifically, the group with No single major ancestry had marginally higher rates of NCI than the European major ancestry group (OR = 2.65; 95% CI = 0.89–8.70;  $p = .08$ ), who, in turn, had marginally higher rates of NCI than those with another major ancestry (African, American, or East Asian combined; OR = 3.83; 95% CI = 0.97–19.72;  $p = .06$ ); with significant differences between this latter group and the No single major ancestry group ( $p < .01$ ). Separate univariable analyses by country of origin group showed proportions of European, African, or American Indian genetic ancestry were not significantly associated with NCI in Puerto Ricans ( $ps = .64, .67, .24$ , respectively) or Mexicans ( $ps = .70, .71, .12$ , respectively).

### Secondary Analyses

#### *Subgroup of Whites matched to Latinos on demographics*


To further control for differences in age, education and gender between ethnic groups, in exploratory analyses, we selected a subset of Whites ( $n = 194$ ) who were one-to-one matched with Latinos on age ( $\pm 5$  years), education ( $\pm 2$  years), and gender (all but one participant were matched on gender), and re-ran all core analyses. Latinos had higher rates of global NCI than the matched White group, as well as higher rates of NCI in the domains of learning, delayed recall and working memory. Results of analyses adjusting for significant covariates in this matched subset of participants (nadir CD4 and alcohol use disorder) on global and domain NCI yielded comparable findings to those of the overall cohort (i.e., while  $p$  values only approached significance in some cases, OR were similar or higher).

We also compared Mexicans and Puerto Ricans to their respective matched group of Whites on global and domain T-scores. Figure 3 shows Cohen's  $d$  effect sizes for these comparisons. For Puerto Ricans, there were medium-to-large effect sizes for global neurocognition, which appeared to be driven primarily by large effect sizes in learning and memory, and small-to-medium effects on verbal fluency, speed of information processing, executive function and working memory. For Mexicans, there were small-to-medium effect sizes on executive function and working memory.

#### *Norms with ethnicity corrections*

For both components of the Processing Speed Index of the WAIS-III (Digit Symbol Coding and Symbol Search subtests; Heaton et al., 2002) we had normative data for English-speaking





**Fig. 3.** Effect sizes on global and domain neurocognitive function (T-scores) by subgroup of Latinos (i.e., Mexicans,  $n=79$ ; and Puerto Ricans,  $n=60$ ) compared with their respective matched subgroups of non-Latino Whites (matched on age, gender, and education). Positive effect sizes indicate worse performance among Latino subgroups. Effect sizes: 0.20 = small; 0.50 = medium; 0.80 = large (Cohen, 1988).

Latinos living in the United States. To explore whether group differences might be attributed to a lack of appropriate norms, we ran a linear regression model on the average adjusted T-scores of the subtests comprising the Processing Speed Index (with Whites as the reference group). Results again showed a significant effect of group,  $F(2,793)=2.77$ ,  $p=.04$ , and indicated that these group differences were primarily driven by Puerto Ricans performing significantly lower than Whites ( $Estimate=-2.96$ ;  $SE=1.39$ ;  $p=.03$ ).

### Impact of study site

Because of the distribution of Latinos in the United States and location of the study sites, ethnicity and country or origin/descent were highly confounded with study site (Tables 1 and 2). We explored the potential effect of study site by running a multivariable model on NCI with predictors being ethnicity and study site among participants tested at sites with reasonable numbers of Whites and Latinos (i.e., San Diego, New York, Galveston, and Seattle;  $n=658$ ). The model showed significant independent effects of ethnicity ( $p=.015$ ) and site ( $p<.001$ ). Given that none of the sites had adequate numbers of both Mexicans and Puerto Ricans, we were unable to run comparable analyses among Latino subgroups. Instead, we ran separate chi-square tests on NCI, comparing Mexicans and Whites in San Diego (Mexicans:  $n=36$ ;  $NCI=47\%$ ; Whites:  $n=165$ ;  $NCI=32\%$ ;  $p=.09$ ) and Puerto Ricans and Whites in New York (Puerto Ricans:  $n=47$ ;  $NCI=79\%$ ; Whites:  $n=61$ ;  $NCI=54\%$ ;  $p=.007$ ).

## DISCUSSION

Present findings support that HIV+ Latinos receiving care at academic medical centers in the United States are at increased

risk for concurrent NCI compared with their White counterparts. This study also highlights that there are important differences among Latino subgroups, and one relevant way in which Latinos differ is their country of origin/descent. In the present study of HIV+ persons, Latinos of Puerto Rican origin/descent showed significantly increased rates of NCI compared with those of Mexican origin/descent.

Our results are consistent with findings of two prior studies (Rivera-Mindt et al., 2008; Wojna et al., 2006), which indicated increased rates of global or domain NCI in HIV+ Latinos of primarily Caribbean descent and with advanced HIV disease. Somewhat inconsistent with present findings, another study (Rivera-Mindt et al., 2014) found rates of NCI were not significantly different among HIV+ Latinos and Whites younger than 50 years of age. Non-significant findings in this prior study might have been partly due to the relatively small sample sizes (Whites:  $n=20$ ; Latinos:  $n=68$ ), particularly given that there were small to medium effect sizes (as measured by Cohen's  $d$ ) on the domains of verbal fluency, attention/working memory, learning, and fine motor skills, and a medium effect size on executive function. In addition, 20% of the present sample was 50 years or older, and this prior study (Rivera-Mindt et al., 2014) found significantly increased rates of NCI among older (50+ years old) HIV+ Latinos than older HIV+ Whites. This suggests that the differences observed in the present study, might be exacerbated in older cohorts.

Ethnic differences on NCI were evident across the domains of executive function, working memory, speed of information processing, learning, and recall, with no significant differences on verbal fluency and fine motor skills. Most of the neurocognitive domains in which there were significant ethnic differences (with the exception of speed of information processing) are among the most commonly impacted by HIV in the combination ART era (Heaton et al., 2011), and are linked with problems in everyday functioning in HIV+ persons (Heaton et al., 2004). This underscores the potential significance of our findings for the daily lives of HIV+ Latinos.

In addition to having different rates of NCI, ethnic groups differed in several other factors, including HIV-disease and non HIV-disease characteristics. Consistent with prior findings (Swindells et al., 2002), Latinos in our study showed worse HIV disease burden than Whites, both in terms of historical and current HIV disease characteristics. While lower nadir CD4 among Latinos significantly accounted for some of the ethnic differences in NCI, differences in NCI were still significant in adjusted models. This suggests that other psychosocial (e.g., acculturation, socioeconomic status, healthcare access, health behaviors) and biomedical factors of particular relevance among Latinos living with HIV need to be considered.

As HIV+ persons are living longer, HIV has become a complex disease with multiple causes of morbidity. While there were no significant ethnic group differences on comorbidity status, there was a trend for Latinos to have higher VACS Index scores. Additionally, comorbidity status

was among the most important predictors of NCI, both in Whites and Latinos, indicating that there might be certain specific comorbidities worth investigating in more detail. One such factor might be metabolic syndrome (Grundy et al., 2004), which is highest among Latinos (Aguilar, Bhuket, Torres, Liu, & Wong, 2015; Ford, Giles, & Dietz, 2002; Heiss et al., 2014) and has been linked to HIV-associated NCI (Fabbiani et al., 2013; McCutchan et al., 2012).

We did not have an HIV-uninfected comparison group, and most of the tests did not have norms available that adjusted for Latino ethnicity. Thus, we cannot rule out that some of the ethnic differences observed in this study of HIV+ persons might be observed in the non HIV-infected population. In the present study, however, group comparisons on neurocognitive tests that do have normative adjustments for Latino ethnicity yielded comparable findings to those in the overall battery of tests. This, together with the findings that significant ethnic differences were observed on neurocognitive domains that are most commonly impacted by HIV, that nadir CD4 partially mediated ethnic differences in NCI, and that NCI within our Latino group(s) was associated with multiple indicators of HIV disease severity, supports the notion that these group differences are, at least partly, HIV-related. Consistently, a large longitudinal study using data on the CHARTER cohort (Heaton et al., 2015), found that Latino ethnicity was an independent predictor of HIV-associated neurocognitive decline.

In line with prior studies among HIV+ persons (Sheehan et al., 2015), we found Puerto Ricans fared worse than Mexicans on HIV-disease burden, comorbidity status and injection drug use. They were also significantly more likely to show global NCI than Mexicans, with these differences being primarily driven by differential rates of NCI in the domains of learning and delayed recall. Findings showing worse NCI among Puerto Ricans than Mexicans in this study are also consistent with other reports indicating worse health outcomes (Velasco-Mondragon, Jimenez, Palladino-Davis, Davis, & Escamilla-Cejudo, 2016), and worse neurocognitive performance (González et al., 2015) among Puerto Ricans than other Latino subgroups living in the United States.

While there might be behavioral factors related to health-care use and adherence that might impact several health outcomes, there might also be some disease-specific genetic/ biomedical vulnerabilities that help explain these disparities. In the present study, there is some indication that both biomedical and behavioral factors might be playing a role. Comorbidity status partially mediated differences in NCI among Latino subgroups, and was a key predictor of NCI in all groups, again highlighting the importance of investigating comorbidities in future studies. Of interest, Puerto Ricans are the subgroup of Latinos with the highest rates of metabolic syndrome (Heiss et al., 2014), and, as mentioned above, metabolic syndrome has been linked to HIV-associated NCI.

Additionally, among Puerto Ricans, being on ART conferred an increased risk for NCI. Being on ART is at least partly a marker of worse disease status, such that those who present to clinic with worse HIV-disease status are more

likely to be prescribed ART. The fact that this factor was associated with NCI among Puerto Ricans, but not other groups in multivariable analyses, might be related to worse HIV-adherence among Puerto Ricans (Robbins et al., 2012). Consistent with this, among those on ART, Puerto Ricans were more likely to have detectable CSF HIV RNA. Regarding Mexicans, nadir CD4 also was an independent predictor of NCI, which is consistent with prior studies of HIV-associated NCI (Ellis et al., 2011) and underscores the importance of early HIV treatment and testing in this group.

In light of prior evidence showing greater NCI risk among HIV+ females than males (Kabuba, Menon, Franklin, Heaton, & Hestad, 2016), it is worth noting that there was a higher percentage of females among Puerto Ricans than Mexicans. However, findings on the association between gender and HIV-associated NCI are inconsistent (Faílde-Garrido, Rodríguez Alvarez, & Simón-López, 2008; Heaton et al., 2011, 2015), and given our methodological approach, it is unlikely that NCI differences between Latino subgroups were accounted for by differential gender composition. Because of the small number of females (Mexican:  $n = 10$ ; Puerto Rican:  $n = 23$ ), we were unable to investigate gender differences in great detail, and it would be important for future studies to do so.

A unique aspect of the present study was the use of genomic data to characterize the genetic ancestry of self-reported ethnic groups. Present findings did not find a significant impact of genetic ancestry on NCI by ethnic or country of origin groups. Yet, genetic ancestry was a significant independent contributor to NCI among Latino subgroups in addition to country of origin/descent. Future larger studies investigating genetic markers of diseases that might contribute to HIV-associated NCI and be more common among subgroups of Latinos, might be key for elucidating the types of exposures that might contribute to ethnic differences in NCI.

Another important aspect of the present study concerns geographic differences in the distribution of the study groups, and the impact of geographic site on NCI. Latino/a ethnicity and country of origin both traveled with study site, which is consistent with U.S. Census Bureau data on the geographic distribution of Latinos in the United States (Ennis et al., 2011). In the present study, study site did not appear to fully explain differences in rates of NCI among Latinos and Whites. Furthermore, we found a 15% increase in rates of NCI in Mexicans compared with Whites in San Diego and a 25% increase in rates of NCI in Puerto Ricans compared with Whites in New York.

While the difference in NCI between Mexicans and Whites was not statistically different, at least partly because of the smaller sample sizes, overall these findings indicate that geographic differences do not fully explain ethnic/cultural disparities in NCI. Of interest, however, there was a significant effect of study site. While this was not the focus of present analyses, understanding the factors driving these site differences might shed light onto other social and environmental factors impacting NCI among HIV-infected persons in the United States.

There are several limitations to our study. We did not have data on several culturally relevant factors, such as language use history, bilingualism, socioeconomic status, country of birth, and acculturation. Future studies incorporating these types of assessments might shed light on factors underlying differences in NCI among subgroups of Latinos. The present study included English-speaking Latinos living in the United States, and thus caution is warranted in generalizing findings to Spanish-speaking and other subgroups of HIV+ Latinos.

In summary, HIV+ Latinos in the United States appear to be at higher risk for NCI compared with HIV+ Whites, with HIV+ Puerto Ricans being at particularly increased risk. In examining these ethnic differences, it is critical to recognize the heterogeneity within the Latino population, as well as the complex interplay of disease and non-disease factors, as well as genetic and environmental risks. Present findings underscore the need for the development of larger studies particularly focused on examining ethnic differences in NCI among HIV+ persons, that include different subgroups of Latinos and systematically consider and incorporate relevant socio-cultural, biomedical, and genetic factors.

## ACKNOWLEDGMENTS

The CNS HIV Anti-Retroviral Therapy Effects Research (CHARTER) is supported by awards N01 MH22005, HHSN271201000036C, and HHSN271201000030C from the National Institutes of Health. M.J. Marquine: K23MH105297. The CNS HIV Anti-Retroviral Therapy Effects Research (CHARTER) group is affiliated with the Johns Hopkins University, Mount Sinai School of Medicine, University of California, San Diego, University of Texas, Galveston, University of Washington, Seattle, Washington University, St. Louis and is headquartered at the University of California, San Diego and includes: Director: Igor Grant, M.D.; Co-Directors: Scott L. Letendre, M.D., Ronald J. Ellis, M.D., Ph.D., Thomas D. Marcotte, Ph.D.; Center Manager: Donald Franklin, Jr.; Neuromedical Component: Ronald J. Ellis, M.D., Ph.D. (P.I.), J. Allen McCutchan, M.D., Terry Alexander, R.N.; Laboratory, Pharmacology and Immunology Component: Scott Letendre, M.D. (P.I.), Edmund Capparelli, Pharm.D.; Neurobehavioral Component: Robert K. Heaton, Ph.D. (P.I.), J. Hampton Atkinson, M.D., Steven Paul Woods, Psy.D., Matthew Dawson; Virology Component: Davey M. Smith, M.D. (P.I.); Imaging Component: Christine Fennema-Notestine, Ph.D. (P.I.), Michael J. Taylor, Ph.D., Rebecca Theilmann, Ph.D.; Data Management Component: Anthony C. Gamst, Ph.D. (P.I.), Clint Cushman; Statistics Component: Ian Abramson, Ph.D. (P.I.), Florin Vaida, Ph.D.; Protocol Coordinating Component: Thomas D. Marcotte, Ph.D. (P.I.), Jennifer Marquie-Beck, M.P.H.; Johns Hopkins University Site: Justin McArthur (P.I.), Vincent Rogalski; Mount Sinai School of Medicine Site: Susan Morgello, M.D. (Co-P.I.) and David Simpson, M.D. (Co-P.I.), Letty Mintz, N.P.; University of California, San Diego Site: J. Allen McCutchan, M.D. (P.I.), Will Toperoff, N.P.; University of Washington, Seattle Site: Ann Collier, M.D. (Co-P.I.) and Christina Marra, M.D. (Co-P.I.), Trudy Jones, M.N., A.R.N.P.; University of Texas, Galveston Site: Benjamin Gelman, M.D., Ph.D. (P.I.), Eleanor Head, R.N., B.S.N.; and Washington University, St. Louis Site: David Clifford, M.D. (P.I.), Muhammad Al-Lozi, M.D., Mengesha Teshome, M.D. The views expressed in this article are those of the authors and do

not reflect the official policy or position of the United States Government. There are no conflicts of interest to report.

## REFERENCES

- Aguilar, M., Bhuket, T., Torres, S., Liu, B., & Wong, R.J. (2015). Prevalence of the metabolic syndrome in the United States, 2003-2012. *Journal of the American Medical Association*, 313(19), 1973-1974. doi: 10.1001/jama.2015.4260
- Alexander, D.H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, 19(9), 1655-1664. doi: 10.1101/gr.094052.109
- American Psychiatric Association. (1994). *Diagnostic and Statistical Manual of Mental Disorders, 4th edition*. Washington; DC: American Psychiatric Association.
- Antinori, A., Arendt, G., Becker, J.T., Brew, B.J., Byrd, D.A., Cherner, M., ... Wojna, V.E. (2007). Updated research nosology for HIV-associated neurocognitive disorders. *Neurology*, 69(18), 1789-1799. doi:10.1212/01.WNL.0000287431.88658.8b.
- Beck, A.T., Steer, R.A., & Brown, G.K. (1996). *Beck Depression Inventory - Second Edition Manual*. San Antonio: The Psychological Corporation.
- Bryc, K., Durand, E.Y., Macpherson, J.M., Reich, D., & Mountain, J.L. (2015). The genetic ancestry of African Americans, Latinos, and European Americans across the United States. *American Journal of Human Genetics*, 96(1), 37-53. doi: 10.1016/j.ajhg.2014.11.010
- Cann, H.M., de Toma, C., Cazes, L., Legrand, M.F., Morel, V., Piouffre, L., ... Cavalli-Sforza, L.L. (2002). A human genome diversity cell line panel. *Science*, 296(5566), 261-262.
- Carey, C.L., Woods, S.P., Gonzalez, R., Conover, E., Marcotte, T. D., & Grant, I., . . . HNRC Group. (2004). Predictive validity of global deficit scores in detecting neuropsychological impairment in HIV infection. *Journal of Clinical and Experimental Neuropsychology*, 26(3), 307-319. doi:10.1080/13803390490510031.
- Centers for Disease Control and Prevention. (2015). HIV surveillance report: Diagnoses of HIV infection in the United States and dependent areas, 2014. 26. <http://www.cdc.gov/hiv/library/reports/surveillance/>
- Chen, N.E., Gallant, J.E., & Page, K.R. (2012). A systematic review of HIV/AIDS survival and delayed diagnosis among Hispanics in the United States. *Journal of Immigrant and Minority Health*, 14(1), 65-81. doi: 10.1007/S10903-011-9497-Y
- Cohen, J. (1988). *Statistical power analyses for the behavioral sciences: Second edition*. New York: Routledge.
- Conomos, M.P., Laurie, C.A., Stilp, A.M., Gogarten, S.M., McHugh, C.P., Nelson, S.C., ... Laurie, C.C. (2016). Genetic diversity and association studies in US Hispanic/Latino populations: Applications in the Hispanic Community Health Study/ Study of Latinos. *American Journal of Human Genetics*, 98(1), 165-184. doi: 10.1016/j.ajhg.2015.12.001
- Dennis, A.M., Napravnik, S., Sena, A.C., & Eron, J.J. (2011). Late entry to HIV care among Latinos compared with non-Latinos in a southeastern US cohort. *Clinical Infectious Diseases*, 53(5), 480-487. doi: 10.1093/Cid/Cir434
- Durvasula, R.S., Miller, E.N., Myers, H.F., & Wyatt, G.E. (2001). Predictors of neuropsychological performance in HIV positive women. *Journal of Clinical and Experimental Neuropsychology*, 23(2), 149-163. doi: 10.1076/Jcen.23.2.149.1211

- Ellis, R.J., Badiee, J., Vaida, F., Letendre, S., Heaton, R.K., Clifford, D., . . . CHARTER Group. (2011). CD4 nadir is a predictor of HIV neurocognitive impairment in the era of combination antiretroviral therapy. *AIDS*, 25(14), 1747–1751. doi: 10.1097/Qad.0b013e32834a40cd
- Ennis, S.R., Rios-Vargas, M., & Albert, N.G. (2011). *The Hispanic population: 2010*. Suitland, MD: U.S. Census Bureau.
- Fabbiani, M., Ciccirelli, N., Tana, M., Farina, S., Balonero, E., Di Cristo, V., . . . Di Giambenedetto, S. (2013). Cardiovascular risk factors and carotid intima-media thickness are associated with lower cognitive performance in HIV-infected patients. *HIV Medicine*, 14(3), 136–144. doi: 10.1111/J.1468-1293.2012.01044
- Fañde-Garrido, J.M., Rodríguez Alvarez, M., & Simón-López, M.A. (2008). Neuropsychological impairment and gender differences in HIV-1 infection. *Psychiatry and Clinical Neurosciences*, 62, 494–502.
- Ford, E.S., Giles, W.H., & Dietz, W.H. (2002). Prevalence of the metabolic syndrome among US adults - Findings from the Third National Health and Nutrition Examination Survey. *Journal of the American Medical Association*, 287(3), 356–359. doi: 10.1001/Jama.287.3.356
- Gardner, E.M., McLees, M.P., Steiner, J.F., del Rio, C., & Burman, W.J. (2011). The spectrum of engagement in HIV care and its relevance to test-and-treat strategies for prevention of HIV infection. *Clinical Infectious Diseases*, 52(6), 793–800. doi: 10.1093/Cid/Ciq243
- González, H.M., Tarraf, W., Gouskova, N., Gallo, L., Penedo, F.J., Davis, S.M., . . . Mosley, T.H. (2015). Neurocognitive function among middle-aged and older Hispanic/Latinos: Results from the Hispanic Community Health Study/Study of Latinos. *Archives of Clinical Neuropsychology*, 30, 68–77.
- Gorman, A.A., Foley, J.M., Ettenhofer, M.L., Hinkin, C.H., & van Gorp, W.G. (2009). Functional consequences of HIV-associated neuropsychological impairment. *Neuropsychology Review*, 19(2), 186–203. doi: 10.1007/s11065-009-9095-0
- Gray, K.M., Valverde, E.E., Tang, T., Siddiqi, A.E.A., & Hall, H.I. (2015). Diagnoses and prevalence of HIV infection among Hispanics or Latinos - United States, 2008–2013. *MMWR Morbidity and Mortality Weekly Report*, 64(39), 1097–1103.
- Grundy, S.M., Brewer, H.B., Cleeman, J.I., Smith, S.C., & Lenfant, C., National Heart, Lung, and Blood Institute; American Heart Association. (2004). Definition of metabolic syndrome - Report of the National Heart, Lung, and Blood Institute/American Heart Association conference on scientific issues related to definition. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 24(2), E13–E18. doi: 10.1161/01.Atv.0000111245.75752.C6
- Guarnaccia, P.J., Martinez Pincay, I., Alegria, M., Shrout, P.E., Lewis-Fernandez, R., & Canino, G. (2007). Assessing diversity among Latinos: Results from the NLAAS. *Hispanic Journal of Behavioral Sciences*, 29(4), 510–534.
- Heaton, R.K., Clifford, D.B., Franklin, D.R., Woods, S.P., Ake, C., Vaida, F., . . . CHARTER Group. (2010). HIV-associated neurocognitive disorders persist in the era of potent antiretroviral therapy CHARTER Study. *Neurology*, 75(23), 2087–2096.
- Heaton, R.K., Franklin, D.R., Ellis, R.J., McCutchan, J.A., Letendre, S.L., LeBlanc, S., . . . the CHARTER; HNRC Group. (2011). HIV-associated neurocognitive disorders before and during the era of combination antiretroviral therapy: Differences in rates, nature, and predictors. *Journal of Neurovirology*, 17(1), 3–16. doi: 10.1007/S13365-010-0006-1
- Heaton, R.K., Franklin, D.R. Jr., Deutsch, R., Letendre, S., Ellis, R.J., Casaletto, K., . . . CHARTER Group. (2015). Neurocognitive change in the era of HIV combination antiretroviral therapy: The longitudinal CHARTER study. *Clinical Infectious Diseases*, 60, 473–480. doi: 10.1093/cid/ciu862
- Heaton, R.K., Marcotte, T.D., Mindt, M.R., Sadek, J., Moore, D.J., Bentley, H., . . . HNRC Group. (2004). The impact of HIV-associated neuropsychological impairment on everyday functioning. *Journal of the International Neuropsychological Society*, 10(3), 317–331. doi: 10.1017/S1355617704102130
- Heaton, R.K., Taylor, M.J., & Manly, J.J. (2002). *Demographic effects and use of demographically corrected norms with the WAIS-III and WMS-III*. San Diego: Academic Press.
- Heiss, G., Snyder, M.L., Teng, Y., Schneiderman, N., Llabre, M.M., Cowie, C., . . . Aviles-Santa, L. (2014). Prevalence of metabolic syndrome among Hispanics/Latinos of diverse background: The Hispanic Community Health Study/Study of Latinos. *Diabetes Care*, 37(8), 2391–2399. doi: 10.2337/dc13-2505
- Imai, K., Keele, L., & Tingley, D. (2010). A general approach to causal mediation analysis. *Psychological Methods*, 15(4), 309–334. doi: 10.1037/a0020761
- International HapMap 3 Consortium, Altshuler, D.M., Gibbs, R.A., Peltonen, L., Altshuler, D.M., Gibbs, R.A., & McEwen, J.E. (2010). Integrating common and rare genetic variation in diverse human populations. *Nature*, 467(7311), 52–58. doi: 10.1038/nature09298
- Justice, A.C., Freiberg, M.S., Tracy, R., Kuller, L., Tate, J.P., Goetz, M.B., . . . VACS Project Team. (2012). Does an index composed of clinical data reflect effects of inflammation, coagulation, and monocyte activation on mortality among those aging With HIV? *Clinical Infectious Diseases*, 54(7), 984–994. doi: 10.1093/Cid/Cir989
- Justice, A.C., Modur, S. P., Tate, J.P., Althoff, K.N., Jacobson, L.P., Gebo, K.A., . . . Gange, S.J. (2013). Predictive accuracy of the Veterans Aging Cohort Study Index for mortality with HIV infection: A North American cross cohort analysis. *Journal of Acquired Immune Deficiency Syndromes (1999)*, 62(2), 149–163. doi: 10.1097/QAI.0b013e31827df36c
- Kabuba, N., Menon, J.A., Frnaklin, D.R., Heaton, R.K., & Hestad, K.A. (2016). HIV- and AIDS-associated neurocognitive functioning in Zambia – a perspective based on differences between the genders. *Neuropsychiatric Disease and Treatment*, 12, 2021–2028.
- Lee, S., & Held, M.L. (2015). Variation in mental health service use among US Latinos by place of origin and service provider type. *Psychiatric Services*, 66(1), 56–64. doi: 10.1176/appi.ps.201300533
- Letendre, S. (2011). Central nervous system complications in HIV disease: HIV-associated neurocognitive disorder. *Topics in Antiviral Medicine*, 19(4), 137–142.
- Letendre, S., Marquie-Beck, J., Capparelli, E., Best, B., Clifford, D., Collier, A.C., . . . Grp, C. (2008). Validation of the CNS penetration-effectiveness rank for quantifying antiretroviral penetration into the central nervous system. *Archives of Neurology*, 65(1), 65–70. doi: 10.1001/Archneuro.2007.31
- Libiger, O., & Schork, N.J. (2013). A method for inferring an individual's genetic ancestry and degree of admixture associated with six major continental populations. *Frontiers in Genetics*, 3, 322. doi: 10.3389/fgene.2012.00322
- Llorrente, A.M. (2008). *Principles of neuropsychological assessment with Hispanics*. New York; NY: Springer.

- McCutchan, J.A., Marquie-Beck, J.A., FitzSimons, C.A., Letendre, S.L., Ellis, R.J., & Heaton, R.K., . . . CHARTER Group. (2012). Role of obesity, metabolic variables, and diabetes in HIV-associated neurocognitive disorder. *Neurology*, 78(7), 485–492. doi: 10.1212/Wnl.0b013e3182478d64
- Nelson, M.R., Bryc, K., King, K.S., Indap, A., Boyko, A.R., Novembre, J., . . . Lai, E.H. (2008). The Population Reference Sample, POPRES: A resource for population, disease, and pharmacological genetics research. *American Journal of Human Genetics*, 83(3), 347–358. doi: 10.1016/j.ajhg.2008.08.005
- Office of Management and Budget. (1997). Revisions to the standards for the classification of federal data on race and ethnicity. Retrieved from [https://http://www.whitehouse.gov/omb/fedreg\\_1997standards](https://http://www.whitehouse.gov/omb/fedreg_1997standards)
- Preacher, K.J., & Hayes, A.F. (2004). SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behavior Research Methods Instruments & Computers*, 36(4), 717–731. doi: 10.3758/Bf03206553
- Rivera-Mindt, M., Byrd, D., Ryan, E.L., Robbins, R., Monzones, J., Arentoft, A., . . . Henniger, D. (2008). Characterization and sociocultural predictors of neuropsychological test performance in HIV+ Hispanic individuals. *Cultural Diversity & Ethnic Minority Psychology*, 14(4), 315–325. doi: 10.1037/A0012615
- Rivera-Mindt, M., Miranda, C., Arentoft, A., Byrd, D., Monzones, J., Fuentes, A., . . . Morgello, S. (2014). Aging and HIV/AIDS: Neurocognitive implications for older HIV-positive Latina/o adults. *Behavioral Medicine*, 40(3), 116–123. doi: 10.1080/08964289.2014.914464
- Robbins, R.N., D'Aquila, E., Morgello, S., Byrd, D., Remien, R.H., & Mindt, M.R. (2012). Cultural influences on antiretroviral therapy adherence among HIV-infected Puerto Ricans. *Journal of the Association of Nurses in AIDS Care*, 23(6), 531–538. doi: 10.1016/j.jana.2011.12.006
- Shapiro, M.F., Morton, S.C., McCaffrey, D.F., Senterfitt, J.W., Fleishman, J.A., Perlman, J.F., . . . Consortium, H. (1999). Variations in the care of HIV-infected adults in the United States - Results from the HIV Cost and Services Utilization Study. *Journal of the American Medical Association*, 281(24), 2305–2315. doi: 10.1001/Jama.281.24.2305
- Sheehan, D.M., Trepka, M.J., & Dillon, F.R. (2015). Latinos in the United States on the HIV/AIDS care continuum by birth country/region: A systematic review of the literature. *International Journal of STD & AIDS*, 26(1), 1–12. doi: 10.1177/0956462414532242
- Swindells, S., Cobos, D.G., Lee, N., Lien, E.A., Fitzgerald, A.P., Pauls, J.S., . . . Anderson, J.R. (2002). Racial/ethnic differences in CD4 T cell count and viral load at presentation for medical care and in follow-up after HIV-1 infection. *AIDS*, 16(13), 1832–1834. doi: 10.1097/00002030-200209060-00020
- Thames, A.D., Arentoft, A., Rivera-Mindt, M., & Hinkin, C.H. (2013). Functional disability in medication management and driving among individuals with HIV: A 1-year follow-up study. *Journal of Clinical and Experimental Neuropsychology*, 35(1), 49–58. doi: 10.1080/13803395.2012.747596
- Turner, B.J., Cunningham, W.E., Duan, N., Andersen, R.M., Shapiro, M.F., Bozzette, S.A., . . . Zierler, S. (2000). Delayed medical care after diagnosis in a US national probability sample of persons infected with human immunodeficiency virus. *Archives of Internal Medicine*, 160(17), 2614–2622. doi: 10.1001/archint.160.17.2614
- U.S. Census Bureau. Population Estimates 2015. Retrieved from <https://www.census.gov/quickfacts/table/PST045215/00>
- Velasco-Mondragon, E., Jimenez, A., Palladino-Davis, A.G., Davis, D., & Escamilla-Cejudo, J.A. (2016). Hispanic health in the USA: A scoping review of the literature. *Public Health Reviews*, 37, 31. doi: 10.1186/s40985-016-0043-2
- Wechsler, D. (1997). *Wechsler Adult Intelligence Scale - Third Edition*. San Antonio, TX: The Psychological Corporation.
- Wittchen, H.U., Robins, L.N., Cottler, L.B., Sartorius, N., Burke, J.D., & Regier, D. (1991). Cross-cultural feasibility, reliability and sources of variance of the Composite International Diagnostic Interview (CIDI). The Multicentre WHO/ADAMHA Field Trials. *The British Journal of Psychiatry*, 159(645–653), 658.
- Wojna, V., Skolasky, R.L., Hechavarría, R., Mayo, R., Selnes, O., McArthur, J.C., . . . Nath, A. (2006). Prevalence of human immunodeficiency virus-associated cognitive impairment in a group of Hispanic women at risk for neurological impairment. *Journal of Neurovirology*, 12(5), 356–364. doi: 10.1080/13550280600964576
- Xing, J., Watkins, W.S., Witherspoon, D.J., Zhang, Y., Guthery, S.L., Thara, R., . . . Jorde, L.B. (2009). Fine-scaled human genetic structure revealed by SNP microarrays. *Genome Research*, 19(5), 815–825. doi: 10.1101/gr.085589.108