

Photoionization modeling of the Galactic planetary nebulae Abell 39 and NGC 7027

Ashkbiz Danehkar¹, David J. Frew¹, Quentin A. Parker^{1,2} and Orsola De Marco¹

¹Department of Physics and Astronomy, Macquarie University, Sydney, NSW 2109, Australia

²Australian Astronomical Observatory, PO Box 296, Epping, NSW 1710, Australia
email: ashkbiz.danehkar@mq.edu.au; david.frew@mq.edu.au

Abstract. We estimate distances to the spherical planetary nebula Abell 39 and the bipolar planetary nebula NGC 7027 by interpolating from a wide grid of photoionization models using the 3-D code, MOCASSIN. We find preliminary distances of 1.5 kpc and 0.9 kpc respectively, with uncertainties of about 30%.

Keywords. planetary nebulae: individual (Abell 39, NGC 7027)

1. Introduction

Accurate distances to planetary nebulae (PNe) are crucial in unraveling the connection between the physical properties of the nebulae with those of their central stars (CSs). Reliable distances facilitate the accurate estimation of fundamental parameters such as the CS mass and luminosity, and the nebular mass and age. We have begun a program of using photoionization modeling to refine the distances to a sample of nearby PNe (see also Danehkar *et al.* 2011). In this work, we study two very different PNe as a proof of concept: Abell 39, a simple spherical shell with no microstructures, and NGC 7027, a well-known, young, luminous bipolar PN with a massive molecular envelope. Using the 3-D photoionization code (MOCASSIN; Ercolano *et al.* 2003), our ultimate aim is to constrain the distance to individual PNe, utilizing the physical PN radius we have calculated and the angular size.

2. Analysis

Our procedure allows us refine the nebular parameters iteratively to provide the best match against the observations, though owing to degeneracies in these parameters, there may be more than one unique solution. Since a black body is not a perfect model for the CS continuum flux, we used various NLTE model atmosphere fluxes from the grid of Rauch (2003). For the initial model inputs for A 39, we adopt the line intensities and abundances from Jacoby *et al.* (2001), and derived the nebular T_e from the [OIII] lines. Because the density of A 39 is very low, we initially determined N_e from the PN diameter and integrated $H\beta$ flux. Our preliminary distance estimate is $D = 1.5$ kpc, and the CS parameters are $T_{\text{eff}} = 160$ kK and $L/L_{\odot} = 1800$, which disagrees with the estimate of $T_{\text{eff}} = 117$ kK from Ziegler (2011, these proceedings). The higher temperature is needed to explain the observed He II and [Ne V] line intensities in the nebular shell.

For NGC 7027, a starting value of $T_e = 11$ kK is adopted, the line intensities and abundances being taken from Bernard Salas *et al.* (2001) and Zhang *et al.* (2005), and refined as necessary. In addition, a bipolar morphology with a mean density ($n_{\text{H}} = 55,000 \text{ cm}^{-3}$; Fig. 1) is needed to model NGC 7027, since a simple assumption of spherical

Table 1. Best-fit parameters (left) and observations versus model outputs (right).

Parameter	NGC 7027		Ion	Line $\lambda(\text{\AA})$	A 39		NGC 7027	
	A 39	NGC 7027			Obs.	Mod.	Obs.	Mod.
T_{eff} (K)	160 000	180 000	[Ne v]	3426	9.8	8.0	154	139
L_{cs} (L_{\odot})	1800	7500	[O II]	3727	40	51	21	21
M_{*} (M_{\odot})	0.59	0.69	[Ne III]	3869	104	92	126	143
R_{out} (pc)	0.64	0.022	H γ	4340	48	47	47	48
δr_{shell} (pc)	0.07	0.015	[O III]	4363	24	20	25	39
D (pc)	1500	900	He II	4686	95	94	49	62
n_{H} (cm^{-3})	36	55 000	[Ar IV]	4740	4.0	4.5	8.1	8.1
T_e (K)	15 400	16 400	H β	4861	100	100	100	100
ϵ	0.45	0.50	[O III]	5007	1131	957	1397	1345
$\log(\text{He}/\text{H})$	-1.02	-1.0	[N II]	5755	...	1.4	5.8	7.2
$\log(\text{N}/\text{H})$	-4.30	-3.89	He I	5876	1.9	1.9	10.9	7.5
$\log(\text{O}/\text{H})$	-3.76	-3.52	H α	6563	286	278	285	275
$\log(\text{Ne}/\text{H})$	-4.45	-4.15	[N II]	6584	12	41	110	118
$\log(\text{S}/\text{H})$	-5.19	-5.08	[S II]	6724	9.7	11	5.6	5.3
$\log(\text{Ar}/\text{H})$	-6.05	-5.80	[Ar III]	7135	5.8	2.2	21	8
			$L_{\text{H}\beta}$ (erg/s)	1E33	0.46	0.50	131	128




Figure 1. (Left) Density distribution, cross section, and morphology used for NGC 7027. (Right) Computed surface brightness in the He II $\lambda 4686$ line compared with the *HST* image.

geometry conflicts with the observed ionization structure and nebular line intensities. Our photoionization model outputs generally agree with the observations, except for some uncertain lines. Our photoionization model of NGC 7027 gives $D = 900$ pc, with the CSPN having $T_{\text{eff}} = 180$ kK and $L/L_{\odot} = 7500$, which reproduces most observations. Our distance agrees with previous determinations (Masson 1989; Volk & Kwok 1997; Zijlstra *et al.* 2008).

Acknowledgements

AD acknowledges receipt of an MQRES PhD Scholarship and an IAU Travel Grant.

References

Bernard Salas, J., Pottasch, S. R., Beintema, D. A., & Wesselius, P. R. 2001, *A&A*, 367, 949
 Danehkar, A., Frew, D. J., Parker, Q., & De Marco, O. 2011, *IAUS 281*, in press, arXiv:1109.2181
 Ercolano, B., Barlow, M. J., Storey, P. J., & Liu, X. W. 2003, *MNRAS*, 340, 1153
 Jacoby, G. H., Ferland, G. J., & Korista, K. T. 2001, *ApJ*, 560, 272
 Masson, C. R. 1989, *ApJ*, 336, 294
 Rauch, T. 2003, *A&A*, 403, 709
 Volk, K. & Kwok, S. 1997, *ApJ*, 477, 722
 Zhang, Y., Liu, X. W., Luo, S. G., Péquignot, D., & Barlow, M. J. 2005, *A&A*, 442, 249
 Zijlstra, A. A., van Hoof, P. A. M., & Perley, R. A. 2008, *ApJ*, 681, 1296