

¹University of Nottingham, Nottingham, United Kingdom and
²Derbyshire Healthcare NHS Foundation Trust, Derby, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.252

Aims. Functional Neurological Disorders (FNDs) affect motor or sensory functions without a detectable underlying disease. FNDs encompass a range of presentations including non-epileptic seizures, cognitive changes, weakness, and sensory symptoms. The prevalence of FND as a diagnosis is increasing rapidly. Following our clinical observations of a high prevalence of Attention-Deficit Hyperactivity Disorder (ADHD) in people referred with a previous diagnosis of FND to our tertiary Neuropsychiatry pilot service in Derbyshire, we conducted an integrative literature review with the aim to investigate the prevalence of ADHD in people diagnosed with FND.

Methods. We conducted an integrative literature review using a systematic approach. A literature search was performed on two databases, PubMed and ScienceDirect. The keywords 'Functional Neurological Disorder', 'Attention-Deficit Hyperactivity Disorder', 'Non-Epileptic Seizures', 'Functional' were used. Databases were searched for initial search on 31 November 2023 and the search was repeated on 31 January 2024. Only articles in English language were included. Studies were eligible if reporting the prevalence of ADHD in FND populations. Studies involving adults and children were included. A further search was conducted on reference lists from the selected articles.

Results. Database searches on PubMed and ScienceDirect had 298 and 11,837 results, respectively. Only seven studies were identified that explored the prevalence of ADHD in individuals diagnosed with a FND and were included. In the adult population an association between a FND diagnosis, and ADHD traits identified on screening, or a final ADHD diagnosis was identified. The findings also demonstrate an increased incidence of comorbid ADHD and FND with the presence of another co-existing neurodevelopmental disorder such as Autism Spectrum Disorder. Furthermore, results indicated that the prevalence of an ADHD diagnosis in children with a FND was higher compared with adults. The literature suggests that, in both adults and children with FND-related functional seizures there is an increased prevalence of comorbid ADHD.

Conclusion. In conclusion, the findings from this review demonstrate a lack of evidence looking into the prevalence of Attention-Deficit Hyperactivity Disorder in complex presentations being labelled as Functional Neurological Disorder. However, the existing literature indicates there is an association between FND and ADHD. These findings highlight the importance of considering potential ADHD comorbidity in the assessment and management of FND, potentially informing targeted treatment approaches for affected individuals. Further research could explore the efficacy of ADHD medication and similar dopamine modulating molecules in treating sub-cohort of people with FND.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

Impact of Covid-19 on Referrals to Paediatric Liaison Psychiatry at Children's Health Ireland (CHI) at Crumlin as the Pandemic Moved to Endemic Status

Mr Bohan Sun^{1*}, Dr Dimitrios Adamis²
 and Dr Fiona McNicholas^{1,3,4}

¹School of Medicine and Medical Science, University College Dublin, Dublin, Ireland; ²Sligo Mental Health Services, Sligo, Ireland;

³Department of Paediatric Liaison Psychiatry, Children's Health Ireland at Crumlin, Dublin, Ireland and ⁴Lucena Clinic Rathgar, St John of God Hospitaller Services, Dublin, Ireland

*Presenting author.

doi: 10.1192/bjo.2024.253

Aims. Rates of acute mental health presentations in youth were rising pre-pandemic internationally. Longitudinal studies following Covid-19 attest to ongoing deterioration in youth mental health, recognising adverse unintended consequences following public health restrictions.

This study aimed to examine whether the initial reported post-Covid-19 increase in mental health presentations persisted following the reclassification of Covid-19 to endemic status, which was accompanied by removal of most restrictions.

Methods. All referrals to paediatric liaison psychiatry (PLP) between Jan 2018–Dec 2022 in a Dublin tertiary children's hospital were included in the study. An interrupted time series analysis was conducted examining referrals with respect to different phases of Covid-19 and application of public health restrictions.

Results. 1,385 referrals to PLP were received over the 5-year study time-period. There was a significant decrease in PLP referrals immediately post Covid-19, following a significant and sustained increase as the pandemic progressed. Public health restriction phases had a unique effect on those presenting with suicidal ideation, with a significant increase in the number of referrals received. There was no effect of restrictions on other clinical profiles.

Conclusion. Increased referrals for youth with mental health difficulties, reported during the Covid-19 pandemic, persisted into the early endemic stage, after Covid-19 public health restriction have ceased. Potential impacts of restrictions on referrals of youth with suicidal ideation require further study. Investment in child and adolescent mental health services remain a priority, and future pandemic responses need to examine unintended consequences of any enforced public health measure.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

IRAMP: Investigation of Risk Assessment and Management Processes Using Datix Report Analysis and Observation of Clinical Team Meetings

Dr Kay Sunderland^{1*}, Dr Emma Drysdale¹, Dr Brian Gillatt¹, Dr Alan Mackenzie¹ and Dr Paula McCahon¹

¹NHS Greater Glasgow and Clyde, Glasgow, United Kingdom

*Presenting author.

doi: 10.1192/bjo.2024.254

Aims. To investigate risk assessment and management processes across a health board in the context of the implementation of a new risk screening tool and risk policy using comparison of DATIX incidents before and after implementation of the CRAFT tool.

In mental health services, risk assessment and management are key responsibilities for clinical staff. A risk management tool that is structured and evidence-based aims to assist staff in managing risks including violence, self-harm, suicide and self-neglect.

It is not clear whether risk tools have clinical utility in influencing risk-related decision making and previous reviews within the