INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, http://www.editorialmanager.com/psm/. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

- 1. S.I. units should be used throughout in text, figures and tables.
- 2. Authors should spell out in full any abbreviations used in their manuscripts.
- 3. Foreign quotations and phrases should be followed by a translation.
- 4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* 286, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* 32, 173–179.

Cleckley HJ (1941). The Mask of Sanity, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In Olfaction and the Brain (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank. org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. Line artwork: Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; Combination artwork (line/tone): Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; Black and white halftone artwork: Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; Colour halftone artwork: Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. Tables Tables should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

DEL//EIA/ ADTIOLE		The section of the section of the first order of the section of	
REVIEW ARTICLE		Unmet needs in patients with first-episode schizophrenia:	
Effects of cognitive therapy <i>versus</i> interpersonal psychotherapy in patients with major depressive disorder: a systematic review of randomized clinical trials with meta-analyses and trial sequential analyses		a longitudinal perspective Landolt K, Rössler W, Burns T, Ajdacic-Gross V, Galderisi S, Libiger J, Naber D, Derks EM, Kahn RS, Fleischhacker WW, for the EUFEST Study Group	1461
Jakobsen JC, Hansen JL, Simonsen S, Simonsen E & Gluud C	1343	Volume increases in putamen associated with positive	
ORIGINAL ARTICLES		symptom reduction in previously drug-naive schizophrenia after 6 weeks antipsychotic treatment	
Guided self-help concreteness training as an intervention for major depression in primary care: a Phase II randomized controlled trial		Li M, Chen Z, Deng W, He Z, Wang Q, Jiang L, Ma X, Wang Y, Chua SE, Cheung C, McAlonan GM, Sham PC, Collier DA, Gong Q & Li T	1475
Watkins ER, Taylor RS, Byng R, Baeyens C, Read R, Pearson K & Watson L	1359	Schizophrenia, culture and neuropsychology: sensory deficits, language impairments and social functioning in Chinese-speaking schizophrenia patients	0
Emotion modulates cognitive flexibility in patients with major depression			
Murphy FC, Michael A & Sahakian BJ	1373	Yang L, Chen S, Chen C-M, Khan F, Forchelli G & Javitt DC	1485
Course trajectories of unipolar depressive disorders identified by latent class growth analysis		Psychotic-like experiences in a community sample of 8000 children aged 9 to 11 years: an item response theory analysis Laurens KR, Hobbs MJ, Sunderland M, Green MJ & Mould GL	1495
Rhebergen D, Lamers F, Spijker J, de Graaf R, Beekman ATF & Penninx BWJH	1383	It's not what you hear, it's the way you think about it:	
Depressed mood enhances anxiety to unpredictable threat Robinson OJ, Overstreet C, Letkiewicz A & Grillon C	1397	appraisals as determinants of affect and behaviour in voice hearers Peters ER, Williams SL, Cooke MA & Kuipers E	1507
Affective symptoms and intra-individual variability in the short-term course of cognitive functioning in bipolar disorder Depp CA, Savla GN, Vergel de Dios LA, Mausbach BT & Palmer BW	1409	Risk of schizophrenia in relation to parental origin and genome-wide divergence Pedersen CB, Demontis D, Pedersen MS, Agerbo E,	
Abnormal anterior cingulate cortical activity during emotional n -back task performance distinguishes bipolar from unipolar depressed females	1417	Mortensen PB, Børglum AD, Hougaard DM, Hollegaard MV, Mors O & Cantor-Graae E	1515
		Pituitary volume, stress reactivity and genetic risk for psychotic disorder	
Bertocci MA, Bebko GM, Mullin BC, Langenecker SA, Ladouceur CD, Almeida JRC & Phillips ML		Habets P, Collip D, Myin-Germeys I, Gronenschild E, Van Bronswijk S, Hofman P, Lataster T, Lardinois M, Nicolson NA,	.=
Cognitive behaviour therapy and supportive therapy for bipolar disorders: relapse rates for treatment period and		van Os J, Marcelis M, for G.R.O.U.P.	1523
2-year follow-up Meyer TD & Hautzinger M	1429	Is maternal smoking during pregnancy a causal environmental risk factor for adolescent antisocial behavior?	
,	1429	Testing etiological theories and assumptions D'Onofrio BM, Van Hulle CA, Goodnight JA, Rathouz PJ	
Home foreclosure and risk of psychiatric morbidity during the recent financial crisis		& Lahey BB	1535
McLaughlin KA, Nandi A, Keyes KM, Uddin M, Aiello AE, Galea S & Koenen KC	1441	Disentangling the relationships between maternal smoking during pregnancy and co-occurring risk factors	
Co-morbid anxiety disorders in bipolar disorder and major depression: familial aggregation and clinical characteristics of co-morbid panic disorder, social phobia, specific phobia and obsessive-compulsive disorder		Ellingson JM, Rickert ME, Lichtenstein P, Långström N & D'Onofrio BM	1547
		Correspondence	1559
Goes FS, McCusker MG, Bienvenu OJ, MacKinnon DF, Mondimore FM, Schweizer B, National Institute of Mental Health			



1449



& Potash JB

Genetics Initiative Bipolar Disorder Consortium, DePaulo Jr. JR

Published online by Cambridge University Press