

RESEARCH ARTICLE

Disparity-persistence and the multistep friendship paradox

Kenneth S. Berenhaut  and Chi M. Zhang

Department of Statistical Sciences, Wake Forest University, Winston Salem, NC, USA

Corresponding author: Kenneth S. Berenhaut; Email: berenhks@wfu.edu

Keywords: friendship paradox; disparity-persistence; random walks; random paths; long-range degree correlation; core-periphery structure

MSC: 05C07; 05C38; 05C50; 05C81; 05C90; 60J10; 91D30

Abstract

In this paper, we consider the friendship paradox in the context of random walks and paths. Among our results, we give an equality connecting long-range degree correlation, degree variability, and the degree-wise effect of additional steps for a random walk on a graph. Random paths are also considered, as well as applications to acquaintance sampling in the context of core-periphery structure.

1. Introduction

The friendship paradox, introduced by Feld in [15], states roughly that your friends have more friends than you do on average (for an explicit statement, see [Theorem 1.1](#)). Following the work of [24], extending the friendship paradox to multiple steps (i.e., iterated friendships), here we quantify an explicit connection between long-range degree correlation, degree variability, and the degree-wise effect of additional steps for random walks on a graph. Results for random paths are also considered.

Throughout, we suppose $G = (\mathcal{V}, \mathcal{E})$ is a connected graph with node set $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$ and undirected edge set \mathcal{E} and define the *degree function* d , so that for $v \in \mathcal{V}$, $d(v)$ is the degree of v (i.e., the number of neighbors of v) in G . Moreover, we denote by $A = [A_{ij}]$ the associated $n \times n$ adjacency matrix; the degree of a node v_i can then be computed via

$$d(v_i) = \sum_{j=1}^n A_{ij}.$$

As we will be interested in the expected degree over random sequences of nodes in the graph, it will be convenient to consider a time-homogeneous random walk $\mathbf{X} = (X_0, X_1, \dots)$ dictated by a transition matrix, $\mathbf{P} = [P_{ij}]$, with

$$P_{ij} \stackrel{\text{def}}{=} \mathbb{P}(X_{k+1} = v_j \mid X_k = v_i) = \frac{A_{ij}}{d(v_i)},$$

for $1 \leq i, j \leq n$, and $k \geq 0$. Importantly, we will assume throughout that X_0 is uniformly selected from \mathcal{V} .

We first restate the friendship paradox formalized in [15].

Theorem 1.1. ([15]) *Suppose X_0 is a node selected uniformly at random from \mathcal{V} , and $E = \{V, W\}$ is an edge pair selected uniformly at random from \mathcal{E} . Then, selecting Y_1 from the nodes in E , each with*

half chance,

$$\mathbb{E}(d(Y_1)) \geq \mathbb{E}(d(X_0)). \tag{1}$$

Similarly, several authors have employed the following counterpart for random walks with a uniformly selected initial node X_0 (see for instance [3, 6, 21]).

Theorem 1.2. *Suppose $X = (X_0, X_1, X_2, \dots)$ is a simple random walk on the graph G , where X_0 is selected uniformly at random from \mathcal{V} . Then*

$$\mathbb{E}(d(X_1)) \geq \mathbb{E}(d(X_0)). \tag{2}$$

The following two results regarding multiple step walks and paths can be found in [24].

Theorem 1.3. ([24]) *Suppose $X = (X_0, X_1, X_2, \dots)$ is a simple random walk on the graph G , where X_0 is selected uniformly at random from \mathcal{V} . Then, for $k \geq 0$,*

$$\mathbb{E}(d(X_k)) \geq \mathbb{E}(d(X_0)). \tag{3}$$

Theorem 1.4. ([24]) *Suppose $k \geq 1$ is odd, X_0 is selected uniformly at random from \mathcal{V} , and $W = \{Y_0, Y_1, \dots, Y_k\}$ is a path selected uniformly at random from the set of all length- k paths on G . Then, we have*

$$\mathbb{E}(d(Y_k)) \geq \mathbb{E}(d(X_0)). \tag{4}$$

For further recent theoretical results regarding the friendship paradox, see for instance [4, 5, 12, 31].

In place of comparisons with $\mathbb{E}(d(X_0))$, as in the results above, the primary concern of this paper is the study of the degree-wise benefit of additional steps on the graph (for both random walks and random paths). Motivation for such considerations is provided by the recent success of applications of the one-step friendship paradox (see [8, 9, 14, 18, 20, 23, 34]).

Several authors have alluded to weakness in expression of the friendship paradox in networks exhibiting positive degree correlation over edges (see for instance [5, 25, 28, 31]). Before formally stating our results, it will be convenient to consider longer-range degree correlation. It is common for networks to exhibit sharing of similar characteristics across edges (i.e., “homophily,” “birds of a feather flock together,” “assortativity”; see [29]). In social networks, in particular, individuals may prefer to be associated with others sharing, for instance, similar age, education, or occupations. In the case where the characteristic of interest is nodal degree, Pearson correlation $\rho = \rho(G)$ refers to the tendency of nodes in a network to be associated with others sharing similar (or different) degrees. Specifically, for the graph G , suppose $E = \{V, W\}$ is an edge selected uniformly at random, then

$$\rho \stackrel{\text{def}}{=} \text{cor}(d(V), d(W)) = \frac{\text{cov}(d(V), d(W))}{\text{sd}(d(V)) \text{sd}(d(W))}. \tag{5}$$

A positive degree correlation, $\rho > 0$, indicates a tendency for high-degree nodes in the graph to connect to other high-degree nodes and similarly for low-degree nodes.

Values of ρ have been studied in various classes of networks (see [7, 16, 26, 33]). For connections between assortativity and other topological network properties, see [11, 13, 36, 37]. For work considering assortativity in the context of the friendship paradox, see for instance [22, 25, 28, 31]. See [30] and the references therein for some further work on assortativity.

Although much previous research has focused on degree correlation among nearest neighbors, some recent considerations of the concept of assortative mixing beyond first neighbors can be found in [1, 2, 17, 27] and can in a general sense be referred to as *long-range degree correlation* (see [2]).

Assortativity, as defined in [29], can be interpreted as a measure of the correlation between nodal degrees based on the first-order adjacency matrix. Here, we define the k th-order path-based degree correlation as the Pearson coefficient, measuring the correlation between degrees for the two end points of a randomly chosen *path* with length k .

Definition 1.5. For $k \geq 0$, let Y_k^- and Y_k^+ be the beginning and terminal nodes for a path selected uniformly at random from the set of all length- k paths on G . The k th-order (path-based) degree correlation $\rho_{0,k}$ is given by

$$\rho_{0,k} \stackrel{\text{def}}{=} \text{cor}(d(Y_k^-), d(Y_k^+)) = \frac{\text{cov}(d(Y_k^-), d(Y_k^+))}{\text{sd}(d(Y_k^-)) \text{sd}(d(Y_k^+))}.$$

In parallel with the definition above, we also consider the walk-based degree correlation as that for the starting node, X_0 , of a random walk $\mathbf{X} = (X_0, X_1, X_2, \dots)$ (with uniform initial distribution) and the node reached at the walk’s k th step, X_k . Note that the use of the uniform distribution here is motivated by the desired applications. This is in contrast to recent work regarding long-range correlation, wherein the initial distribution is stationary for the corresponding Markov chain (see [2, 19, 32]).

Definition 1.6. Consider a random walk, $\mathbf{X} = \{X_0, X_1, X_2, \dots\}$, on the graph G (with uniform initial distribution). The k th-order (walk-based) degree correlation, $\phi_{0,k} = \phi_{0,k}(\mathbf{X})$, is given by

$$\phi_{0,k} \stackrel{\text{def}}{=} \text{cor}(d(X_0), d(X_k)) = \frac{\text{cov}(d(X_0), d(X_k))}{\text{sd}(d(X_0)) \text{sd}(d(X_k))}.$$

Note. As an aside, it may be of some value to consider applications for which $\phi_{0,1}$ as in Definition 1.6 may be of interest. The correlation computed involves the same node pairs as in (5) but with weighting in a manner that places equal emphases on each node regardless of degree. Ties in a sense are weaker (or diluted) for nodes of large degree. □

Now, for the random walk $\mathbf{X} = (X_0, X_1, \dots)$, let X_∞ be a node selected according to the stationary distribution $\pi = (\pi_1, \dots, \pi_n)$ of the random walk \mathbf{X} , that is, $\pi_i = d(v_i) / \sum_j d(v_j)$. A main quantity of interest will be what we term the *proportional residual benefit* at time k for the walk, which measures the degree-wise (remaining) benefit of additional steps of a random walk.

Definition 1.7. The *proportional residual benefit* for the walk \mathbf{X} at time k , $\tau_k = \tau_k(\mathbf{X})$ is given by

$$\tau_k \stackrel{\text{def}}{=} \frac{\mathbb{E}(d(X_\infty)) - \mathbb{E}(d(X_k))}{\mathbb{E}(d(X_k))}.$$

We now have the following quantitative relationship between the walk-based long-range degree correlation and the proportional residual benefit. The proof is provided in Section 2. Throughout, for a random variable U with finite second moment, we denote by $c_v(U)$, the *coefficient of variation* of U , that is,

$$c_v(U) \stackrel{\text{def}}{=} \frac{\text{sd}(U)}{\mathbb{E}(U)}. \tag{6}$$

Theorem 1.8. Suppose $X = (X_0, X_1, \dots)$ is a random walk on the graph G . The proportional residual benefit τ_k can be written as the product of the k th-order degree correlation $\phi_{0,k}$ and the two coefficients of variation, $c_v(d(X_0))$ and $c_v(d(X_k))$, that is

$$\begin{aligned} \tau_k &= \text{cor}(d(X_0), d(X_k)) \cdot \frac{\text{sd}(d(X_0))}{\mathbb{E}(d(X_0))} \cdot \frac{\text{sd}(d(X_k))}{\mathbb{E}(d(X_k))} \\ &= \phi_{0,k} c_v(d(X_0)) c_v(d(X_k)). \end{aligned} \tag{7}$$

Now, as in Definition 1.5, for $k \geq 0$, let Y_k^- and Y_k^+ be the beginning and terminal nodes for a path selected uniformly at random from the set of all length- k paths. We have the following definition and result.

Definition 1.9. Suppose $k \geq 0$. The proportional one-step benefit at length k , γ_k , is given by

$$\gamma_k \stackrel{\text{def}}{=} \frac{\mathbb{E}(d(Y_{k+1}^+)) - \mathbb{E}(d(Y_k^+))}{\mathbb{E}(d(Y_k^+))}.$$

Theorem 1.10. The proportional one-step benefit, γ_k , at length k can be written as the product of the k th-order degree correlation $\rho_{0,k}$ and the two coefficients of variation, $c_v(d(Y_k^-))$ and $c_v(d(Y_k^+))$, that is,

$$\begin{aligned} \gamma_k &= \text{cor}(d(Y_k^-), d(Y_k^+)) \cdot \frac{\text{sd}(d(Y_k^-))}{\mathbb{E}(d(Y_k^-))} \cdot \frac{\text{sd}(d(Y_k^+))}{\mathbb{E}(d(Y_k^+))} \\ &= \rho_{0,k} c_v(d(Y_k^-)) c_v(d(Y_k^+)). \end{aligned} \tag{8}$$

In terms of residual benefit, in the case of random paths, we will also prove the following.

Theorem 1.11. Suppose G is a non-bipartite (and connected) graph. If (a) $k \geq 0$ is even or (b) $k \geq 1$ is odd and the corresponding k th-order path-based degree correlation $\rho_{0,k}$ is nonnegative, then the limiting expected degree of Y_i^+ is no less than that of Y_k^+ , that is,

$$\lim_{i \rightarrow \infty} \mathbb{E}(d(Y_i^+)) \geq \mathbb{E}(d(Y_k^+)). \tag{9}$$

Note. (Disparity persistence and core-periphery structure) Theorems 1.8, 1.10, and 1.11 provide insight into when it may be beneficial in acquaintance sampling and elsewhere to continue on to neighbors of neighbors in an iterated fashion. In fact, the benefit can be high in networks wherein the degree correlation is positive out to a longer range, but, globally, the degree variability is high. We refer to such a phenomenon as *disparity-persistence*, since disparity in degree persists over extended neighborhoods. A prime example wherein such behavior can occur is social networks exhibiting strong core-periphery structure, with a large core of high-degree nodes and a periphery of loosely connected nodes, with low degree. □

The remainder of the paper proceeds as follows. In Section 2, we prove Theorem 1.8, while in Section 3, we prove Theorems 1.10 and 1.11.

2. Proof of Theorem 1.8 (The random walk case)

In this section, we will prove Theorem 1.8.

Proof of Theorem 1.8 Set $d_i = d(v_i)$, $\mathbf{d} = (d_1, d_2, \dots, d_n)'$, and $\mathbf{1} = (1, 1, \dots, 1)'$. Let \mathbf{P} be the transition matrix for the random walk, \mathbf{X} , and note that $\mathbf{d} = \mathbf{A}\mathbf{1}$. Now, note that

$$\mathbb{E}(d(X_0) d(X_k)) = \frac{1}{n} \sum_i \sum_j d_i d_j (\mathbf{P}^k)_{ij} = \frac{\mathbf{d}' \mathbf{P}^k \mathbf{d}}{\mathbf{1}' \mathbf{1}}.$$

Hence, letting \mathbf{D} be the diagonal matrix with diagonal \mathbf{d} and noting that $\mathbf{P} = \mathbf{D}^{-1} \mathbf{A}$ gives

$$\begin{aligned} \mathbf{d}' \mathbf{P}^k \mathbf{d} &= \mathbf{d}' \mathbf{D}^{-1} \mathbf{A} \mathbf{P}^{k-1} \mathbf{d} = \mathbf{1}' \mathbf{A} \mathbf{P}^{k-1} \mathbf{d} = \mathbf{d}' \mathbf{P}^{k-1} \mathbf{d} \\ &= \dots = \mathbf{d}' \mathbf{d}. \end{aligned}$$

Thus,

$$\mathbb{E}(d(X_0) d(X_k)) = (\mathbf{d}' \mathbf{d}) / (\mathbf{1}' \mathbf{1}).$$

Now,

$$\begin{aligned} \mathbb{E}(d(X_\infty)) &= \sum_i \left(\frac{d_i}{\sum_j d_j} \right) d_i = \frac{\mathbf{d}' \mathbf{d}}{\mathbf{1}' \mathbf{A} \mathbf{1}}, \\ \mathbb{E}(d(X_0)) &= \frac{1}{n} \sum_i d_i = \frac{\mathbf{1}' \mathbf{A} \mathbf{1}}{\mathbf{1}' \mathbf{1}}, \end{aligned}$$

and hence

$$\mathbb{E}(d(X_\infty)) \cdot \mathbb{E}(d(X_0)) = \frac{\mathbf{d}' \mathbf{d}}{\mathbf{1}' \mathbf{A} \mathbf{1}} \frac{\mathbf{1}' \mathbf{A} \mathbf{1}}{\mathbf{1}' \mathbf{1}} = \frac{\mathbf{d}' \mathbf{d}}{\mathbf{1}' \mathbf{1}} = \mathbb{E}(d(X_0) d(X_k)).$$

Thus,

$$\begin{aligned} \tau_k &= \frac{\mathbb{E}(d(X_\infty)) - \mathbb{E}(d(X_k))}{\mathbb{E}(d(X_k))}, \\ &= \frac{1}{\mathbb{E}(d(X_k))} \left(\frac{\mathbb{E}(d(X_0) d(X_k))}{\mathbb{E}(d(X_0))} - \mathbb{E}(d(X_k)) \right), \\ &= \frac{\text{cov}(d(X_0), d(X_k))}{\mathbb{E}(d(X_k)) \cdot \mathbb{E}(d(X_0))}, \\ &= \phi_{0,k} \left(\frac{\text{sd}(d(X_0))}{\mathbb{E}(d(X_0))} \right) \cdot \left(\frac{\text{sd}(d(X_k))}{\mathbb{E}(d(X_k))} \right) = \phi_{0,k} c_v(d(X_0)) c_v(d(X_k)). \end{aligned} \quad \square$$

We easily derive the following corollary.

Corollary 2.1. *Suppose $\mathbf{X} = (X_0, X_1, \dots)$ is a random walk on the graph $G = (\mathcal{V}, \mathcal{E})$. For $k \geq 1$, the expected degree of X_∞ is no less than that of X_k if and only if the k th-order degree correlation $\phi_{0,k}$ is nonnegative, that is,*

$$\mathbb{E}(d(X_\infty)) \geq \mathbb{E}(d(X_k)) \quad \text{if and only if} \quad \phi_{0,k} = \text{cor}(d(X_0), d(X_k)) \geq 0. \tag{10}$$

3. Proof of Theorems 1.10 and 1.11 (The random path case)

In this section, we will prove Theorem 1.10 and further discuss the relationship between the proportional one-step benefit γ_k and the path-based degree correlation $\rho_{0,k}$ in parallel with the random-walk case, $(X_k)_{k \geq 0}$. As per convention, throughout, we will take the zeroth power of the adjacency matrix, \mathbf{A} , of a graph G (i.e., \mathbf{A}^0) to be the $n \times n$ identity matrix, and hence for $k \geq 0$, the entry $(\mathbf{A}^k)_{ij}$ counts the number of distinct paths of length k connecting nodes v_i and v_j .

As before, suppose for $i \geq 0$, Y_i^- and Y_i^+ are the beginning and terminal nodes of a path selected uniformly at random from the set of all length- i paths on G . We have the following lemma (see also Lemma 4 in [24]).

Lemma 3.1. *Suppose $k \geq 0$. The expected degree of Y_k^+ can be written in terms of successive entries in the sequence (N_0, N_1, \dots) , where N_j is the number of length j paths on G , that is,*

$$\mathbb{E}(d(Y_k^+)) = \frac{N_{k+1}}{N_k}. \tag{11}$$

Proof. We have

$$\mathbb{E}(d(Y_k^+)) = \sum_{j=1}^n d_j \frac{\sum_{i=1}^n A_{i,j}^k}{N_k} = \frac{\mathbf{d}'(\mathbf{A}^k)\mathbf{1}}{N_k} = \frac{\mathbf{1}'\mathbf{A}\mathbf{A}^k\mathbf{1}}{N_k} = \frac{N_{k+1}}{N_k},$$

where the third equality follows from the symmetry of the adjacency matrix, \mathbf{A} . □

Now we turn to a proof of Theorem 1.10.

Proof of Theorem 1.10 Suppose $k \geq 0$ and Y_k^- and Y_k^+ are two nodes connected by a randomly chosen path with length k . We have

$$\mathbb{E}(d(Y_k^-) d(Y_k^+)) = \sum_{i=1}^n \sum_{j=1}^n d_i d_j \frac{A_{i,j}^k}{N_k} = \frac{\mathbf{d}'\mathbf{A}^k\mathbf{d}}{N_k} = \frac{N_{k+2}}{N_k}. \tag{12}$$

Note that by Lemma 3.1 and symmetry,

$$\mathbb{E}(d(Y_k^-)) = \frac{N_{k+1}}{N_k} = \mathbb{E}(d(Y_k^+)). \tag{13}$$

Hence, using Eqs. (12) and (13),

$$\begin{aligned} \text{cov}(d(Y_k^-), d(Y_k^+)) &= \frac{N_{k+2}}{N_k} - \left(\frac{N_{k+1}}{N_k}\right)^2 = \frac{N_{k+1}}{N_k} \left(\frac{N_{k+2}}{N_{k+1}} - \frac{N_{k+1}}{N_k}\right), \\ &= \mathbb{E}(d(Y_k^+)) \cdot (\mathbb{E}(d(Y_{k+1}^+)) - \mathbb{E}(d(Y_k^+))), \end{aligned} \tag{14}$$

and hence by the definition of γ_k , and Eqs. (13) and (14),

$$\gamma_k = \frac{\mathbb{E}(d(Y_{k+1}^+)) - \mathbb{E}(d(Y_k^+))}{\mathbb{E}(d(Y_k^+))} \cdot \frac{\mathbb{E}(d(Y_k^+))}{\mathbb{E}(d(Y_k^-))} \cdot \frac{\text{sd}(d(Y_k^-))}{\text{sd}(d(Y_k^+))} \cdot \frac{\text{sd}(d(Y_k^+))}{\text{sd}(d(Y_k^-))},$$

$$\begin{aligned}
 &= \frac{\text{cov}(d(Y_k^-), d(Y_k^+))}{\text{sd}(d(Y_k^-)) \text{sd}(d(Y_k^+))} \cdot \frac{\text{sd}(d(Y_k^-))}{\mathbb{E}(d(Y_k^-))} \cdot \frac{\text{sd}(d(Y_k^+))}{\mathbb{E}(d(Y_k^+))}, \\
 &= \rho_{0,k} c_v(d(Y_k^-)) c_v(d(Y_k^+)). \quad \square
 \end{aligned}
 \tag{15}$$

Therefore, the proportional increase in relative expected degree for an increase of one in the length of our random path can be determined by the corresponding path-based degree correlation, along with the two coefficients of variation. We have the following corollary.

Corollary 3.2. *Suppose $k \geq 0$. The expected degree of Y_{k+1}^+ is no less than that of Y_k^+ if and only if the k -th-order degree correlation $\rho_{0,k}$ is nonnegative, that is,*

$$\mathbb{E}(d(Y_{k+1}^+)) \geq \mathbb{E}(d(Y_k^+)) \quad \text{if and only if} \quad \rho_{0,k} = \text{cor}(d(Y_k^-), d(Y_k^+)) \geq 0. \tag{16}$$

Note that employing an inequality on the number of paths, developed by [35] (with $c = 1$ and $b = 0$), we have

$$\frac{N_{2a+1}}{N_{2a}} \leq \frac{N_{2a+2}}{N_{2a+1}} \tag{17}$$

for $a \geq 0$, and hence by Lemma 3.1, when k is even, $\mathbb{E}(d(Y_{k+1}^+)) \geq \mathbb{E}(d(Y_k^+))$. In the case $k = 0$, this is simply the result of Feld in Theorem 1.1. In addition, by Corollary 3.2, for any network G , G is assortative (i.e., $\rho_{0,1} > 0$) if and only if $\mathbb{E}(d(Y_2^+)) > \mathbb{E}(d(Y_1^+))$. For consideration of degree comparison for X_1 and X_2 under a random graph model, see [4].

The following lemma follows directly from eigen decomposition of the matrix A and Lemma 3.1 (see for instance [10]).

Lemma 3.3. ([10]) *Suppose $k \geq 0$ and the graph G is non-bipartite. The expected degree of Y_k^+ tends to λ_1 , the largest eigenvalue of the adjacency matrix, A , as k tends to infinity, that is,*

$$\lim_{k \rightarrow \infty} \mathbb{E}(d(Y_k^+)) = \lambda_1. \tag{18}$$

We now turn to a proof of Theorem 1.11

Proof of Theorem 1.11 Suppose $m \geq 0$. From Rayleigh’s inequality, we have

$$\mathbb{E}(d(Y_{2m}^+)) = \frac{N_{2m+1}}{N_{2m}} = \frac{(A^m \mathbf{1})' A (A^m \mathbf{1})}{(A^m \mathbf{1})' \cdot (A^m \mathbf{1})} \leq \lambda_1. \tag{19}$$

The result for k even follows by Lemma 3.3. Now, suppose k is odd and $\rho_{0,k} \geq 0$. Employing Corollary 3.2 and Eq. (19) then gives

$$\mathbb{E}(d(Y_k^+)) \leq \mathbb{E}(d(Y_{k+1}^+)) \leq \lambda_1,$$

and the result follows. □

Competing interests. The authors declare no conflict of interest.

References

[1] Allen-Perkins, A., Pastor, J.M., & Estrada, E. (2017). Two-walks degree assortativity in graphs and networks. *Applied Mathematics and Computation* **311**: 262–271.

- [2] Arcagni, A., Grassi, R., Stefani, S., & Torriero, A. (2017). Higher order assortativity in complex networks. *European Journal of Operational Research* **262**(2): 708–719.
- [3] Berenhaut, K.S. & Jiang, H. (2019). The friendship paradox for weighted and directed networks. *Probability in the Engineering and Informational Sciences* **33**(1): 136–145.
- [4] Berenhaut, K.S., Jiang, H., McNab, K.M., & Krizay, E.J. (2018). The degree-wise effect of a second step for a random walk on a graph. *Journal of Applied Probability* **55**(4): 1203–1210.
- [5] Cantwell, G.T., Kirkley, A., & Newman, M. (2021). The friendship paradox in real and model networks. *Journal of Complex Networks* **9**(2): cnab011.
- [6] Cao, Y. & Ross, S.M. (2016). The friendship paradox. *Mathematical Scientist* **41**(1): 61–64.
- [7] Chen, N. & Olvera-Cravioto, M. (2013). Directed random graphs with given degree distributions. *Stochastic Systems* **3**(1): 147–186.
- [8] Christakis, N.A. & Fowler, J.H. (2010). Social network sensors for early detection of contagious outbreaks. *PLoS One* **5**(9): e12948.
- [9] Cohen, R., Havlin, S., & Ben-Avraham, D. (2003). Efficient immunization strategies for computer networks and populations. *Physical Review Letters* **91**(24): 247901.
- [10] Cvetkovic, D., Cvetković, D.M., Rowlinson, P., & Simic, S. (1997). *Eigenspaces of graphs* **66**, Cambridge: Cambridge University Press.
- [11] D'Agostino, G., Scala, A., Zlatić, V., & Caldarelli, G. (2012). Robustness and assortativity for diffusion-like processes in scale-free networks. *Europhysics Letters* **97**(6): 68006.
- [12] Dizon-Ross, R. & Ross, S.M. (2022). A probabilistic friendship network model. *Probability in the Engineering and Informational Sciences* **36**(1): 1–16.
- [13] Ellens, W., Spieksma, F.M., Van Mieghem, P., Jamakovic, A., & Kooij, R.E. (2011). Effective graph resistance. *Linear Algebra and its applications* **435**(10): 2491–2506.
- [14] Eom, Y.-H. & Jo, H.-H. (2015). Tail-scope: Using friends to estimate heavy tails of degree distributions in large-scale complex networks. *Scientific Reports* **5**: 9752. doi:[10.1038/srep09752](https://doi.org/10.1038/srep09752).
- [15] Feld, S.L. (1991). Why your friends have more friends than you do. *American Journal of Sociology* **96**(6): 1464–1477.
- [16] Foster, J.G., Foster, D.V., Grassberger, P., & Paczuski, M. (2010). Edge direction and the structure of networks. *Proceedings of the National Academy of Sciences of the United States of America* **107**(24): 10815–10820.
- [17] Fujiki, Y., Takaguchi, T., & Yakubo, K. (2018). General formulation of long-range degree correlations in complex networks. *Physical Review E* **97**(6): 062308.
- [18] Garcia-Herranz, M., Moro, E., Cebrian, M., Christakis, N.A., & Fowler, J.H. (2014). Using friends as sensors to detect global-scale contagious outbreaks. *PLoS One* **9**(4): e92413.
- [19] Gutiérrez-Gómez, L. & Delvenne, J.-C. (2019). Multi-hop assortativities for network classification. *Journal of Complex Networks* **7**(4): 603–622.
- [20] Herrera, J.L., Srinivasan, R., Brownstein, J.S., Galvani, A.P., & Meyers, L.A. (2016). Disease surveillance on complex social networks. *PLoS Computational Biology* **12**(7): e1004928.
- [21] Jackson, M.O. (2019). The friendship paradox and systematic biases in perceptions and social norms. *Journal of Political Economy* **127**(2): 777–818. doi:[10.1086/701031](https://doi.org/10.1086/701031).
- [22] Jo, H.-H. & Eom, Y.-H. (2014). Generalized friendship paradox in networks with tunable degree-attribute correlation. *Physical Review E* **90**(2): 022809.
- [23] Kim, D.A., Hwong, A.R., Stafford, D., Hughes, D.A., O'Malley, A.J., Fowler, J.H., & Christakis, N.A. (2015). Social network targeting to maximise population behaviour change: A cluster randomised controlled trial. *The Lancet* **386**(9989): 145–153.
- [24] Kramer, J.B., Cutler, J., & Radcliffe, A. (2016). The multistep friendship paradox. *American Mathematical Monthly* **123**(9): 900–908.
- [25] Lee, E., Lee, S., Eom, Y.-H., Holme, P., & Jo, H.-H. (2019). Impact of perception models on friendship paradox and opinion formation. *Physical Review E* **99**(5): 052302.
- [26] Li, C., Wang, H., & Van Mieghem, P. (2013). Epidemic threshold in directed networks. *Physical Review E* **88**(6): 062802.
- [27] Mayo, M., Abdelzahr, A., & Ghosh, P. (2015). Long-range degree correlations in complex networks. *Computational Social Networks* **2**(1): 1–13.
- [28] Momeni, N. & Rabbat, M.G. (2018). Effectiveness of alter sampling in social networks. Preprint, arXiv:1812.03096.
- [29] Newman, M.E. (2002). Assortative mixing in networks. *Physical Review letters* **89**(20): 208701.
- [30] Noldus, R. & Van Mieghem, P. (2015). Assortativity in complex networks. *Journal of Complex Networks* **3**(4): 507–542.
- [31] Pal, S., Yu, F., Novick, Y., Swami, A., & Bar-Noy, A. (2019). A study on the friendship paradox—quantitative analysis and relationship with assortative mixing. *Applied Network Science* **4**(1): 1–26.
- [32] Peel, L., Delvenne, J.-C., & Lambiotte, R. (2018). Multiscale mixing patterns in networks. *Proceedings of the National Academy of Sciences of the United States of America* **115**(16): 4057–4062.
- [33] Piraveenan, M., Prokopenko, M., & Zomaya, A.Y. (2012). Assortative mixing in directed biological networks *IEEE/ACM Transactions on Computational Biology and Bioinformatics* **9**(1): 66–78. doi:[10.1109/TCBB.2010.80](https://doi.org/10.1109/TCBB.2010.80).
- [34] Singer, Y. (2016). Influence maximization through adaptive seeding. *ACM SIGecom Exchanges* **15**(1): 32–59.

- [35] Täubig, H., Weihmann, J., Kosub, S., Hemmecke, R., & Mayr, E.W. (2013). Inequalities for the number of walks in graphs. *Algorithmica* **66**(4): 804–828.
- [36] Van Mieghem, P., Ge, X., Schumm, P., Trajanovski, S., & Wang, H. (2010). Spectral graph analysis of modularity and assortativity. *Physical Review E* **82**(5): 056113.
- [37] Van Mieghem, P., Wang, H., Ge, X., Tang, S., & Kuipers, F.A. (2010). Influence of assortativity and degree-preserving rewiring on the spectra of networks. *The European Physical Journal B* **76**(4): 643–652.