

# Blind Search for Variability in Planck Data

Jörg P. Rachen<sup>1</sup>, Elina Keihänen<sup>2</sup> and Martin Reinecke<sup>3</sup>

<sup>1</sup>Department of Astrophysics/IMAPP, Radboud University Nijmegen, The Netherlands

<sup>2</sup>Department of Physics, University of Helsinki, Finland

<sup>3</sup>MPI for Astrophysics, 85741 Garching, Germany

**Abstract.** The sky is full of variable and transient sources on all time scales, from milliseconds to decades. *Planck*'s regular scanning strategy makes it an ideal instrument to search for variable sky signals in the millimetre and submillimetre regime, on time scales from hours to several years. A precondition is that instrumental noise and systematic effects, caused in particular by non-symmetric beam shapes, are properly removed. We present a method to perform a full sky blind search for variable and transient objects at all Planck frequencies.

**Keywords.** space vehicles: instruments, methods: statistical, techniques: miscellaneous

## 1. Introduction

*Planck* (Tauber *et al.* 2010, Planck Collaboration I. 2011) provides 8 full sky surveys at the frequencies of its LFI instrument (30, 44, and 70 GHz), and 5 full surveys at the 6 HFI frequencies in the range 100–857 GHz. The satellite rotates with 1 rpm around an axis kept fixed for a pointing period (PID) of about 50 minutes, then shifts along the ecliptic by 2°. This keeps a point source near the ecliptic in the main beam (2 FWHM) for about 5–30 PIDs (depending on frequency) for each survey; for sources near the ecliptic poles the coverage can be much larger. Analysing the time information in Planck data for a particular sky direction thus allows to search for variability in the sky signal.

## 2. Mapping of time ordered information


Variability mapping is based on four-dimensional Healpix (Górski *et al.* 2005) constructs called 4D-maps, which record for every sky pixel  $k$  all contributions of a given detector at times  $t_j$  and beam orientation  $b_j$ , where the index  $j$  refers to Planck PIDs. To construct an average sky signal  $S_k$  free from beam orientation effects, we use the *ArtDeco* beam deconvolution code (Keihänen & Reinecke 2012). A variability map is then a two-dimensional Healpix map of a quantity

$$X_k = \frac{1}{2} \operatorname{sgn}(\hat{\chi}_k^2 - 1) N_k \left[ \hat{\chi}_k^2 - 1 - \ln \hat{\chi}_k^2 \right]$$

where

$$\hat{\chi}_k^2 \equiv \frac{\chi_k^2}{N_k} = \frac{1}{N_k} \sum_j \frac{(I_{kj} - (S*b)_{kj})^2}{\sigma_k^2} \quad \text{with} \quad \sigma_{kj}^2 = \sigma_n^2 + \xi^2 (S*b)_{kj}^2.$$

$N_k$  is the number of entries for pixel  $k$  in the 4D signal map  $I_{kj}$ ,  $(S*b)_{kj}$  is the beam-reconvolved average 4D-map based on the *ArtDeco* map  $S_k$ ,  $\sigma_n$  is the detector white noise, and  $\xi$  combines all instrumental fluctuations which factor on the signal (e.g., calibration, inaccuracies of the beam model). The definition of  $X$  is motivated by the Chernoff bound on the CDF  $P_N(\chi^2)$ , with  $X \leq -\ln(1 - P_N(\chi^2))$  for  $\hat{\chi}^2 \geq 1$ , and  $X > \ln P_N(\chi^2)$  for



**Figure 1.** Histogram of an X-map for the LFI 44 GHz detectors for  $\xi = 0.01$ , and detector noise  $n_s$  as given in Planck Collaboration II. (2014), thin dotted lines show the distribution expected for pure Gaussian noise. The tail to large values of  $X$  indicates the presence of true sky variability. A tail to large negative values of  $X$  would indicate an overestimation of  $\xi$ , an offset of the peak from  $X = 0$  an incorrect estimation of  $n_s$ .

$\hat{\chi}^2 < 1$ . The distribution of X-values over a sky map is indicative not only of variability, but also to incorrect estimations of noise and instrumental variations (see Fig. 1).

### 3. Time resolved Planck fluxes, status and outlook

As the analysis of time variations in the sky is essentially background-free, our method provides a way to extract time resolved Planck fluxes down to a time resolution of a few hours. If the position of a variable point source is known, the residual 4D-map,  $R_{kj} = I_{kj} - (S*b)_{kj}$ , can be beam-deconvolved to the source position by a simple division and thus provide an estimate of the flux variation,  $\langle \Delta S(t_j) \rangle_k = R_{kj}/b_{kj}$ , where  $b_{kj}$  is a *beamfactor map* expressing the measured flux of a unit emitter at the source position in a beam centered at pixel  $k$  and time  $t_j$ . This method is used, e.g., to extract time-resolved Planck fluxes for co-eval monitoring of blazars with the F-GAMMA program (Fuhrmann *et al.* 2007, Rachen *et al.* 2015).

Our analysis tools work on unified data structures, which ensures that all methods for variability analysis and flux extraction are applicable to LFI and HFI data in the same way. Interfaces for initial 4D-mapping are in place at both the LFI and HFI DPC. We expect that all Planck frequencies will be analysed for variability, and science results prepared for publication well before the Planck Legacy release.

### References

- Fuhrmann, L., Zensus, J. A., Krichbaum, T. P., Angelakis, E., & Readhead, A. C. S. 2007, *AIPC*, 921,249
- Keihänen, E. & Reinecke, M. 2012, *A&A* 548, A110
- Górski, K. M., Hivon, E., Banday, A. J., *et al.* 2005, *ApJ*, 622, 759
- Planck Collaboration I. 2011, *A&A* 536, A1
- Planck Collaboration I. I. 2011, *A&A* 571, A2
- Rachen, J. P., Fuhrmann, L., Krichbaum, T. P., *et al.* 2015, IAU GA FP5p.43, and 28<sup>th</sup> Texas Symposium on Relativistic Astrophysics, Geneva
- Tauber, J. A., Mandolesi, N., Puget, J.-L., *et al.* 2010, *A&A*, 520, A1