

2. It was decided that pharmacy would take a leading role in ensuring the annual risk assessment forms are updated.

Following 6 months, this patient cohort was re-audited, with further results obtained.

These results showed a reduction in patients prescribed sodium valproate from 28 to 19. 37% of all patients prescribed sodium valproate had forms, but of these, only 30% were up to date. Therefore, only 11% of patients had correct annual risk acknowledgement forms in place which were up to date.

These results showed an improvement in those who had at one time had a form in place, but roughly similar compliance with availability of up to date forms.

**Conclusion.** Overall it appears that there is a real lack of consistency in ensuring the annual risk assessment forms are in place. A very low percentage of patients have the correct form in place within expiry date, despite interventions which have taken place as part of this audit. Further efforts should therefore be made by teams to ensure that these forms are in place and up to date for all women of childbearing age on valproate.

On a positive note, it may be possible to surmise that the reduction in number of patients on sodium valproate may be linked to raised awareness of the risks to women of childbearing age.

Additionally, a significant number of patients had been sent forms in the post, but had not returned them. Some of these same patients had, in the time that they should have had updated risk forms put in place, been admitted to an inpatient psychiatric hospital. It is therefore important to consider in future if these forms should be updated whilst the patients are admitted to hospital, to increase the number of patients with these forms up to date and therefore improve safety for this cohort of patients. Given that many of these patients have a diagnosis of Bipolar Affective Disorder, they may find it more challenging to return the forms whilst in the community, particularly if unwell; the ICD-11 criteria does note that patients with this condition may suffer from “distractibility, impulsive behaviour and rapid changes in mood state.” Further research could therefore be carried out to determine if utilising inpatient admissions to discuss risks of their medications would be a beneficial way to improve compliance with Annual Risk Assessment Forms.

Lastly, we know that there has been discussion around implementation of such monitoring for men as well as for women. Should this be implemented within the NICE guidelines, further audits should be carried out to determine our compliance with this.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

## Assessing the Compliance of Glucose Monitoring for Patients Prescribed Antipsychotics With Intellectual Disabilities

Dr Melissa Bremner\*

Woodland View Hospital, Irvine, United Kingdom

\*Presenting author.

doi: 10.1192/bjo.2024.547

**Aims.** To assess the compliance of glucose monitoring for patients prescribed antipsychotics in the local outpatient Learning Disability Team.

**Methods.** A review was conducted of all outpatients seen in a 6 week period during September and October 2023. Each patient was reviewed to check their diagnosis or diagnoses, the anti-psychotic medication they are currently on and if they have had the required tests done. These tests were considered, as per the NICE guidelines, to be a plasma glucose test or HbA1c test. It was checked to confirm if these tests had been carried out within the past 12 months, as per NICE.

### Results.

- 46/79 patients seen in a 6 week period at the outpatient clinic were found to be currently prescribed an antipsychotic.
- 30 of those prescribed an antipsychotic were on risperidone (65%).
- Of those prescribed any antipsychotic, 18 out of 46 had not had their glucose or HbA1c checked within the past 12 months (39%).
- This therefore demonstrates 61% of patients on an antipsychotic had appropriate glucose monitoring within the time period audited.

**Conclusion.** Monitoring glucose levels for patients on anti-psychotic medication is very important for patients with an Intellectual Disability. Patients in this cohort are known to be more likely to have diabetes and obesity than the general population. In addition, there are higher levels of inactivity and multimorbidity. It is also important to note that over-prescribing of psychotropic medication to individuals with learning disabilities, particularly antipsychotic medications such as risperidone and olanzapine, may be contributing to levels of obesity and diabetes within this population.

NICE guidelines state that for patients prescribed an antipsychotic, plasma glucose or HbA1c should be checked 3 months after commencement of treatment, and then every 12 months whilst on treatment. For olanzapine and clozapine, levels should be checked after 1 month of commencing treatment. Symptoms of hyperglycaemia should also be asked about (such as polydipsia, polyuria and increased appetite).

The results from this audit demonstrate there is definitely room for improvement in our monitoring of glucose levels for these patients. From discussing this with colleagues, it appears that a multidisciplinary approach is needed to promote this change.

Going forward, therefore, interventions should include asking the nursing staff within the outpatient team to monitor for increased appetite, polydipsia and polyuria amongst patients, especially if they are on an antipsychotic. Additionally, for any patients seen in OPC who are prescribed an antipsychotic, it should be routinely checked when they last had a glucose or HbA1c test. If this was not within the past 12 months, this should be carried out by the GP or another appropriate member of the team.

Overall, the physical health of our patients with Intellectual Disabilities is paramount. Given the nature of their Intellectual Disability and depending on the severity, it may be very challenging for them to identify any new symptoms of diabetes themselves or to report these to carers. Therefore, when prescribing antipsychotic medication, it is vital to monitor the effects of this to ensure optimal patient care utilising a multidisciplinary team approach.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.