

Concept generation techniques change patterns of brain activation during engineering design

Tripp Shealy ¹, John Gero ², Mo Hu ¹ and Julie Milovanovic ³

¹ Department of Civil and Environmental Engineering, Virginia Tech, Blacksburg, USA

² Department of Computer Science and School of Architecture, University of North Carolina at Charlotte, Charlotte, USA

³ UMR AAU-CRENAU, Graduate School of Architecture of Nantes, Nantes, France

Abstract

This paper presents the results of studying the brain activations of 30 engineering students when using three different design concept generation techniques: brainstorming, morphological analysis, and TRIZ. Changes in students' brain activation in the prefrontal cortex were measured using functional near-infrared spectroscopy. The results are based on the area under the curve analysis of oxygenated hemodynamic response as well as an assessment of functional connectivity using Pearson's correlation to compare students' cognitive brain activations using these three different ideation techniques. The results indicate that brainstorming and morphological analysis demand more cognitive activation across the prefrontal cortex (PFC) compared to TRIZ. The highest cognitive activation when brainstorming and using morphological analysis is in the right dorsolateral PFC (DLPFC) and ventrolateral PFC. These regions are associated with divergent thinking and ill-defined problem-solving. TRIZ produces more cognitive activation in the left DLPFC. This region is associated with convergent thinking and making judgments. Morphological analysis and TRIZ also enable greater coordination (i.e., synchronized activation) between brain regions. These findings offer new evidence that structured techniques like TRIZ reduce cognitive activation, change patterns of activation and increase coordination between regions in the brain.

Key words: concept generation, design neurocognition, functional near-infrared spectroscopy, prefrontal cortex

Received 06 February 2020

Revised 09 October 2020

Accepted 13 October 2020

Corresponding author

Tripp Shealy

tshealy@vt.edu

© The Author(s), 2020. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Des. Sci., vol. 6, e31

journals.cambridge.org/dsj

DOI: [10.1017/dsj.2020.30](https://doi.org/10.1017/dsj.2020.30)



1. Introduction

Engineering design is an iterative process of problem exploration, concept generation and evaluation (Cross 1989). This process of design is not linear (Lawson 2006). It is an activity that evolves through time (Dorst & Cross 2001). Arguably the most critical time is during concept generation (French 1999). The quality and quantity of concepts generated in this intermediate phase ultimately determine the outcome (Shah, Smith & Vargas-Hernandez 2003; Bryant 2005; Helm *et al.* 2016). There are numerous techniques to enhance the concept generation process (Bohm, Vucovich & Stone 2005; Jablolkow *et al.* 2015; Helm *et al.* 2016). Concept generation techniques rely on diverse procedures, classified into broad categories based on their intuitiveness (intuition and logical steps) (Shah, Smith & Vargas-Hernandez 2003), their structure (structured, partially structured or unstructured) (Gero *et al.* 2012) and the amount of motivation

required from the designer (intrinsic motivation or goal-directed motivation) (Taura & Nagai 2013; Shealy & Gero 2019).

Recently, design research has shown a growing interest in using neuroscience tools and methods to better understand the cognition of design ideation (Seitamaa-Hakkarainen *et al.* 2016; Borgianni & Maccioni 2020; Gero & Milovanovic 2020). The purpose of the research presented in this paper is to explore how three different techniques, brainstorming (Osborn 1953), morphological analysis (Allen 1962) and TRIZ (Altshuller 1984), influence design cognition in the brain. Concept generation techniques shape design outcomes through their steps and procedures, for example, by increasing or decreasing abstract reasoning, memory retrieval or uncertainty processing (Shealy & Gero 2019). These changes in cognition are observable in the patterns of activation in the brain (Alexiou, Zamenopoulos & Gilbert 2011; Hu & Shealy 2019). For instance, creative tasks rely heavily on the right prefrontal cortex (Gilbert 2010). Neuroscience offers methods to explore activation patterns in brain regions associated with critical cognitive functions for design (Liang 2017; Shealy & Gero 2019). The paper begins by providing the background for why variability in neurocognition is expected when using these concept generation techniques. This is followed by an outline of the methods used to measure cognitive activation during concept generation. Results provide evidence of significant differences in cognitive activation between the three concept generation methods studied: brainstorming, morphological analysis and TRIZ. The paper concludes with a discussion of potential explanation for these differences, and the conclusion presents opportunities for future studies.

2. Background

2.1. Concept generation

Three well-known design ideation techniques: brainstorming (Osborn 1953), morphological analysis (Allen 1962) and TRIZ (Altshuller 1984) encapsulate different characteristics in terms of ideation intuitiveness and motivation as well as technique structuredness, as Table 1 shows. Such techniques can be used either individually or in groups (Gero, Jiang & Williams 2013). Brainstorming was originally developed by Osborn (1953) as a group concept generation technique, but it can also be used by individuals in solitary situations (Harari & Graham 1975). Brainstorming is characterized as intuitive, unstructured and an inner sense-driven process (Shealy & Gero 2019). Brainstorming requires the designer to be intrinsically motivated (Shai *et al.* 2009). In practice, it is the fluid ideation of concepts. A general guideline for brainstorming is to generate as many ideas as possible and suspend evaluation until the next design phase (Daly *et al.* 2012). Contrary to our use of brainstorming, TRIZ is a logical, structured and problem-driven technique (Gero, Jiang & Williams 2013; Shah, Kulkarni & Vargas-Hernandez 2000; Shealy & Gero 2019). TRIZ requires users to decompose and analyse the problem systematically before generating new concepts. TRIZ offers engineering principles and cataloged solutions (i.e., design reference of 39 engineering parameters and 40 innovative principles).

Morphological analysis has some similar attributes to both brainstorming and TRIZ (Gero, Jiang & Williams 2013). Morphological analysis, like brainstorming, is an intuitive technique. It relies heavily on association rather than standardized

Table 1. Comparisons of concept generation techniques.

Techniques	Brainstorming	Morphological analysis	TRIZ
Intuitiveness	Intuitive	Intuitive	Logical
Motivation	Inner-sense driven	Problem-driven	Problem-driven
Structure	Unstructured	Partially structured	Structured
Steps	1	Generate as many solutions as possible and suspend evaluation	Define and decompose the problem
	2		Define the problem
	3		Search for standard engineering parameters
	4		Search for standard cataloged solutions
			Generate final solutions

engineering principles (Shah, Smith & Vargas-Hernandez 2003). Morphological analysis is also problem-driven, similar to TRIZ. In morphological analysis, a final design is predetermined through decomposition, forced association and a structured combination (Zwicky 1969).

Concept generation techniques like TRIZ and brainstorming often involve opposing cognitive structures (Gero, Jiang & Williams 2012). TRIZ tends to increase focus among designers (Gero, Jiang & Williams 2013) and can lead to a mental fixation on problem constraints (Gero 2011). This fixation occurs because TRIZ is problem-driven and follows logical steps guided by analysis, situational context and constraints (Cross 2006; Crilly 2015). Such fixation can unintentionally hinder potential creative leaps that are needed during design (Storm & Hickman 2015). In contrast to TRIZ, brainstorming enables potential creative leaps by encouraging designers to suspend evaluation and relax constraints. However, the quality of design proposals that are developed through brainstorming is often doubted because of the lack of structure and lack of intermediate evaluation in its process (Howard, Dekoninck & Culley 2010; Shah, Kulkarni & Vargas-Hernandez 2000).

Mental processes in the brain regulate the ability to generate design concepts when using TRIZ and brainstorming (Fink *et al.* 2009). An assumption about brain function during design is that information is stored in separate cortical modules that have not been previously associated (Alexiou *et al.* 2009). Composing new concepts elicits connections and communication between disparate regions of the brain (Heilman, Nadeau & Beversdorf 2003). How and where activations occur in the brain can provide new insight into concept generation (Liu, Nguyen & Zeng 2016; Sweller 1994).

2.2. Cognitive functions in the prefrontal cortex (PFC) relating to concept generation

A critical region for new connections and communication during concept generation is the prefrontal cortex (PFC) (Gibson, Folley & Park 2009, Gilbert *et al.* 2010,

Goel 2014). The PFC is the region of the brain associated with executive control functions (Schneider, Owen & Duncan 2012), attention (Dias, Robbins & Roberts 1996), working memory (Lara & Wallis 2015), planning and inhibition (Dietrich 2004). Subregions within the PFC are especially necessary for creative tasks like concept generation (Beaty *et al.* 2016; Dietrich & Kanso 2010; Dietrich 2004; Goldschmidt 2016). The right PFC plays an active role in divergent thinking (Aziz-Zadeh, Liew & Dandekar 2013; Heilman, Nadeau & Beversdorf 2003; Wu *et al.* 2015; Zmigrod, Colzato & Hommel 2015) and sustained attention (Cabeza & Nyberg 2000). Designers who display high originality in solution generation exhibit strong synchronization within the right PFC (Fink *et al.* 2009). The left PFC plays a more active role when supporting rule-based design, goal-directed planning (Aziz-Zadeh, Liew & Dandekar 2013) and making analytic judgments (Hoefft *et al.* 2007; Gabora 2010; Luft *et al.* 2017). The left PFC also plays a critical role in solving math problems (Poldrack *et al.* 1999).

The left and right dorsolateral prefrontal cortex (DLPFC) is bilaterally active when performing creativity tasks that require new associations and evaluations (Funahashi 2017). For instance, activation in the left DLPFC decreases (Tachibana *et al.* 2019) and activation increases in the right DLPFC during improvisation (De Dreu *et al.* 2012; Kleibeuker *et al.* 2013). The medial PFC (mPFC) and ventrolateral PFC (VLPFC), are also involved in creative design tasks. The function of the mPFC is to learn associations and is observed to play a role in the retrieval of 'remote' memories (Euston, Gruber, & McNaughton 2012). Increased activation in the mPFC is associated with improved ability to simulate future imaginative events (Meyer *et al.* 2019). The VLPFC is critical for combining existing information into new ideas (Dietrich 2004; Wu *et al.* 2015). The ability to detect similarity between items activates the right VLPFC (Garcin *et al.* 2012).

2.3. Identifying coactivation of PFC subregions with brain network

One approach to understand the relationship between patterns of activation in subregions of the PFC is through neural networks. Neural networks are used to describe how and where connections are made spatially between brain regions, and this is used to develop frameworks about brain processing, the activation level of these regions and patterns of coactivation among regions during design (Martindale 1995). For example, distinct patterns of activation in the right parietal and right prefrontal cortex occurred among females during spatial-cognition tasks and left hippocampus in males (Grön *et al.* 2000). The difference in activation patterns by gender is expressed by their neural network connections between brain regions (Grön *et al.* 2000).

Identifying interconnected brain regions that are central for each concept generation technique can also provide evidence about what engineers are doing and thinking during design (Alexiou, Zamenopoulos & Gilbert 2011). For instance, TRIZ requires cognitive flexibility to switch between evaluating design principles and imagining the use of these principles with given problem constraints (Savransky 2000). Cognitive flexibility is observed in the brain by higher oscillation between left and right hemisphere dominance in the brain compared to brainstorming (Shealy, Hu & Gero 2018).

A new concept might be missed if requisite brain regions are not sufficiently engaged, and this is also observable in patterns of activation described by neural

network connections (Grabner *et al.* 2009). For example, an increase in the connections associated with the right DLPFC corresponds to an increase in the number of solutions generated (Hu 2018). Performance in the ability to develop new associations when concept mapping is also observable in network connections. Concept maps can reduce the need for coordination in the brain because of a reduction in demand from working memory and an increase in activation in the region of the brain associated with divergent thinking (Hu *et al.* 2019).

2.4. Neuroimaging techniques to measure cognitive activation

Several neuroimaging techniques are available to quantify neurocognitive activation in the brain during concept generation and build models of neural networks. These methods include electroencephalograms (EEGs), functional magnetic resonance imaging (fMRI) and function near-infrared spectroscopy (fNIRS). EEG and fMRI are widely used to study creativity (see Pidgeon *et al.* 2016 for a review) and design studies using such tools focussed on diverse topics such as comparing the neurocognition of mechanical engineers and architects (Vieira *et al.* 2019a; 2019b), evaluating mental effort and mental stress while designing (Nguyen & Zeng 2014), the influence of design problem constraints on workload and convergent and divergent thinking (Liu *et al.* 2018), the difference between design and problem-solving in the neurological basis (Alexiou *et al.* 2009), or the role of dorsolateral prefrontal cortex in ill-structured design cognition (Gilbert *et al.* 2010). fNIRS is a more recently developed neuroimaging technique. It has gained popularity because of its usability in naturalistic environment and resilience to motion artefacts (Balardin *et al.* 2017; Brockington *et al.* 2018).

EEG has a high temporal resolution (i.e., ability to detect quick changes on the order of milliseconds), mobility and a relatively low initial purchase price (Hu & Shealy 2018). EEG, however, is limited in spatial resolution (i.e., ability to detect where the change in cognitive activation occurs) because the electrical activity measured by EEG goes through multiple layers in the brain and is a mixture of signals from underlying brain sources. The ability to pinpoint specific brain regions with EEG is a challenge (Burle *et al.* 2015) and is limited to macro and even hemispherical scales. Recent advances in EEG technology have increased the spatial resolution considerably. In contrast to EEG, fMRI has high spatial resolution with the ability to display cognitive activation in the whole brain. fMRI measures the changes in blood oxygenation level, which is linked to cognitive activity (Gramann *et al.* 2014). The temporal resolution of fMRI is on the order of seconds due to the blood flow change over time and the time needed for net magnetization recovery before the next sampling (Eysenck & Keane 2015). Data collection with an fMRI machine requires participants to remain still and lay down while partially enclosed inside the fMRI scanner and this can be constraining. While studying design with fMRI, a solution is for participants to verbalize their design solutions and subsequently sketch them once out of the fMRI scanner (Hay *et al.* 2019).

Considering the limited spatial resolution of EEG and less naturalistic experiment environment of fMRI, the study presented in this paper adopted the use of fNIRS. It has relatively high spatial and temporal resolution and is portable. Participants can operate a computer or perform a task in an upright sitting position, similar to EEG. fNIRS has a good spatial resolution compared to EEG but low spatial resolution compared to fMRI. fNIRS does not measure cognitive

activity directly rather it measures metabolic demands (oxygen consumption) of active neurons (Herold *et al.* 2018). fNIRS is worn as a cap where light is emitted from sources at specific wavelengths (between 700 and 900 nm) into the scalp. The light scatters before reflecting back to light receivers. The oxy-hemoglobin (oxy-Hb) and deoxy-hemoglobin (deoxy-Hb) absorb more light than water and other tissue in the brain. The change in the difference between the emitted light and reflected light is used to calculate the change in oxygenated blood using a modified Beer–Lambert law. The oxy-Hb and deoxy-Hb are inversely related. Typically, only oxy-Hb is reported because of its relatively higher amplitudes and sensitivity to cognitive activities (Chu *et al.* 2008; Cazzell *et al.* 2012; Zhang *et al.* 2017; Hu & Shealy 2019).

A drawback of fNIRS is the limited power of light emitter, which makes it unable to capture subcortical activation in the brain, unlike fMRI. However, areas relevant for design neurocognition, such as the PFC, associated with executive function and working memory, are sufficiently accessible with fNIRS (Fuster 1988). For example, fNIRS can adequately capture the ability to think in systems (Hu *et al.* 2019) and make decisions (Hu & Shealy 2019; Shealy & Hu 2017).

The research reported in this paper aimed to assess how the attributes associated with the three concept generation techniques change how information is cognitively processed and influence the dominant use of specific regions in the brain. The use of fNIRS enables measuring neurocognitive activation during design. It acts as a proxy for neurocognition by measuring change oxy-Hb (Herold *et al.* 2018). Change in oxy-Hb provides evidence of the changes in cognitive demand patterns and functional coordination (e.g., abstract reasoning and evaluation) when designers generate concepts and how patterns of neurocognition and neurocoordination vary between techniques.

3. Research questions

The study described in this paper aimed to assess how brainstorming, morphological analysis and TRIZ changes how specific brain regions are activated in the PFC. The specific research questions are:

1. What is the effect of brainstorming, morphological analysis and TRIZ on cognitive activation in the prefrontal cortex?
2. What regions within the prefrontal cortex are most central during concept generation when using brainstorming, morphological analysis and TRIZ?
3. How does cognitive coordination across regions in the prefrontal cortex change over time when using brainstorming, morphological analysis and TRIZ?

4. Methods

4.1. Experimental design

Thirty graduate engineering students (all right-handed, 22–26 years old, 10 females and 20 males) were recruited to participate in the study. The procedures followed for this study were approved by the Institutional Review Board. There was no incentive provided for participation. Recruitment occurred through multiple graduate engineering courses at the same institution. All participants reported prior course work in engineering design and were first-year graduate students. Participants completed all three concept generation tasks individually using a

different technique (brainstorming, morphological analysis and TRIZ) for each task. None of the participants indicated they had formal training with morphological analysis or TRIZ. Pretask training was provided to introduce the three techniques to participants. The pretask training included verbally explaining the steps of both morphological analysis and TRIZ. Participants were allowed to review the written instructions provided with each design technique. The experiment began with a 15-second baseline period. This baseline asked participants to keep their mind in a rested state. Participants then received one of three engineering design tasks and completed the task at their own pace using one of the three techniques as instructed. The sequence of techniques and design tasks were assigned randomly to each participant. Each participant completed all three design tasks using one of the three techniques. The 15-second baseline period commenced before each design task.

The instructions for brainstorming were for participants to generate solutions for the design task and suspend evaluation of their design. Participants were not provided any additional tool or aid during the brainstorming task. The instructions for morphological analysis were to define and decompose the problem, generate multiple subsolutions and then develop a solution. The instructions for TRIZ were to define the problem, review standard engineering parameters that fit this problem, compare these parameters with cataloged solutions and then generate a solution (see [Table 1](#)). The steps for the design process follow previously developed methods (see Gero, Jiang & Williams [2012](#); Gero, Jiang & Williams [2013](#) for more details).

The design tasks were not discipline specific and previously demonstrated to require similar cognitive processes to generate a solution (Gero, Jiang & Williams [2013](#)). In one of the design tasks, participants were instructed to design a device to assist the elderly with raising and lowering windows. Another design task required participants to design an alarm clock for the hearing impaired. The final design task asked participants to design a kitchen measuring tool for the blind. Participants were instructed to sketch on paper to illustrate their design solutions. Participants were instructed to raise their hand when they were done developing their final solutions and data collection with fNIRS would stop. Observations of participants during the tasks provided some indication about whether participants continued to make progress during the design task. Any participants that appeared to stop during the design or disengage were noted, though this was not an issue for any of the 30 participants. The average time to generate a solution when brainstorming, using morphological analysis and TRIZ, was 7.53 min (SD = 3.25 min), 11.02 min (SD = 4.70 min) and 13.34 min (SD = 5.03 min), respectively. Most participants generated one to three design solutions or subsolutions when using brainstorming, morphological analysis and TRIZ.

Participants were outfitted with the fNIRS cap from LIGHTNIRS fNIRS system (Shimadzu Co., Kyoto, Japan) with a sampling frequency of 4.44 Hz. LIGHTNIRS uses a three-wavelength absorbance calculation (780, 805 and 830 nm) to record a change in participants' oxygenated hemoglobin. Change in participants' oxygenated hemoglobin is an indicator of cognitive activation in their PFC as they generated a solution to each design task. The sensor placement on the fNIRS cap is shown in [Figure 1](#). A total of 16 sensors (8 emitters and 8 detectors) were located using the 10/20 international systems and formed a total of 22 channels. A channel is the combination of a light source and a nearby light receiver. This is indicated in




Figure 1. A participant with fNIRS cap and sensor configuration.

Figure 1b with light source and light receiver numbers. It captures the change in oxygenated cortical blood in the brain. These channels cover multiple subregions in the PFC, including dorsolateral prefrontal cortex (DLPFC: channel 1, 2, 3, 9 and 10 in the right hemisphere and channel 5, 6, 7, 13 and 14 in the left hemisphere), ventrolateral prefrontal cortex (VLPFC: channel 16 and 17 in the right hemisphere and channel 21 and 22 in the left hemisphere), orbitofrontal cortex (OFC: channel 18 in the right hemisphere and channel 20 in the left hemisphere) and medial prefrontal cortex (mPFC: channel 4, 11, 12 and 19) in both hemispheres.

4.2. Data analysis

Three out of the 30 participants were removed from the analysis because of a weak signal during the experiment. In this study, we compare the idea generation phases for each technique. Therefore, for brainstorming, we considered the entire session, for morphological analysis, we analysed the third step of this technique of multiple subsolutions generation (see Table 1) and for TRIZ, and we analysed the fourth step that focusses on generating final solutions (see Table 1). This way, we are able to compare the ideation phase of each technique.

fNIRS raw data for the remaining 27 subjects were processed using a bandpass filter (frequency ranging between 0.01 and 0.1 Hz, third-order Butterworth filter) to remove high-frequency instrumental and low-frequency psychological noise (Huppert *et al.* 2009). To reduce motion artifacts, participants were instructed to keep their head motion to a minimum, additionally an independent component analysis with a coefficient of spatial uniformity of 0.5 was applied to remove motion artifacts. The steps of noise and motion artifacts removal are critical to avoid false discovery in brain network and connectivity analysis (Santosa *et al.* 2017). The parameters in data processing are based on prior research (Sato, Hokari & Wade 2011; Naseer & Hong 2015). The filtering process was conducted using Shimadzu fNIRS software, and the following analysis was conducted using Python (NetworkX package was used for the network analysis). Only oxygenated hemoglobin (oxy-Hb) in the filtered data is reported in the results because oxy-Hb generally has a higher amplitude and is more sensitive to cognitive activities than deoxygenated hemoglobin (deoxy-Hb) (Chu *et al.* 2008; Cazzell *et al.* 2012; Zhang *et al.* 2017; Hu & Shealy 2019). Then, the baseline correction is applied in which the




Figure 2. Area under the curve and mean value of Oxy-Hb.

mean oxy-Hb during the baseline rest period was subtracted from the oxy-Hb during the tasks for each channel.

To answer the first research question, two methods were used to measure neurocognitive activation in the prefrontal cortex when using brainstorming, morphological analysis and TRIZ. First, the positive area under the curve (AUC), as illustrated by the shading in [Figure 2](#), was calculated for each participant when using each design technique. Blood oxygenation level dependent-local field potential (BOLD-LFP) coupling model suggests that positive BOLD responses (i.e., increased oxy-Hb) correspond to actively actuated increase in blood flow in support of neural activity (Ekstrom 2010; Bartra, McGuire, & Kable 2013). Therefore, the cumulated amplitudes of oxy-Hb (i.e., AUC) were used as an indicator of cognitive load, which was used in prior literature to evaluate cognitive load (Manfredini *et al.* 2009; Agbangla, Audiffren & Albinet 2017; Suzuki *et al.* 2018). In addition to the positive AUC, the absolute AUC was also calculated. Similar findings between the positive AUC and absolute AUC were observed. The positive AUC is a better predictor to classify high and low mental workload in prior brain-computer interface studies (Verdière *et al.* 2018). So, the positive AUC results are reported. The positive AUC for all channels was used for comparison between the concept generation techniques and is described in the results as the overall cognitive load in the PFC (Manfredini *et al.* 2009; Agbangla, Audiffren & Albinet 2017; Suzuki *et al.* 2018).

AUC was also used to compare hemisphere asymmetry between the left and right PFC (Toga & Thompson 2003; Runco 2014). The AUC of 10 channels in the right PFC and 10 channels in the left PFC were averaged respectively to calculate a proxy for cognitive load in the right and left hemispheres. Analysis of variance (ANOVA) was used to measure the statistical difference in the AUC across the PFC and the left and right PFC for each concept generation technique. Significance was defined as $p < 0.05$. The effect size for the significant difference was measured by η^2 (Eta squared) for ANOVA. The difference is regarded as large when η^2 is greater than 0.138 (Cohen 1977). We performed a normality check using the Shapiro-Wilk test before the ANOVA analysis. The purpose of the normality check was to confirm the data were normally distributed.

The second measure for cognitive activation was the mean value of oxy-Hb illustrated by the dotted line in Figure 2. We created one mean oxy-Hb for each design technique. To do this, we used a fractioning technique based on the function-behaviour-structure (FBS) design ontology framework (Gero 2010; Gero, Jiang & Williams 2013). The purpose of this fractioning was to normalize the concept generation sessions over time. This normalization was necessary because each concept generation phase had a different length of time. The fractioning technique divided the design session for each design task for each participant into 20 equal and nonoverlapping segments or ventiles. Participants' mean oxy-Hb was then calculated for each ventile. The length of ventiles varied for each participant because the time they spent during concept generation varied. All of the participants' ventiles were then averaged together to create an average oxy-Hb for each of the design techniques. The use of 20 segments for the average oxy-Hb follows prior design cognition studies (e.g., EEG studies and design protocol studies) (Gero, Jiang & Williams 2013; Jiang *et al.* 2014; Kan & Gero 2017; Milovanovic & Gero 2018; Shealy & Gero 2019). ANOVA was then used to measure the difference in the patterns of oxy-Hb in the PFC for the 20 ventiles (including left and right DLPFC, VLPFC and mPFC). Significance was defined as $p < 0.05$.

To answer research question 2, graph theory (Wijk, Stam & Daffertshofer 2010) was used to understand what regions within the prefrontal cortex are most central and the coordination required between brain regions during concept generation (Bullmore & Sporns 2009; De Vico Fallani *et al.* 2014). Pearson's correlation matrices were developed using the change of oxy-Hb in all channels following the common steps in prior studies (Achard & Bullmore 2007; Bullmore & Sporns 2009) during each design task for each participant. Correlation matrices were averaged across participants when using the same design technique. A range of plausible global threshold coefficients (incrementally from 0.6 to 0.7) as used in prior studies (Achard & Bullmore 2007; Bullmore & Sporns 2009; Bressler & Menon 2010) were considered as connective functions (De Vico Fallani *et al.* 2014; Fornito, Zalesky & Bullmore 2016; Bassett & Sporns 2017). Correlations higher than the threshold coefficients indicate a correlative and potentially functional relationship between synchronized activation of different brain regions. Links were drawn between channels (called nodes in a network) when the correlation coefficient was higher than the threshold. These steps are illustrated in Figure 3. All the links (i.e., connections between channels) and nodes (i.e., 22 channels) form a network. For each ventile during brainstorming, morphological analysis and TRIZ, a PFC




Figure 3. Brain networks and metrics

functional network was developed based on the participants average Pearson correlation matrix following the steps illustrated in Figure 3.

The centrality and network density were then calculated to provide descriptive measures of the network. Node centrality describes the nodes with the most edges in the network. Central nodes are critical to efficient communication for task completion (Bullmore & Sporns 2009; Fornito, Zalesky & Bullmore 2016). The density of connections was used to answer research question 3. Network density is the proportion of the number of actual connections to the number of possible connections in a network. It provides an estimate of cognitive coordination within the network (Achard & Bullmore 2007). A low network density means low coordination between brain regions. Then the average network density in each ventile for each technique among all participants was calculated. Network densities were compared between the three techniques using ANOVA followed by posthoc analysis using paired t-tests to compare network density between design task for each participant.

5. Results

5.1. TRIZ demands significantly less cognitive load in the prefrontal cortex compared to brainstorming and morphological analysis

The neurocognitive activations when generating new concepts through brainstorming, morphological analysis and TRIZ are significantly different ($F(2,57) = 29.5, p < 0.001, \eta^2 = 0.509$) with a large effect size. The positive area under the curve (AUC) of oxy-Hb in the PFC is lower when using TRIZ compared to brainstorming ($t = 4.68, p < 0.001$) and morphological analysis ($t = 7.62, p < 0.001$). Morphological analysis elicited significantly more AUC in the PFC than brainstorming ($t = 2.94, p = 0.013$). AUC is used as one indicator of cognitive load associated with working memory, cognitive flexibility and reasoning. We observed that TRIZ reduces the cognitive load (i.e., the positive AUC) required in the PFC compared to brainstorming and morphological analysis.

These results are consistent when isolating the right PFC. TRIZ reduces the cognitive load ($F(2,57) = 36.6, p < 0.001, \eta^2 = 0.465$) required in the right PFC compared to brainstorming ($t = 7.38, p < 0.001$) and morphological analysis ($t = 7.43, p < 0.001$). The effect size is large. TRIZ also demands significantly ($F(2,57) = 25.5, p < 0.001, \eta^2 = 0.472$) less cognitive load in the left hemisphere when generating concepts compared to when using brainstorming ($t = 2.33, p = 0.025$) and morphological analysis ($t = 6.86, p < 0.001$). Morphological analysis elicited significantly more cognitive load in the left PFC than brainstorming ($t = 5.13, p < 0.001$). To summarize these results, for our participants, TRIZ requires significantly less cognitive load than morphological analysis and brainstorming in the right and left PFC. Morphological analysis demands a higher cognitive load in the left hemisphere compared to brainstorming and TRIZ. These results are illustrated in Figure 4.

5.2. Brainstorming, morphological analysis and TRIZ produce significantly different patterns of cognitive activation over time

Consistent with the area under the curve, mean oxy-Hb over time, which is a proxy for cognitive activation, in the right DLPFC ($F(2,57) = 58.9, p < 0.001, \eta^2 = 0.674$) and right VLPFC ($F(2,57) = 9.78, p < 0.001, \eta^2 = 0.255$) is significantly less when




Figure 4. Difference in area under the oxy-Hb after baseline correction when using brainstorming, morphological analysis and TRIZ; (a) Average area under the curve (AUC) in the left and right prefrontal cortex (PFC); (b) AUC in the left PFC; (c) AUC in the right PFC.




Figure 5. Differences in patterns of cognitive activation in the right dorsolateral prefrontal cortex (a) and right ventrolateral prefrontal cortex (b) when brainstorming, using morphological analysis and TRIZ.

using TRIZ compared to brainstorming and morphological analysis. Mean oxy-Hb is significantly different in the right DLPFC between TRIZ and brainstorming ($t = 10.39, p < 0.001$) and morphological analysis ($t = 7.91, p < 0.001$). Patterns of cognitive activation are similar when using brainstorming and morphological analysis with no significant difference. TRIZ also demands significantly less cognitive activation in the right VLPFC compared to brainstorming ($t = 4.28, p < 0.001$) and morphological analysis ($t = 3.09, p = 0.008$). Figure 5 depicts the patterns of cognitive activation in both the right DLPFC and right VLPFC. Both TRIZ and brainstorming demand more cognitive activation early in the concept generation process, but this activation declines more quickly with TRIZ.




Figure 6. Differences in patterns of cognitive activation in the left dorsolateral prefrontal cortex when brainstorming, using morphological analysis and TRIZ.

Morphological analysis tends to demand more cognitive activation in the middle of the concept generation process with two distinct peaks around ventiles 6 and 12.

Significant differences ($F(2,57) = 10.70$, $p = 0.003$, $\eta^2 = 0.181$) in patterns of cognitive activation when using brainstorming, morphological analysis and TRIZ are also observed in the left DLPFC. TRIZ ($t = 3.43$, $p = 0.003$) and morphological analysis ($t = 2.51$, $p = 0.039$) demand more cognitive activation compared to brainstorming with a large effect size ($\eta^2 > 0.138$). TRIZ and morphological analysis elicit similar patterns of cognitive activation and produce multiple peaks of cognitive activation in the left DLPFC that is higher in amplitude than brainstorming. Some activation is observed at the beginning and end during brainstorming, but the amplitude of activation is lower compared to TRIZ and morphological analysis, illustrated in Figure 6.

A significant and large ($F(2,57) = 20.7$, $p < 0.001$, $\eta^2 = 0.420$) difference is also observed in the medial PFC (mPFC) (channels 11 and 19). Brainstorming ($t = 6.35$, $p < 0.001$) and morphological analysis ($t = 4.04$, $p < 0.001$) demand more cognitive activation over time in the mPFC than TRIZ, and the difference is large ($\eta^2 > 0.138$). Brainstorming demands more cognitive activation both at the beginning and end of the concept generation process. Neurocognitive activation gradually increases when using morphological analysis for the first 15 ventiles. TRIZ demands more neurocognitive activation early and late in the concept generation process, but the amplitude of activation is less than both brainstorming and morphological analysis, illustrated in Figure 7.

5.3. Node centrality varies by hemisphere between brainstorming, morphological analysis and TRIZ

Brain network analysis for the entire length of the design ideation phase suggests that node centrality varies when using brainstorming, morphological analysis and TRIZ. A sequence of increasing threshold coefficients within the range of 0.6–0.7 was used to measure node centrality. The channels with the highest centrality (average under all thresholds) and their associated regions are shown in Table 2. The network graphs in Table 2 illustrate the brain network with a global threshold of 0.6 and 0.7 when using brainstorming, morphological analysis and TRIZ.





Figure 7. Difference in patterns of cognitive activation (mean value of Oxy-Hb) in the subregions of medial prefrontal cortex among techniques.

Table 2. Network graphs and centrality when concept generation.

Technique	Network graph, Threshold = 0.6	Network graph, Threshold = 0.7	Channel: central regions
Brainstorming			Channel 10: 0.351 (Right DLPFC) Channel 3: 0.253 (Right DLPFC)
Morphological			Channel 11: 0.310 (Right DLPFC) Channel 14: 0.230 (Left DLPFC)
TRIZ			Channel 12: 0.344 (Left DLPFC) Channel 11: 0.307 (Right DLPFC) Channel 10: 0.270 (Right DLPFC) Channel 4: 0.264 (Medial PFC)

Abbreviations: DLPFC, dorsolateral prefrontal cortex; PFC, prefrontal cortex.




Figure 8. Network density change over time during concept generation (correlation threshold equals 0.7).

When brainstorming, the most central node is in the right DLPFC. When using morphological analysis, the most central node is in both the right and left DLPFC. When using TRIZ, the most central nodes are in the right DLPFC, left DLPFC and medial PFC. TRIZ also elicits the most network connections compared to morphological analysis and brainstorming. Morphological analysis elicits more network connections than brainstorming. Network connections are one proxy for coordination between brain regions.

5.4. Coordination between brain regions increases when using morphological analysis and TRIZ compared to brainstorming

The network density was calculated for each ventile when using brainstorming, morphological analysis and TRIZ. The purpose of this network density was to understand the coordination between brain regions over time. Figure 8 shows the change in density for each ventile. There are significant $F(2,57) = 8.86, p < 0.001, \eta^2 = 0.237$ differences in the network density when using brainstorming, morphological analysis and TRIZ. The density when brainstorming is significantly lower than morphological analysis (e.g., $t = -2.71, p = 0.013$ when threshold = 0.7) and TRIZ (e.g., $t = -4.76, p = 0.001$ when threshold = 0.7). Morphological analysis and TRIZ have no significant difference in network density. TRIZ and morphological analysis significantly increase the brain regions that are in coordination during concept generation compared to brainstorming, especially in the early and middle phase of concept generation as Figure 8 shows.

6. Discussion

These results offer empirical evidence about the neurocognitive differences when using brainstorming, morphological analysis and TRIZ. The results relate to the structuredness of each technique. Fundamental cognitive functions of the PFC include working memory, cognitive flexibility and reasoning (Lara & Wallis 2015;

Funahashi 2017). The cumulative cognitive activation (described in the results as the positive area under the curve for oxy-Hb) in the PFC is a proxy for cognitive load associated with the cognitive functions in this region. The results indicate that the use of TRIZ demands less cognitive load in the PFC than brainstorming and morphological analysis. This trend also appears in the right and left PFC. This trend is consistent with prior research that says TRIZ is likely to occupy less space in students' short-term memory based on self-report surveys and student reflections (Belski 2011; Belski & Belski 2015). A reason why TRIZ demands less cognitive load might be that the structuredness of TRIZ offers strong cues and an organized information retrieval process between short-term and long-term memory systems. With reference to the 39 Engineering Parameters and 40 Innovative Principles in TRIZ, students break down the problem, focus on a single principle at a time and attend to one possible solution before moving to the next parameter and principle. This process of shifting attention between principles and solution reduces cognitive complexity in design. This alleviation in cognitive load seems to align with human cognitive structures as cognitive load theory suggests (Jong 2010). The lower demand in cognitive load during TRIZ should be further explored. None of the students in this study were familiar with TRIZ. Considering the learning curve for TRIZ and possible higher levels of familiarity with brainstorming, this significant alleviation in cognitive load when using TRIZ is promising and may become even more pronounced with additional practice using TRIZ.

Conversely, the lack of cues while brainstorming might result in less focussed attention. When brainstorming, students in our cohort had significantly higher cognitive load in their PFC. The excess of information or distractions from other thoughts could be the result of the un-structuredness of this technique (Kohn & Smith 2011). Cognitive resources are limited in short-term memory (Artino 2008). Brainstorming appears to consume more of these resources (Kirschner 2002; Santanen, Briggs & de Vreede 2000), and this is consistent with prior results in the literature (Gabora 2010; Lara & Wallis 2015).

Using morphological analysis also appears to result in a higher cognitive load than TRIZ. Morphological analysis follows a process of breaking down the problem and then concept association, which can stimulate more concepts than brainstorming (Keong *et al.* 2012). However, without any engineering parameters or principles like in TRIZ, each step of morphological analysis requires intuitive thinking, likely demanding more cognitive resources. To summarize these findings, the logical rule-based technique of TRIZ provides a design tool that reduces cognitive load compared to the intuitive techniques of brainstorming and morphological analysis.

The second finding is that each technique relies on specific subregions of the PFC. Previous studies about creative tasks find limited evidence of differential activation between hemispheres and subregions (Colombo *et al.* 2015). Brainstorming and morphological analysis demand more cognitive activation (described as the mean value of oxy-Hb) in the right lateral PFC compared to TRIZ. The right lateral PFC (including DLPFC and VLPFC) is generally associated with divergent thinking (Aziz-Zadeh, Liew & Dandekar 2013; Wu *et al.* 2015; Zmigrod, Colzato & Hommel 2015) and maintaining divergent ideas with sustained attention (Cabeza & Nyberg 2000). Intuitive ideas that suddenly come to mind are associated with increased activation in the right DLPFC (Pisapia *et al.* 2016). The right DLPFC is a critical region for ill-structured design cognition (Gilbert *et al.* 2010).

The right VLPFC plays a critical role related to hypotheses generation and maintenance of divergent thinking (Goel & Vartanian 2005). A possible explanation for the higher activation in the right DLPFC and right VLPFC when using brainstorming and morphological analysis compared to TRIZ is that students tend to continually rely on divergent thinking during brainstorming and morphological analysis to generate multiple new, unconnected concepts. This reliance on divergent thinking appears to lead to higher sustained activation in the right lateral PFC to maintain these isolated small chunks of information in the working memory (Gilbert *et al.* 2010).

Another possible explanation for the higher activation in the right DLPFC when using brainstorming and morphological analysis is that the problem appears to be more ill-defined for brainstorming and morphological analysis. A design study found that the right DLPFC showed significantly higher activation in ill-structured problems than well-structured problems (Gilbert *et al.* 2010). Brainstorming begins with a random and intuitive exploration of the solution space without explicit identification of the design problem, and morphological analysis provides no parameters or principles for designers to formulate a problem like TRIZ. This explanation seems consistent with prior findings that reasoning about the design problem is increased when applying TRIZ compared to brainstorming and morphological analysis (Gero, Jiang & Williams 2013).

Patterns of high cognitive activation (described in the results as the mean oxy-HB) in the right lateral PFC occur at the beginning of concept generation when using TRIZ. A possible explanation is students might think divergently to generate many ideas, but the pattern of neurocognitive activation shifts from the right DLPFC to the left DLPFC later in the concept generation process when using TRIZ. The left DLPFC is generally associated with making judgments (Birdi, Leach & Magadley 2012) and fixation (Cross 2006). The left DLPFC is also associated with controlling convergent judgments about whether ideas generated in the right hemisphere meet constraints (Luft *et al.* 2017). This region also shows more activation in goal-directed planning of novel solutions (Aziz-Zadeh, Liew & Dandekar 2013). The higher activation in the left DLPFC when using TRIZ compared to brainstorming and morphological analysis might indicate that students reserve cognitive attention to evaluate concepts by applying filters and affirm solutions to satisfy the constraints or meet the design goals. This shift from right to left DLPFC enables cognitive flexibility and might lead to increased attention (Goldschmidt 2016), which seems to support the claim that TRIZ can increase attention (Gero, Jiang & Williams 2013).

In contrast, when using brainstorming and morphological analysis, more cognitive resources are allocated to the right DLPFC. Possibly, maintaining divergent thinking in the right hemisphere means fewer resources are available for convergent thinking and evaluating concepts in the left DLPFC. Of course, this result might not be surprising since the general instruction for brainstorming is to suspend or delay judgments when generating solutions (Keong *et al.* 2012). For morphological analysis, students might not have had adequate cognitive resources for concept evaluation allocated to the left DLPFC, which is suggested by the lower activation in this region.

Higher cognitive activation was also observed in the medial PFC (mPFC) when using brainstorming. The function of the mPFC is to learn associations and is observed to play a critical role in the retrieval of « remote » memories (Euston, Gruber, & McNaughton 2012). The higher activation in this region when

brainstorming might suggest more cognitive resources are required to make associations between divergent ideas or linking known concepts with new ones. In the case of morphological analysis, students decomposed the problem based on functions, so the association processing could seem more manageable and require less activation in the mPFC than brainstorming. The fewest cognitive resources were required when using TRIZ. Similar to morphological analysis, this logical process relies on decomposition and analysis.

In addition to the changes in cognitive load and patterns of activation in subregions, the brain network analysis in our dataset revealed potential connections between the structuredness of each concept generation technique and the central regions for cognitive coordination. The right DLPFC is the most central region needed for communication across brain regions during brainstorming. The right and left DLPFC are the two most central regions for communication across the brain during morphological analysis, and the right and left DLPFC and the medial PFC are the most central for communication across brain regions during TRIZ. The same regions were also detected with high centrality for concept generation in a prior study investigating design cognition (Shealy, Hu & Gero 2018).

The common brain region with high centrality when using all three techniques is the right DLPFC. The right DLPFC plays a crucial role in efficient communication (i.e., correlation to other nodes) during concept generation. This finding is consistent with previous research, which finds coordination in the right DLPFC is crucial to design cognition (Gilbert *et al.* 2010). The differences found in this study, compared to previous studies, is the cognitive correlation (described in the results as the network density) across the PFC is higher for TRIZ and morphological analysis than brainstorming. In other words, using the problem-driven approaches that require decomposition and analysis activate more correlation, which is a proxy for communication across regions in the brain (Achard & Bullmore 2007; Bullmore & Sporns 2009). This might be because these techniques direct more reasoning about the problem, binding of different knowledge sets and information retrieval from long-term memory (Heilman, Nadeau & Beversdorf 2003). Another possible explanation is the relative unfamiliarity with TRIZ among participants, which resulted in higher brain network communication compared to brainstorming.

The results presented in this paper provide new insights to better understand the relationship between concept generation techniques and cognitive processes through the analysis of neurocognitive activation. Brainstorming, morphological analysis and TRIZ change engineering students' neurocognitive behaviour. There are several limitations to this study that are worth mentioning. fNIRS data only include the change of oxygenated hemoglobin in the PFC. Other brain regions (e.g., parietal cortex) might also contribute to creative design cognition. This limit is characteristic in all neuroimaging studies that do not capture whole-brain activation (Ayaz *et al.* 2011; Cazzell *et al.* 2012). Another limitation is that this study focussed on neurocognitive differences between three distinct design techniques and did not include a comparison of the outcomes among engineering students. Future research could explore neurocognitive differences among students who produce more or less novel design solutions. The 27-person sample size is another limitation (Schönbrodt & Perugini 2013), although the number of participants does meet the average sample size of 27 in similar studies (Hu & Shealy 2019). Future research should replicate the results with a larger sample size (Shrout & Rodgers 2018). Additionally, the evenly fractioned and averaging technique focussed

on the group-level analysis of design dynamics while ignoring individual differences. Using a sliding window instead of a nonoverlapping window provides a higher granularity and more continuous data (Allen *et al.* 2014; Zhang & Zhu 2020). A sliding window can better capture the temporal dynamics of cognitive activation among individuals and remove the assumption that participants follow a similar path of cognitive activation (Allen *et al.* 2014; Zhang & Zhu 2020). Rather than 20 segments, a sliding window may include 300 segments. The downside of a sliding window is handling each participant's data individually without averaging them together over time. This creates a more complex data set with greater challenges in comparing between and within subjects. Future research can also begin to explore the neurocognitive effects of other design instruments. Many techniques exist for design, and variations within tool use are also rampant. For example, many different TRIZ tools are available (Separation Principles, Su-Fields, Standards, ARIZ). These tools can lead to varying outcomes (Ilevbare, Probert & Phaal 2013; Spreafico & Russo 2016) and likely lead to varying effects on patterns of neurocognition. In addition, future research can explore the differences between novice and experts during design. Previous neurocognition studies suggest differences in mental abilities based on experience are observable with fNIRS (Harrison *et al.* 2013).

7. Conclusion

The neuroimaging methods adopted in this study explored how concept generation techniques influence neurocognition during design. Significant differences are observed in cognitive activation when using brainstorming, morphological analysis and TRIZ. Brainstorming and morphological analysis induce more cognitive load across the PFC compared to TRIZ. Higher cognitive activation associated with divergent thinking and ill-defined problem-solving is observed in the right DLPFC and VLPFC when using brainstorming and morphological analysis. TRIZ demands more cognitive activation in the left dorsolateral PFC. This region is associated with controlling judgments and convergent thinking.

Centrality and correlation between regions in the PFC also varied with each technique. The right DLPFC plays a central role in network analysis across brain regions when using all three techniques. The left DLPFC also plays a central role in network analysis across brain regions when morphological analysis and TRIZ are used, and the mPFC also plays a role when using TRIZ. Morphological analysis and TRIZ significantly increase the number of brain regions that correlate during concept generation.

These multiple analyses indicate that TRIZ, compared to brainstorming and morphological analysis, increases correlation, which is a proxy for coordination, between brain regions, and decreases the cognitive load during concept generation. This insight about the neurocognitive benefits of using TRIZ offers new supporting evidence for the use of structured and goal-direct concept generation techniques. It motivates the development of new techniques and offers a more in-depth explanation about how these techniques inform creative thought and behaviour. Future research should explore the correlation between the neurocognitive response, design behaviour and creative design outcomes during concept generation. By combining theory about design behaviour and measurements from neurocognition, this type of study and future studies can contribute to design science by providing a framework and methods to enhance concept generation.

Acknowledgements

This material is based in part on work supported by The National Science Foundation, through Grant EEC-1929892 and EEC-1929896. Any opinions, findings and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

References

- Achard, S. & Bullmore, E. (2007) Efficiency and cost of economical brain functional networks. *PLOS Computational Biology* 3 (2), e17.
- Agbangla, N. F., Audiffren, M. & Albinet, C. T. (2017) Assessing muscular oxygenation during incremental exercise using near-infrared spectroscopy: comparison of three different methods. *Physiological Research* 66, 979–985.
- Alexiou, K., Zamenopoulos, T. & Gilbert, S. (2011) Imaging the designing brain: a neurocognitive exploration of design thinking. In *Design Computing and Cognition'10* (ed. J. S. Gero), pp. 489–504. Springer Netherlands.
- Alexiou, K., Zamenopoulos, T., Johnson, J. H. & Gilbert, S. J. (2009) Exploring the neurological basis of design cognition using brain imaging: some preliminary results. *Design Studies* 30 (6), 623–647.
- Allen, M. S. (1962) *Morphological Creativity: The Miracle of Your Hidden Brain Power: A Practical Guide to the Utilization of Your Creative Potential*. Prentice-Hall.
- Allen, E. A., Damaraju, E., Plis, S. M., Erhardt, E. B., Eichele, T., & Calhoun, V. D. (2014) Tracking whole-brain connectivity dynamics in the resting state. *Cerebral Cortex* 24 (3), 663–676; doi:10.1093/cercor/bhs352.
- Altshuller, G. S. (1984) *Creativity as an Exact Science*. CRC Press.
- Artino, A. (2008) Cognitive load theory and the role of learner experience: an abbreviated review for educational practitioners. *AACE Journal* 16 (4), 425–439.
- Ayaz, H., Shewokis, P. A., Curtin, A., Izzetoglu, M., Izzetoglu, K. & Onaral, B. (2011) Using MazeSuite and functional near infrared spectroscopy to study learning in spatial navigation. *Journal of Visualized Experiments: JoVE* (56), e3443, doi:10.3791/3443.
- Aziz-Zadeh, L., Liew, S.-L. & Dandekar, F. (2013) Exploring the neural correlates of visual creativity. *Social Cognitive and Affective Neuroscience* 8 (4), 475–480.
- Balardin, J. B., Morais, G. A. Z., Furucho, R. A., Trambaiolli, L. R., & Sato, J. R. (2017) Impact of communicative head movements on the quality of functional near-infrared spectroscopy signals: negligible effects for affirmative and negative gestures and consistent artifacts related to raising eyebrows. *Journal of Biomedical Optics* 22 (4), 046010; doi:10.1117/1.JBO.22.4.046010.
- Bartra, O., McGuire, J. T. & Kable, J. W. (2013) The valuation system: a coordinate-based meta-analysis of BOLD fMRI experiments examining neural correlates of subjective value. *NeuroImage* 76, 412–427.
- Bassett, D. S. & Sporns, O. (2017) Network neuroscience. *Nature Neuroscience*, 20 (3), 353–364.
- Beaty, R. E., Benedek, M., Silvia, P. J. & Schacter, D. L. (2016) Creative cognition and brain network dynamics. *Trends in Cognitive Sciences* 20 (2), 87–95.
- Belski, I. (2011) TRIZ course enhances thinking and problem solving skills of engineering students. *Procedia Engineering*, Proceeding of the ETRIA World TRIZ Future Conference 9, 450–460.

- Belski, I. & Belski, I.** (2015) Application of TRIZ in improving the creativity of engineering experts. *Procedia Engineering* **131**, 792–797.
- Birdi, K., Leach, D. & Magadley, W.** (2012) Evaluating the impact of TRIZ creativity training: an organizational field study. *R&D Management* **42** (4), 315–326.
- Bohm, M. R., Vucovich, J. P. & Stone, R. B.** (2005) Capturing creativity: using a design repository to drive concept innovation. In *International Design Engineering Technical Conferences and Computers and Information in Engineering Conference*, pp. 331–342. ASME; doi:[10.1115/DETC2005-85105](https://doi.org/10.1115/DETC2005-85105).
- Borgianni, Y. & Maccioni, L.** (2020) Review of the use of neurophysiological and biometric measures in experimental design research. *Artificial Intelligence for Engineering Design, Analysis and Manufacturing* **34**, 248–285; doi:[10.1017/S0890060420000062](https://doi.org/10.1017/S0890060420000062).
- Bressler, S. L. & Menon, V.** (2010) Large-scale brain networks in cognition: emerging methods and principles. *Trends in Cognitive Sciences* **14** (6), 277–290.
- Brockington, G., Balardin, J. B., Zimeo Morais, G. A., Malheiros, A., Lent, R., Moura, L. M., & Sato, J. R.** (2018) From the laboratory to the classroom: the potential of functional near-infrared spectroscopy in educational neuroscience. *Frontiers in Psychology* **9**, 1840; doi:[10.3389/fpsyg.2018.01840](https://doi.org/10.3389/fpsyg.2018.01840).
- Bryant, C, Stone, R, McAdams, D, et al** (2005) Concept generation from the functional basis of design. *International conference engineering design of ICED*, pp 15–18.
- Bullmore, E. & Sporns, O.** (2009) Complex brain networks: graph theoretical analysis of structural and functional systems. *Nature Reviews Neuroscience*, **10** (3), 186–198.
- Burle, B., Spieser, L., Roger, C., Casini, L., Hasbroucq, T. & Vidal, F.** (2015) Spatial and temporal resolutions of EEG: is it really black and white? A scalp current density view. *International Journal of Psychophysiology* **97** (3), 210–220.
- Cabeza, R. & Nyberg, L.** (2000) Imaging cognition II: an empirical review of 275 PET and fMRI Studies. *Journal of Cognitive Neuroscience* **12** (1), 1–47.
- Cazzell, M., Li, L., Lin, Z.-J., Patel, S. J. & Liu, H.** (2012) Comparison of neural correlates of risk decision making between genders: an exploratory fNIRS study of the Balloon Analogue Risk Task (BART). *NeuroImage* **62** (3), 1896–1911.
- Chu, H., Breite, A., Ciraolo, P., Franco, R. S. & Low, P. S.** (2008) Characterization of the deoxyhemoglobin binding site on human erythrocyte band 3: implications for O₂ regulation of erythrocyte properties. *Blood* **111** (2), 932–938.
- Cohen, J.** (1977) *Statistical Power Analysis for the Behavioral Sciences*. Elsevier.
- Colombo, B., Bartesaghi, N., Simonelli, L. & Antonietti, A.** (2015) The combined effects of neurostimulation and priming on creative thinking. A preliminary tDCS study on dorsolateral prefrontal cortex. *Frontiers in Human Neuroscience* **9**, 403. doi:[10.3389/fnhum.2015.00403](https://doi.org/10.3389/fnhum.2015.00403).
- Crilly, N.** (2015) Fixation and creativity in concept development: the attitudes and practices of expert designers. *Design Studies* **38**, 54–91.
- Cross, N.** (1989) *Engineering Design Methods*. Wiley.
- Cross, N.** (2006) *Designerly Ways of Knowing*. Springer-Verlag.
- Daly, S. R., Yilmaz, S., Christian, J. L., Seifert, C. M. & Gonzalez, R.** (2012) Design heuristics in engineering concept generation. *Journal of Engineering Education* **101** (4), 601–629.
- De Dreu, C. K. W., Nijstad, B. A., Baas, M., Wolsink, I. & Roskes, M.** (2012) Working memory benefits creative insight, musical improvisation, and original ideation through maintained task-focused attention. *Personality and Social Psychology Bulletin* **38** (5), 656–669.

- De Vico Fallani, F., Richiardi, J., Chavez, M. & Achard, S.** (2014) Graph analysis of functional brain networks: practical issues in translational neuroscience. *Philosophical Transactions of the Royal Society B: Biological Sciences* **369** (1653).
- Dias, R., Robbins, T. W. & Roberts, A. C.** (1996) Dissociation in prefrontal cortex of affective and attentional shifts. *Nature* **380**(6569), 69–72.
- Dietrich, A.** (2004) The cognitive neuroscience of creativity. *Psychonomic Bulletin & Review* **11** (6), 1011–1026.
- Dietrich, A. & Kanso, R.** (2010) A review of EEG, ERP, and neuroimaging studies of creativity and insight. *Psychological Bulletin* **136** (5), 822–848.
- Dorst, K. & Cross, N.** (2001) Creativity in the design process: co-evolution of problem–solution. *Design Studies* **22** (5), 425–437.
- Ekstrom, A.** (2010) How and when the fMRI BOLD signal relates to underlying neural activity: the danger in dissociation. *Brain Research Reviews* **62** (2), 233–244.
- Euston, D. R., Gruber, A. J. & McNaughton, B. L.** (2012) The role of medial prefrontal cortex in memory and decision making. *Neuron* **76** (6), 1057–1070.
- Eysenck, M. W. & Keane, M. T.** (2015) *Cognitive Psychology: A Student's Handbook*. Psychology Press.
- Fink, A., Grabner, R. H., Benedek, M., Reishofer, G., Hauswirth, V., Fally, M., Neuper, C., Ebner, F. & Neubauer, A. C.** (2009) The creative brain: investigation of brain activity during creative problem solving by means of EEG and FMRI. *Human Brain Mapping* **30** (3), 734–748.
- Fornito, A., Zalesky, A. & Bullmore, E.** (2016) *Fundamentals of Brain Network Analysis*. Academic Press.
- French, J. M.** (1999) *Conceptual Design for Engineers*, Springer.
- Funahashi, S.** (2017) Working memory in the prefrontal cortex. *Brain Sciences* **7** (5).
- Fuster, J. M.** (1988) Prefrontal cortex. *Comparative Neuroscience and Neurobiology*, Readings from the Encyclopedia of Neuroscience, pp. 107–109. Birkhäuser Boston.
- Gabora, L.** (2010) Revenge of the ‘neurds’: characterizing creative thought in terms of the structure and dynamics of memory. *Creativity Research Journal* **22** (1), 1–13.
- Garcin, B., Volle, E., Dubois, B. & Levy, R.** (2012) Similar or different? The role of the ventrolateral prefrontal cortex in similarity detection. *PLOS ONE* **7** (3), e34164.
- Gero, J. S.** (2010) Generalizing design cognition research. In *DTRS8: Interpreting Design Thinking*, Edited by: Dorst, K., Stewart, S. C., Staudinger, I., Paton, B. and Dong, A. 187–198. NSW: University of Technology Sydney.
- Gero, J. S.** (2011) Fixation and commitment while designing and its measurement. *The Journal of Creative Behavior* **45** (2), 108–115.
- Gero, J. S., Jiang, H. & Williams, C. B.** (2012) Design cognition differences when using structured and unstructured concept generation creativity techniques. In *Design Creativity 2012* (Ed. A. Duffy, Y. Nagai & T. Taura), pp. 3–12. The Design Society.
- Gero, J. S., Jiang, H. & Williams, C. B.** (2013) Design cognition differences when using unstructured, partially structured, and structured concept generation creativity techniques. *International Journal of Design Creativity and Innovation* **1** (4), 196–214.
- Gero, J.S. & Milovanovic J.** (2020) A framework for studying design thinking through measuring designers minds, bodies and brains. *Design* **6**, E19; doi:[10.1017/dsj.2020.15](https://doi.org/10.1017/dsj.2020.15).
- Gibson, C., Folley, B. S. & Park, S.** (2009) Enhanced divergent thinking and creativity in musicians: a behavioral and near-infrared spectroscopy study. *Brain and Cognition* **69** (1), 162–169. doi:[10.1016/j.bandc.2008.07.009](https://doi.org/10.1016/j.bandc.2008.07.009).

- Gilbert, S. J., Zamenopoulos, T., Alexiou, K. & Johnson, J. H. (2010) Involvement of right dorsolateral prefrontal cortex in ill-structured design cognition: an fMRI study. *Brain Research* **1312**, 79–88.
- Goel, V. (2014) Creative brains: designing in the real world. *Frontiers in Human Neuroscience* **8**, 241. doi:[10.3389/fnhum.2014.00241](https://doi.org/10.3389/fnhum.2014.00241).
- Goel, V. & Vartanian, O. (2005) Dissociating the roles of right ventral lateral and dorsal lateral prefrontal cortex in generation and maintenance of hypotheses in set-shift problems. *Cerebral Cortex* **15** (8), 1170–1177.
- Goldschmidt, G. (2016) Linkographic evidence for concurrent divergent and convergent thinking in creative design. *Creativity Research Journal* **28** (2), 115–122.
- Grabner, R. H., Ansari, D., Koschutnig, K., Reishofer, G., Ebner, F. & Neuper, C. (2009) To retrieve or to calculate? Left angular gyrus mediates the retrieval of arithmetic facts during problem solving. *Neuropsychologia* **47** (2), 604–608.
- Gramann, K., Jung, T.-P., Ferris, D. P., Lin, C.-T. & Makeig, S. (2014) *Towards a New Cognitive Neuroscience: Modeling Natural Brain Dynamics*. Frontiers E-books.
- Grön, G., Wunderlich, A. P., Spitzer, M., Tomczak, R. & Riepe, M. W. (2000) Brain activation during human navigation: gender-different neural networks as substrate of performance. *Nature Neuroscience* **3** (4), 404–408.
- Harari, O. & Graham, W. K. (1975) Tasks and task consequences as factors in individual and group brainstorming. *The Journal of Social Psychology* **95** (1), 61–65.
- Harrison, J., Izzetoglu, K., Ayaz, H., Willems, B., Hah, S., Woo, H., Shewokis, P. A., Bunce, S. C., & Onaral, B. (2013) Human performance assessment study in aviation using functional near infrared spectroscopy. In *Foundations of Augmented Cognition* (Ed. D. D. Schmorow & C. M. Fidopiastis), (pp. 433–442). Springer; doi:[10.1007/978-3-642-39454-6_46](https://doi.org/10.1007/978-3-642-39454-6_46).
- Hay, L., Duffy, A. H. B., Gilbert, S. J., Lyall, L., Campbell, G., Coyle, D. & Grealy, M. A. (2019) The neural correlates of ideation in product design engineering practitioners. *Design Science* **5**, e29; doi:[10.1017/dsj.2019.27](https://doi.org/10.1017/dsj.2019.27).
- Heilman, K. M., Nadeau, S. E. & Beversdorf, D. O. (2003) Creative innovation: possible brain mechanisms. *Neurocase* **9** (5), 369–379.
- Herold, F., Wiegel, P., Scholkmann, F. & Mueller, N. (2018) Applications of functional near-infrared spectroscopy (fNIRS) neuroimaging in exercise–cognition science: a systematic, methodology-focused review. *Journal of Clinical Medicine* **7** (12), 466.
- Helm, K. C., Jablokow, K. W., Daly, S. R., Silk, E. M., Yilmaz, S., & Suero, R. (2016, June), Evaluating the Impacts of Different Interventions on Quality in Concept Generation Paper presented at 2016 ASEE Annual Conference & Exposition, New Orleans, Louisiana. doi:[10.18260/p.26766](https://doi.org/10.18260/p.26766).
- Hoefl, F., Meyler, A., Hernandez, A., Juel, C., Taylor-Hill, H., Martindale, J. L., McMillon, G., Kolchugina, G., Black, J. M., Faizi, A., Deutsch, G. K., Siok, W. T., Reiss, A. L., Whitfield-Gabrieli, S. & Gabrieli, J. D. E. (2007) Functional and morphometric brain dissociation between dyslexia and reading ability. *Proceedings of the National Academy of Sciences of the United States of America* **104** (10), 4234–4239.
- Howard, T. J., Dekoninck, E. A. & Culley, S. J. (2010) The use of creative stimuli at early stages of industrial product innovation. *Research in Engineering Design* **21** (4), 263–274.
- Hu, M. (2018) *Neuroscience for engineering sustainability: measuring cognition during design ideation and systems thinking among students in engineering*. Master's Thesis, Virginia Tech.
- Hu, M. & Shealy, T. (2019) Application of functional near-infrared spectroscopy to measure engineering decision-making and design cognition: literature review and synthesis of methods. *Journal of Computing in Civil Engineering* **33** (6), 04019034.

- Hu, M., Shealy, T., Grohs, J. & Panneton, R. (2019) Empirical evidence that concept mapping reduces neurocognitive effort during concept generation for sustainability. *Journal of Cleaner Production* **238**, 117815.
- Huppert, T. J., Diamond, S. G., Franceschini, M. A. & Boas, D. A. (2009) HomER: a review of time-series analysis methods for near-infrared spectroscopy of the brain. *Applied Optics* **48** (10), D280–D298.
- Ilevbare, I. M., Probert, D. & Phaal, R. (2013) A review of TRIZ, and its benefits and challenges in practice. *Technovation* **33** (2), 30–37
- Jablokow, K.W., Teerlink, W., Yilmaz, S., Daly, S.R. & Silk, E.M. (2015) The Impact of Teaming and Cognitive Style on Student Perceptions of Design Ideation Outcomes Paper presented at 2015 ASEE Annual Conference & Exposition. Seattle, Washington. doi:10.18260/p.24885.
- Jiang, H., Gero, J. S., & Yen, C.-C. (2014). Exploring Designing Styles Using a Problem-Solution Division. In J. S. Gero (Ed.), *Design Computing and Cognition '12* (pp. 79–94). Springer Netherlands. https://doi.org/10.1007/978-94-017-9112-0_5.
- Jong, T. de. (2010) Cognitive load theory, educational research, and instructional design: some food for thought. *Instructional Science* **38** (2), 105–134.
- Kan, W. T., & Gero, J. (2017). Quantitative Methods for Studying Design Protocols. Springer Netherlands. <https://doi.org/10.1007/978-94-024-0984-0>.
- Keong, T. C., Wah, L. K., Aris, B. & Harun, J. (2012) Enhancing and assessing student teachers' creativity using brainstorming activities and ICT-based morphological analysis method. *Academic Research International* **2**, 241–250.
- Kirschner, P. A. (2002) Cognitive load theory: implications of cognitive load theory on the design of learning. *Learning and Instruction* **12** (1), 1–10.
- Kleibeuker, S. W., Koolschijn, P. C. M. P., Jolles, D. D., Schel, M. A., De Dreu, C. K. W. & Crone, E. A. (2013) Prefrontal cortex involvement in creative problem solving in middle adolescence and adulthood. *Developmental Cognitive Neuroscience* **5**, 197–206.
- Kohn, N. W. & Smith, S. M. (2011) Collaborative fixation: effects of others ideas on brainstorming. *Applied Cognitive Psychology* **25** (3), 359–371.
- Lara, A. H. & Wallis, J. D. (2015) The role of prefrontal cortex in working memory: a mini review. *Frontiers in Systems Neuroscience* **9**, 173.
- Lawson, B. (2006) *How Designers Think: The Design Process Demystified*. Elsevier/Architectural Press.
- Liu, L., Li, Y., Xiong, Y., Cao, J., & Yuan, P. (2018) An EEG study of the relationship between design problem statements and cognitive behaviors during conceptual design. *Artificial Intelligence for Engineering Design, Analysis and Manufacturing* **32** (3), 351–362; doi:10.1017/S0890060417000683.
- Liu, L., Nguyen, T. A., Zeng, Y. & Hamza, A. B. (2016) “Identification of Relationships Between Electroencephalography (EEG) Bands and Design Activities.” *Proceedings of the ASME 2016 International Design Engineering Technical Conferences and Computers and Information in Engineering Conference*. Volume 7: 28th International Conference on Design Theory and Methodology. Charlotte, North Carolina, USA. August 21–24, V007T06A019. ASME. <https://doi.org/10.1115/DETC2016-59104>.
- Liang, C. (2017) Brain electrical activation among experienced designers engaging in tasks that involve transforming imagination. *International Journal of Neuroscience and Behavior Studies* **1** (1), 22–33.
- Luft, C. D. B., Zioga, I., Banissy, M. J. & Bhattacharya, J. (2017) Relaxing learned constraints through cathodal tDCS on the left dorsolateral prefrontal cortex. *Scientific Reports* **7** (1), 2916.

- Manfredini, F., Malagoni, A. M., Felisatti, M., Mandini, S., Mascoli, F., Manfredini, R., Basaglia, N. & Zamboni, P.** (2009) A dynamic objective evaluation of peripheral arterial disease by near-infrared spectroscopy. *European Journal of Vascular and Endovascular Surgery* **38** (4), 441–448.
- Martindale, C.** (1995) Creativity and connectionism. *The Creative Cognition Approach*, pp. 249. MIT Press.
- Meyer, M. L., Hershfield, H. E., Waytz, A. G., Mildner, J. N. & Tamir, D. I.** (2019) Creative expertise is associated with transcending the here and now. *Journal of Personality and Social Psychology* **116** (4), 483–494.
- Milovanovic, J. & Gero, J. S.** (2018) Exploration of cognitive design behavior during design critiques. In D. Marjanovic, P. J. Clarkson, U. Lindemann, T. McAloone & C. Weber (Eds), *Human Behavior in Design Vol. 5*, pp.2099–2110. <https://doi.org/10.21278/idc.2018.0547>.
- Naseer, N. & Hong, K.-S.** (2015) Corrigendum fNIRS-based brain-computer interfaces: a review. *Frontiers in Human Neuroscience* **9**, 3. <https://doi.org/10.3389/fnhum.2015.00003>
- Nguyen, T. A., & Zeng, Y.** (2014) A physiological study of relationship between designer's mental effort and mental stress during conceptual design. *Computer-Aided Design* **54**, 3–18. doi:[10.1016/j.cad.2013.10.002](https://doi.org/10.1016/j.cad.2013.10.002).
- Osborn, A. F.** (1953) *Applied Imagination; Principles and Procedures of Creative Thinking*. Scribner.
- Pidgeon, L. M., Grealy, M., Duffy, A. H. B., Hay, L., McTeague, C., Vuletic, T., Coyle, D., & Gilbert, S. J.** (2016) Functional neuroimaging of visual creativity: a systematic review and meta-analysis. *Brain and Behavior* **6** (10), e00540. doi:[10.1002/brb3.540](https://doi.org/10.1002/brb3.540).
- Pisapia, N. D., Bacci, F., Parrott, D. & Melcher, D.** (2016) Brain networks for visual creativity: a functional connectivity study of planning a visual artwork. *Scientific Reports* **6**, 39185.
- Poldrack, R. A., Wagner, A. D., Prull, M. W., Desmond, J. E., Glover, G. H. & Gabrieli, J. D.** (1999) Functional specialization for semantic and phonological processing in the left inferior prefrontal cortex. *NeuroImage* **10** (1), 15–35.
- Runco, M. A.** (2014) Chapter 3 - Biological perspectives on creativity. *Creativity (2nd Edition)* (Ed. M. A. Runco), pp.69–108 Academic Press.
- Santanen, E. L., Briggs, R. O. & de Vreede, G.** (2000) “The cognitive network model of creativity: a new causal model of creativity and a new brainstorming technique,” *Proceedings of the 33rd Annual Hawaii International Conference on System Sciences*, Maui, HI, USA, pp. 10 pp. vol.1-, doi:[10.1109/HICSS.2000.926895](https://doi.org/10.1109/HICSS.2000.926895).
- Santosa, H., Aarabi, A., Perlman, S. B. & Huppert, T. J.** (2017) Characterization and correction of the false-discovery rates in resting state connectivity using functional near-infrared spectroscopy. *Journal of Biomedical Optics* **22** (5), 55002.
- Sato, T., Hokari, H. & Wade, Y.** (2011) “Independent component analysis technique to remove skin blood flow artifacts in functional near-infrared spectroscopy signals.” *Annual Conference of the Japanese Neural Network Society*, Nagoya, Japan.
- Savransky, S. D.** (2000) *Engineering of Creativity: Introduction to TRIZ Methodology of Inventive Problem Solving*. CRC Press.
- Schneider, W. X., Owen, A. M. & Duncan, J.** (2012) *Executive Control and the Frontal Lobe: Current Issues*. Springer Science & Business Media.
- Schönbrodt, F. D. & Perugini, M.** (2013) At what sample size do correlations stabilize? *Journal of Research in Personality* **47** (5), 609–612.
- Seitamaa-Hakkarainen, P., Huottilainen, M., Mäkelä, M., Groth, C., & Hakkarainen, K.** (2016) How can neuroscience help understand design and craft activity? The promise of

- cognitive neuroscience in design studies. *FORMakademisk - Forskningstidsskrift for Design Og Designdidaktikk* 9:1–16; doi:[10.7577/formakademisk.1478](https://doi.org/10.7577/formakademisk.1478)
- Shah, J. J., Kulkarni, S. V. & Vargas-Hernandez, N.** (2000) Evaluation of idea generation methods for conceptual design: effectiveness metrics and design of experiments. *Journal of Mechanical Design* 122 (4), 377–384.
- Shah, J. J., Smith, S. M. & Vargas-Hernandez, N.** (2003) Metrics for measuring ideation effectiveness. *Design Studies* 24 (2), 111–134.
- Shai, O., Reich, Y., Hatchuel, A., Subrahmanian, E.** (2009) Creativity theories and scientific discovery: a study of C-K theory and infused design. In *Proceedings of the Design Society: International Conference on Engineering Design – ICED'09*, 24–27 Aug 2009, Stanford CA.
- Shealy, T. & Gero, J.S.** (2019) The neurocognition of three engineering concept generation techniques. *Proceedings of the Design Society: International Conference on Engineering Design* 1 (1), 1833–1842.
- Shealy, T. & M. Hu.** (2017) “Evaluating the potential of neuroimaging methods to study engineering cognition and project-level decision making. In *Proc., Engineering Project Organization Conf.-5th Int. Megaprojects Workshop: Theory meets Practice*. Boulder, CO: The Engineering Project Organization Society.
- Shealy, Tripp, Mo Hu & John Gero.** (2018) “Patterns of cortical activation when using concept generation techniques of brainstorming, morphological analysis, and TRIZ.” In Volume 7: *30th International Conference on Design Theory and Methodology*, 1–9. ASME. doi:[10.1115/DETC2018-86272](https://doi.org/10.1115/DETC2018-86272).
- Shrout, P. E. & Rodgers, J. L.** (2018) Psychology, science, and knowledge construction: broadening perspectives from the replication crisis. *Annual Review of Psychology* 69 (1), 487–510.
- Spreafico, C., Russo, D.** (2016) Triz industrial case studies: a critical survey. *Procedia CIRP* 39, 51–56
- Storm, B. C. & Hickman, M. L.** (2015) Mental fixation and metacognitive predictions of insight in creative problem solving. *Quarterly Journal of Experimental Psychology* 68 (4), 802–813.
- Suzuki, K., Suzuki, T., Shimada, S., Tachibana, A. & Ono, Y.** (2018) Investigation of appropriate fNIRS feature to evaluate cognitive load. *Proceedings of the Annual Conference of JSAI*, Volume JSAI2018, https://doi.org/10.11517/pjsai.JSAI2018.0_2F2OS4a05.
- Sweller, J.** (1994) Cognitive load theory, learning difficulty, and instructional design. *Learning and Instruction* 4 (4), 295–312.
- Tachibana, A., Noah, J. A., Ono, Y., Taguchi, D. & Ueda, S.** (2019) Prefrontal activation related to spontaneous creativity with rock music improvisation: a functional near-infrared spectroscopy study. *Scientific Reports* 9 (1), 1–13.
- Taura, T. & Nagai, Y.** (2013) Perspectives on concept generation and design creativity. *Concept Generation for Design Creativity*, pp. 9–20. Springer.
- Toga, A. W. & Thompson, P. M.** (2003) Mapping brain asymmetry. *Nature Reviews Neuroscience* 4 (1), 37–48.
- Vieira, S. L. da S., Gero, J. S., Delmoral, J., Gattol, V., Fernandes, C. & Fernandes, A. A.** (2019a) Comparing the design neurocognition of mechanical engineers and architects: a study of the effect of designers domain. *Proceedings of the Design Society: International Conference on Engineering Design* 1 (1), 1853–1862.
- Vieira, S., Gero, J., Delmoral, J., Gattol, V., Fernandes, C., Parente, M. & Fernandes, A.** (2019b) Understanding the design neurocognition of mechanical engineers when designing and problem-solving. *Presented at the ASME 2019 International Design*

Engineering Technical Conferences and Computers and Information in Engineering Conference, American Society of Mechanical Engineers DigitalCollection. <https://doi.org/10.1115/DETC2019-97838>.

- Van Wijk, B. C. M., Stam, C. J. & Daffertshofer, A.** (2010) Comparing brain networks of different size and connectivity density using graph theory. *PLOS ONE* 5 (10), e13701.
- Verdière, K. J., Roy, R. N., & Dehais, F.** (2018). Detecting Pilot's Engagement Using fNIRS Connectivity Features in an Automated vs. Manual Landing Scenario. *Frontiers in Human Neuroscience*, 12. <https://doi.org/10.3389/fnhum.2018.00006>.
- Wu, X., Yang, W., Tong, D., Sun, J., Chen, Q., Wei, D., Zhang, Q., Zhang, M. & Qiu, J.** (2015) A meta-analysis of neuroimaging studies on divergent thinking using activation likelihood estimation. *Human Brain Mapping* 36 (7), 2703–2718.
- Zhang, M., Liu, T., Pelowski, M. & Yu, D.** (2017) Gender difference in spontaneous deception: a hyperscanning study using functional near-infrared spectroscopy. *Scientific Reports* 7 (1), 7508.
- Zhang, Y., & Zhu, C.** (2020) Assessing brain networks by resting-state dynamic functional connectivity: an fNIRS-EEG Study. *Frontiers in Neuroscience* 13, 1430; doi:[10.3389/fnins.2019.01430](https://doi.org/10.3389/fnins.2019.01430).
- Zmigrod, S., Colzato, L. S. & Hommel, B.** (2015) Stimulating creativity: modulation of convergent and divergent thinking by transcranial Direct Current Stimulation (tDCS). *Creativity Research Journal* 27 (4), 353–360.
- Zwicky, F.** (1969) *Discovery, Invention, Research Through the Morphological Approach*. MacMillan.