

RESEARCH ARTICLE

Holomorphic anomaly equation for (\mathbb{P}^2, E) and the Nekrasov-Shatashvili limit of local \mathbb{P}^2

Pierrick Bousseau¹, Honglu Fan², Shuai Guo³ and Longting Wu⁴

¹Laboratoire de mathématiques d’Orsay, Université Paris-Saclay, CNRS, 91405, Orsay, France;
 E-mail: pierrick.bousseau@u-psud.fr.

²Department of Mathematics, University of Geneva, 24, rue du Général Dufour, Case postale 64, 1211 Geneva 4, Switzerland;
 E-mail: honglu.fan@unige.ch.

³School of Mathematical Sciences, Peking University, No.5 Yiheyuan Road Haidian District, 100871, Beijing, China;
 E-mail: guoshuai@math.pku.edu.cn.

⁴Department of Mathematics, ETH Zurich, Rämistrasse 101, 8092 Zurich, Switzerland; E-mail: longting.wu@math.ethz.ch.

Received: 25 January 2020; Revised: 17 January 2021; Accepted: 2 April 2021

2020 Mathematics Subject Classification: 14N35

Abstract

We prove a higher genus version of the genus 0 local-relative correspondence of van Garrel-Graber-Ruddat: for (X, D) a pair with X a smooth projective variety and D a nef smooth divisor, maximal contact Gromov-Witten theory of (X, D) with λ_g -insertion is related to Gromov-Witten theory of the total space of $\mathcal{O}_X(-D)$ and local Gromov-Witten theory of D .

Specializing to $(X, D) = (S, E)$ for S a del Pezzo surface or a rational elliptic surface and E a smooth anticanonical divisor, we show that maximal contact Gromov-Witten theory of (S, E) is determined by the Gromov-Witten theory of the Calabi-Yau 3-fold $\mathcal{O}_S(-E)$ and the stationary Gromov-Witten theory of the elliptic curve E . Specializing further to $S = \mathbb{P}^2$, we prove that higher genus generating series of maximal contact Gromov-Witten invariants of (\mathbb{P}^2, E) are quasimodular and satisfy a holomorphic anomaly equation. The proof combines the quasimodularity results and the holomorphic anomaly equations previously known for local \mathbb{P}^2 and the elliptic curve.

Furthermore, using the connection between maximal contact Gromov-Witten invariants of (\mathbb{P}^2, E) and Betti numbers of moduli spaces of semistable one-dimensional sheaves on \mathbb{P}^2 , we obtain a proof of the quasimodularity and holomorphic anomaly equation predicted in the physics literature for the refined topological string free energy of local \mathbb{P}^2 in the Nekrasov-Shatashvili limit.

Contents

1	Introduction	2
1.1	Higher genus local-relative correspondence	2
1.2	The case of log Calabi-Yau surfaces with smooth boundary	3
1.3	Finite generation for (\mathbb{P}^2, E)	5
1.4	Quasimodularity for (\mathbb{P}^2, E)	7
1.5	Holomorphic anomaly equation for (\mathbb{P}^2, E)	8
1.6	Conifold gap conjecture	10
1.7	Nekrasov-Shatashvili limit of local \mathbb{P}^2	11
1.8	Outline of the paper	13

2	Higher genus local-relative correspondence	13
2.1	Relative Gromov-Witten theory	13
2.2	Statement of the local-relative correspondence	14
2.3	Degeneration	17
2.4	Localization	18
3	The case of log Calabi-Yau surfaces	24
3.1	Specializing Theorem 2.7 to the pair (S, E)	24
3.2	Twisted Gromov-Witten theory of the elliptic curve	27
3.3	Conclusion of the proof of Theorem 1.2	31
3.4	Stationary Gromov-Witten theory of the elliptic curve	33
4	Finite generation and quasimodularity for (\mathbb{P}^2, E)	34
4.1	Quasimodular forms for $\Gamma_1(3)$	34
4.2	Mirror of local \mathbb{P}^2 and quasimodular forms	36
4.3	Quasimodular forms: from $SL(2, \mathbb{Z})$ to $\Gamma_1(3)$	39
4.4	Genus 0 invariants of (\mathbb{P}^2, E)	39
4.5	Genus 1 invariants of (\mathbb{P}^2, E)	40
4.6	Proof of finite generation and quasimodularity	41
4.7	Genus 2 invariants of (\mathbb{P}^2, E)	43
5	Holomorphic anomaly equation for (\mathbb{P}^2, E)	44
5.1	Holomorphic anomaly equation for the elliptic curve	44
5.2	Holomorphic anomaly equation for local \mathbb{P}^2	46
5.3	Proof of the holomorphic anomaly equation for (\mathbb{P}^2, E)	46
5.4	S -degree bound on $F_g^{\mathbb{P}^2/E}$	51
Appendix A	Product formulae for the relative theory	51

1. Introduction

1.1. Higher genus local-relative correspondence

Let X be a smooth projective complex variety and D a smooth effective divisor on X . We assume that D is nef: that is, $C \cdot D \geq 0$ for every curve C on X . The main topic of the present paper is the comparison of the relative Gromov-Witten theory of the pair (X, D) and of the local Gromov-Witten theory of the total space $\text{Tot}(\mathcal{O}_X(-D))$ of the line bundle $\mathcal{O}_X(-D)$.

Let β be a curve class on X such that $\beta \cdot D > 0$. We denote by $\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)$ the moduli space of genus g stable maps of class β to the total space of $\mathcal{O}_X(-D)$, and by $\overline{\mathcal{M}}_g(X/D, \beta)$ the moduli space of genus g relative stable maps of class β to (X, D) with only one contact condition of maximal tangency along D .

As D is nef and $\beta \cdot D > 0$, $\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)$ coincides with the moduli space $\overline{\mathcal{M}}_g(X, \beta)$ of genus g stable maps of class β to X . On the other hand, there is a natural morphism $F: \overline{\mathcal{M}}_g(X/D, \beta) \rightarrow \overline{\mathcal{M}}_g(X, \beta)$ obtained by forgetting the relative marking and stabilizing. Therefore, it makes sense to try to compare the virtual fundamental classes $[\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)]^{\text{vir}}$ and $F_*[\overline{\mathcal{M}}_g(X/D, \beta)]^{\text{vir}}$ both living on $\overline{\mathcal{M}}_g(X, \beta)$.

In genus 0, van Garrel, Graber and Ruddat [29] proved that

$$[\overline{\mathcal{M}}_0(\mathcal{O}_X(-D), \beta)]^{\text{vir}} = \frac{(-1)^{\beta \cdot D - 1}}{\beta \cdot D} F_*[\overline{\mathcal{M}}_0(X/D, \beta)]^{\text{vir}} .$$

Our first main result is a generalization of this formula in arbitrary genus.

Theorem 1.1 (=Theorem 2.7). *For every $g \geq 0$, we have*

$$\begin{aligned} [\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)]^{\text{vir}} &= \frac{(-1)^{\beta \cdot D - 1}}{\beta \cdot D} F_* \left((-1)^g \lambda_g \cap [\overline{\mathcal{M}}_g(X/D, \beta)]^{\text{vir}} \right) \\ &+ \sum_{\mathcal{G} \in \overline{\mathcal{G}}_{g, \beta}} \frac{1}{|\text{Aut}(\mathcal{G})|} (\tau_{\mathcal{G}})_* \left(C_{\mathcal{G}} \cap [\overline{\mathcal{M}}_{\mathcal{G}}]^{\text{vir}} \right). \end{aligned}$$

The right-hand side of Theorem 1.1 is the sum of a leading term and correction terms. The leading term is obtained by capping $[\overline{\mathcal{M}}_g(X/D, \beta)]^{\text{vir}}$ with the top Chern class λ_g of the Hodge bundle. The correction terms are explicitly described in Section 2.2 in terms of the classes $(-1)^{g'} \lambda_{g'} \cap [\overline{\mathcal{M}}_{g'}(X/D, \beta')]^{\text{vir}}$ with $g' < g$, $\beta' \leq \beta$ (see Remark 2.8) and the Gromov-Witten theory of the rank 2 vector bundle $\mathcal{O}_D(D) \oplus \mathcal{O}_D(-D)$ over D .

The genus 0 result of [29] is proved by an application of the degeneration formula in Gromov-Witten theory. We prove Theorem 1.1 using the same strategy. The main novelty for $g > 0$ is that the degeneration formula contains new terms that are not present for $g = 0$ and come from the bubble geometry $\mathbb{P}(\mathcal{O}_D \oplus \mathcal{O}_D(D))$. We compute these correction terms using the relative virtual localization formula in Gromov-Witten theory applied to the scaling action of \mathbb{C}^* on the fibers of the \mathbb{P}^1 -bundle $\mathbb{P}(\mathcal{O}_D \oplus \mathcal{O}_D(D))$.

1.2. The case of log Calabi-Yau surfaces with smooth boundary

We specialize the higher genus local-relative correspondence given by Theorem 1.1 to $(X, D) = (S, E)$, where S is a smooth projective surface over \mathbb{C} and E is a smooth effective anticanonical divisor on S , which is nef. By the adjunction formula, E is a genus 1 curve. Examples of such surface S include del Pezzo surfaces and rational elliptic surfaces.

The local geometry $\text{Tot}(\mathcal{O}_S(-E))$ is the total space of the canonical line bundle $\mathcal{O}_S(-E) = K_S$ and is a non-compact Calabi-Yau 3-fold. Let β be a curve class on S such that $\beta \cdot E > 0$. The moduli space $\overline{\mathcal{M}}_g(\mathcal{O}_S(-E), \beta)$ has virtual dimension 0, and we define Gromov-Witten invariants

$$N_{g, \beta}^{K_S} := \int_{[\overline{\mathcal{M}}_g(\mathcal{O}_S(-E), \beta)]^{\text{vir}}} 1.$$

The moduli space $\overline{\mathcal{M}}_g(S/E, \beta)$ has virtual dimension g , and we define maximal contact relative Gromov-Witten invariants

$$N_{g, \beta}^{S/E} := \int_{[\overline{\mathcal{M}}_g(S/E, \beta)]^{\text{vir}}} (-1)^g \lambda_g.$$

The stationary Gromov-Witten invariants of the elliptic curve E are defined as

$$\langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n} \rangle_{g, n, d}^E := \int_{[\overline{\mathcal{M}}_{g, n}(E, d)]^{\text{vir}}} \prod_{j=1}^n \psi_j^{a_j} \text{ev}_j^*(\omega) \tag{1}$$

for every $\mathbf{a} = (a_1, \dots, a_n)$ and $g, d \in \mathbb{Z}_{\geq 0}$, where $\overline{\mathcal{M}}_{g, n}(E, d)$ is the moduli space of n pointed genus g degree d stable maps to E and $\omega \in H^2(E)$ is the (Poincaré dual) class of a point.

We show that the local invariants $N_{g, \beta}^{K_S}$ are explicitly determined by the relative invariants $N_{g, \beta}^{S/E}$ and the stationary theory of E . Moreover, this relation can be inverted: the relative invariants $N_{g, \beta}^{S/E}$ are explicitly determined by the local invariants $N_{g, \beta}^{K_S}$ and the stationary Gromov-Witten theory of E . Okounkov and Pandharipande [67] have completely solved the stationary Gromov-Witten theory of E . Therefore, we establish the complete equivalence of the local theory of K_S and of the relative theory of (S, E) with maximal contact and λ_g insertion.

In order to write down the formula, we introduce some notation. For every effective $\beta \in H_2(S, \mathbb{Z})$, we denote by Q^β the corresponding monomial in the algebra of the monoid of effective curve classes. In particular, we denote by Q^E the monomial Q^β for β the class of E . We form the generating series

$$F_g^{K_S} := \delta_{g,0} F_{\text{classical}}^{K_S} + \delta_{g,1} F_{\text{unstable}}^{K_S} + \sum_{\beta, \beta \cdot E > 0} N_{g,\beta}^{K_S} Q^\beta, \tag{2}$$

$$F_g^{S/E} := \delta_{g,0} F_{\text{classical}}^{K_S} + \delta_{g,1} F_{\text{unstable}}^{S/E} + \sum_{\beta, \beta \cdot E > 0} \frac{(-1)^{\beta \cdot E + g - 1}}{\beta \cdot E} N_{g,\beta}^{\mathbb{P}^2/E} Q^\beta, \tag{3}$$

where

$$F_{\text{classical}}^{K_S} := -\frac{1 - \delta_{(E \cdot E), 0}}{3!(E \cdot E)^2} (\log Q^E)^3,$$

$$F_{\text{unstable}}^{K_S} := \left(\frac{1 - \delta_{(E \cdot E), 0}}{(E \cdot E)} \frac{\chi(S)}{24} - \frac{1}{24} \right) \log Q^E,$$

$$F_{\text{unstable}}^{S/E} := -\frac{1 - \delta_{(E \cdot E), 0}}{(E \cdot E)} \frac{\chi(S)}{24} \log Q^E.$$

Here $\chi(S)$ is the Euler characteristic of S , and we adopt the convention that $\frac{1 - \delta_{(E \cdot E), 0}}{(E \cdot E)^2} = 0$ and $\frac{1 - \delta_{(E \cdot E), 0}}{(E \cdot E)} = 0$ if $E \cdot E = 0$.

According to the genus 0 result of [29], we have

$$N_{0,\beta}^{S/E} = (-1)^{\beta \cdot E - 1} (\beta \cdot E) N_{0,\beta}^{K_S},$$

that is

$$F_0^{S/E} = F_0^{K_S}.$$

For every $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n$, we consider the generating series

$$F_{g,\mathbf{a}}^E := \delta_{g,1} \delta_{n,0} F_{\text{unstable}}^E + \sum_{d \geq 0} \tilde{Q}^d \langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n,d}^E \tag{4}$$

of stationary Gromov-Witten invariants of E , where

$$F_{\text{unstable}}^E = -\frac{1}{24} \log \left((-1)^{E \cdot E} \tilde{Q} \right).$$

We view the series $F_{g,\mathbf{a}}^E$ as a function of the variables Q^β through the change of variables

$$\tilde{Q} = (-1)^{E \cdot E} Q^E \exp \left(\sum_{\beta, \beta \cdot E > 0} (-1)^{\beta \cdot E} (\beta \cdot E) N_{0,\beta}^{S/E} Q^\beta \right) \tag{5}$$

$$= (-1)^{E \cdot E} Q^E \exp \left(- \sum_{\beta, \beta \cdot E > 0} (\beta \cdot E)^2 N_{0,\beta}^{K_S} Q^\beta \right) \tag{6}$$

$$= (-1)^{E \cdot E} \exp \left(-D^2 F_0^{K_S} \right), \tag{7}$$

where D is the differential operator defined by $DQ^\beta = (\beta \cdot E)Q^\beta$.

Theorem 1.2. For every $g \geq 0$, we have

$$F_g^{K_S} = (-1)^g F_g^{S/E} + \sum_{n \geq 0} \sum_{\substack{g=h+g_1+\dots+g_n, \\ \mathbf{a}=(a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n \\ (a_j, g_j) \neq (0, 0), \sum_{j=1}^n a_j = 2h-2}} \frac{(-1)^{h-1} F_{h, \mathbf{a}}^E}{|\text{Aut}(\mathbf{a}, \mathbf{g})|} \prod_{j=1}^n (-1)^{g_j-1} D^{a_j+2} F_{g_j}^{S/E}.$$

According to Theorem 1.2, the local series $F_g^{K_S}$ is completely determined by the relative series $F_{g'}^{S/E}$ with $g' \leq g$ and the stationary theory of E . This relation is clearly invertible: that is, the relative series $F_g^{S/E}$ is completely determined by the local series $F_{g'}^{K_S}$ with $g' \leq g$ and the stationary theory of E .

Theorem 1.2 is a corollary of the specialization of Theorem 1.1 to $(X, D) = (S, E)$. The non-trivial part of the proof is to express explicitly the correction terms present in Theorem 1.1 in terms of the stationary Gromov-Witten theory of E . This is done using the quantum Riemann-Roch theorem in the form given by Coates and Givental [16]. The relatively simple form of Theorem 1.2 relies on several algebraic identities and in particular on the presence of the classical and unstable terms in the definition of $F_g^{K_S}$, $F_g^{S/E}$ and $F_{g, \mathbf{a}}^E$.

1.3. Finite generation for (\mathbb{P}^2, E)

Finite generation, quasimodularity and the holomorphic anomaly equation for the local series $F_g^{K_{\mathbb{P}^2}}$ have been recently proven using various techniques by Lho and Pandharipande [51], Coates and Iritani [17], and Fang, Ruan, Zhang and Zhou [28]. We show that similar properties also hold for the relative series $F_g^{\mathbb{P}^2/E}$.

Specializing the formulae (2) and (3) to \mathbb{P}^2 , we get

$$F_g^{K_{\mathbb{P}^2}} := -\frac{\delta_{g,0}}{18} (\log Q)^3 - \frac{\delta_{g,1}}{12} \log Q + \sum_{d \geq 1} N_{g,d}^{K_{\mathbb{P}^2}} Q^d, \tag{8}$$

$$F_g^{\mathbb{P}^2/E} := -\frac{\delta_{g,0}}{18} (\log Q)^3 - \frac{\delta_{g,1}}{24} \log Q + \sum_{d \geq 1} \frac{(-1)^{d+g-1}}{3d} N_{g,d}^{\mathbb{P}^2/E} Q^d. \tag{9}$$

The I -function for the Gromov-Witten theory of local \mathbb{P}^2 is

$$I^{K_{\mathbb{P}^2}}(z, q) = \sum_{k=0}^2 I_k(q) H^k z^{1-k} := z e^{\frac{H}{z} \log q} \sum_{d \geq 0} q^d \frac{\prod_{k=0}^{3d-1} (-3H - kz)}{\prod_{k=1}^d (H + kz)^3}, \tag{10}$$

where H is the hyperplane class of \mathbb{P}^2 . In particular, we have $I_0 = 1$, $I_1 = \log q + \bar{I}_1$, $I_2 = \frac{1}{2}(\log q)^2 + \bar{I}_1 \log q + O(q)$, where

$$\bar{I}_1 = 3 \sum_{k \geq 1} \frac{(3k-1)!}{(k!)^3} (-1)^k q^k.$$

The functions I_0 , I_1 and I_2 form a basis of solutions of the linear differential equation

$$\left[\left(q \frac{d}{dq} \right)^3 + 3q \left(q \frac{d}{dq} \right) \left(3 \left(q \frac{d}{dq} \right) + 1 \right) \left(3 \left(q \frac{d}{dq} \right) + 2 \right) \right] I = 0. \tag{11}$$

The variables q and Q are related by the mirror map

$$Q = e^{I_1}. \tag{12}$$

Explicitly, we have

$$Q = q - 6q^2 + 63q^3 - 866q^4 + 13899q^5 - 246366q^6 + \dots$$

and

$$q = Q + 6Q^2 + 9Q^3 + 56Q^4 - 300Q^5 + 3942Q^6 + \dots$$

In particular, we have $\frac{d}{dI_1} = Q \frac{d}{dQ}$. The genus 0 mirror theorem for $K_{\mathbb{P}^2}$ [30, 57, 14] computes the generating series $F_0^{K_{\mathbb{P}^2}}$ in terms of I_1 and I_2 :

$$-3Q \frac{dF_0^{K_{\mathbb{P}^2}}}{dQ} = I_2. \tag{13}$$

More precisely, we have

$$F_0^{K_{\mathbb{P}^2}} = -\frac{(\log Q)^3}{18} + 3Q - \frac{45}{8}Q^2 + \frac{244}{9}Q^3 - \frac{12333}{64}Q^4 + \frac{211878}{125}Q^5 + \dots$$

In order to describe higher genus invariants, we introduce the functions¹

$$X := (1 + 27q)^{-1}, \tag{14}$$

$$I_{11} := q \frac{dI_1}{dq}, \tag{15}$$

$$S := q \frac{d}{dq} \left(\log I_{11} - \frac{1}{3} \log X \right) = q \frac{d}{dq} \log I_{11} - \frac{1}{3}(X - 1). \tag{16}$$

The functions S , X and I_{11} are algebraically independent over \mathbb{C} (see Lemma 4.2). Therefore, the ring of functions generated by S and X is the polynomial ring

$$\mathbf{R} := \mathbb{Q}[X, S].$$

We define a grading on \mathbf{R} by $\deg X = \deg S = 1$ and denote by \mathbf{R}_k the subspace of polynomials with degree no more than k .

The finite generation property for the Gromov-Witten theory of $K_{\mathbb{P}^2}$, proved in [51, 52, 17] states that, for every $g \geq 2$, we have

$$F_g^{K_{\mathbb{P}^2}} \in [X^{-(g-1)} \cdot \mathbf{R}_{3g-3}]^{\text{reg}},$$

where $[-]^{\text{reg}}$ (the ‘orbifold regularity’ condition) is defined by

$$[-]^{\text{reg}} := \{f(X, S) : 3 \deg_X f + \deg_S f \geq 0\}. \tag{17}$$

We prove a finite generation result for the series $F_g^{\mathbb{P}^2/E}$ of relative Gromov-Witten invariants of (\mathbb{P}^2, E) .

Theorem 1.3. *For every $g \geq 2$, we have*

$$F_g^{\mathbb{P}^2/E} \in [X^{-(g-1)} \cdot \mathbf{R}_{3g-3}]^{\text{reg}}.$$

¹Our S and X are related to the A_2 and L of [51] by $A_2 = \frac{3S}{X} + \frac{1}{2}$ and $L = X^{\frac{1}{3}}$. We are defining our generators in the current way in order to have a X -degree bound $3g - 3$ for the genus g generating function.

Moreover, we have $\deg_S F_g^{\mathbb{P}^2/E} \leq 2g - 3$.

The bound of the S -degree of $F_g^{\mathbb{P}^2/E}$ by $2g - 3$ is specific to the relative theory. In general, the local series $F_g^{K_{\mathbb{P}^2}}$ is of S -degree $3g - 3$.

For small genera, the series $F_g^{K_{\mathbb{P}^2}}$ are explicitly known. For example, we have [37]

$$F_1^{K_{\mathbb{P}^2}} = -\frac{1}{12} \log q - \frac{1}{2} \log I_{11} - \frac{1}{12} \log(1 + 27q)$$

and [51]

$$F_2^{K_{\mathbb{P}^2}} = \frac{1}{X} \left(\frac{5}{8} S^3 + \frac{X}{8} S^2 + \frac{X^2}{96} S + \frac{X^3}{4320} + \frac{X^2}{4320} - \frac{X}{2160} \right).$$

Note that these formulae were first predicted in the string theory literature [2, 3, 36, 47]. We prove similar explicit formulae for the series $F_g^{\mathbb{P}^2/E}$ in low genera.

Theorem 1.4 (=Theorem 4.11). *We have*

$$F_1^{\mathbb{P}^2/E} = -\frac{1}{24} \log q + \frac{1}{24} \log(1 + 27q).$$

Theorem 1.5. *We have*

$$F_2^{\mathbb{P}^2/E} = \frac{X}{384} S - \frac{X^2}{360} + \frac{X}{240} - \frac{1}{720}. \tag{18}$$

The proof of Theorems 1.3, 4.11, and 1.5 relies on Theorem 1.2, the corresponding properties for local \mathbb{P}^2 , and results on the Gromov-Witten theory of the elliptic curve. In particular, the proof of the finite generation in Theorem 1.3 uses the quasimodularity properties of the stationary theory of the elliptic curve [67] and a rewriting of the generators S and X in terms of quasimodular forms discussed in Section 1.4 below. The proof of the S -degree bound in Theorem 1.3 is more difficult and needs the holomorphic anomaly equation described in Section 1.5.

1.4. Quasimodularity for (\mathbb{P}^2, E)

The mirror geometry of local \mathbb{P}^2 is naturally related to modular forms. Indeed, the functions $I_{11} = q \frac{dI_1}{dq}$ and $I_{12} = q \frac{dI_2}{dq}$ are periods of the fibers of the universal family of elliptic curves over the modular curve $Y_1(3) \simeq \{q \in \mathbb{C} \mid q \neq -\frac{1}{27}, 0\} \cup \{\infty\}$ ². In the context of mirror symmetry, where $Y_1(3)$ is viewed as the stringy Kähler moduli space of local \mathbb{P}^2 , the point $q = 0$ is the large volume point, $q = -\frac{1}{27}$ is the conifold point and $q = \infty$ is the orbifold point.

The modular curve $Y_1(3)$ is the quotient of the upper half-plane $\mathbb{H} = \{\tau \in \mathbb{C} \mid \text{Im } \tau > 0\}$ by the action of the congruence subgroup $\Gamma_1(3)$. The identification between $Y_1(3)$ and $\{q \in \mathbb{C} \mid q \neq -\frac{1}{27}, 0\} \cup \{\infty\}$ is given by

$$\tau = \frac{1}{2} + \frac{1}{2\pi i} \frac{I_{12}(q)}{I_{11}(q)}.$$

We denote $\mathcal{Q} = e^{2\pi i \tau}$. Remark that we have $\mathcal{Q}^3 = \tilde{\mathcal{Q}}$, where $\tilde{\mathcal{Q}} = -\exp(-D^2 F_0^{K_{\mathbb{P}^2}})$ was introduced in (7). Indeed, we have $D = 3Q \frac{d}{dQ} = \frac{3}{I_{11}} q \frac{d}{dq}$, so $-D^2 F_0^{K_{\mathbb{P}^2}} = 3 \frac{I_{12}}{I_{11}}$ follows from the mirror theorem (13).

²More precisely, it is a description of the coarse moduli space of $Y_1(3)$. As a stack, $Y_1(3)$ has a $\mathbb{Z}/3$ -orbifold point at $q = \infty$.

We define³

$$A(\tau) := \left(\frac{\eta(\tau)^9}{\eta(3\tau)^3} + 27 \frac{\eta(3\tau)^9}{\eta(\tau)^3} \right)^{\frac{1}{3}}, \quad B(\tau) := \frac{1}{4}(E_2(\tau) + 3E_2(3\tau)),$$

$$C(\tau) := \frac{\eta(\tau)^9}{\eta(3\tau)^3},$$

where

$$\eta(\tau) := \mathbb{Q}^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - \mathbb{Q}^n)$$

is the Dedekind eta function and

$$E_2(\tau) := 1 - 24 \sum_{n=1}^{\infty} \frac{n\mathbb{Q}^n}{1 - \mathbb{Q}^n}$$

is the weight 2 Eisenstein series. The functions A , B , and C are quasimodular forms for $\Gamma_1(3)$. More precisely, A and C are modular and, respectively, of weight 1 and 3, and B is quasimodular of weight 2 and depth 1. In fact, A , B , and C freely generate the ring of quasimodular forms of $\Gamma_1(3)$:

$$\text{QMod}(\Gamma_1(3)) = \mathbb{C}[A, B, C]. \tag{19}$$

Going from the variable τ to the variable q , we can express the quasimodular forms A , B , C in terms of the functions X , I_{11} , S introduced in Section 1.3 ([4, 58, 59, 76]):

$$A = I_{11}, \quad B = \frac{I_{11}^2}{X}(X + 6S), \quad C = \frac{I_{11}^3}{X}.$$

The space $[X^{-(g-1)}\mathbf{R}_{3g-3}]^{\text{reg}}$ of polynomials in S and X^{\pm} introduced in Theorem 1.3 has a very natural interpretation in terms of modular forms. Indeed, we show in Proposition 4.3 that it can be identified with the space of quasimodular forms for $\Gamma_1(3)$ of weight $6g - 6$ in the following way:

$$[X^{-(g-1)}\mathbf{R}_{3g-3}]^{\text{reg}} = C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}. \tag{20}$$

Therefore, we can rephrase Theorem 1.3 as follows.

Theorem 1.6. *For every $g \geq 2$, we have*

$$F_g^{\mathbb{P}^2/E} \in C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}.$$

Moreover, we have $\text{deg}_B F_g^{\mathbb{P}^2/E} \leq 2g - 3$.

1.5. Holomorphic anomaly equation for (\mathbb{P}^2, E)

The holomorphic anomaly equation is a remarkable structure conjecturally underlying Gromov-Witten theory of Calabi-Yau varieties [5, 6]. In the past few years, a series of works has led to the proof of several instances of the holomorphic anomaly equation: local \mathbb{P}^2 [51, 17], $\mathbb{C}^3/\mathbb{Z}^3$ [52, 17], local $\mathbb{P}^1 \times \mathbb{P}^1$ [50, 71], toric Calabi-Yau 3-folds (through the Eynard-Orantin topological recursion) [21, 22, 27, 28], the formal quintic 3-fold [53], the quintic 3-fold [35, 13], the formal elliptic curve [72], elliptic orbifold projective lines [62], elliptic curves [65] and elliptic fibrations [66]. We combine two a priori distinct

³The A , B , C defined here are respectively denoted by A , E , B^3 in [4, 76].

directions of this wave of progress, the cases of local \mathbb{P}^2 and of the elliptic curve, to formulate and prove a holomorphic equation for the maximal contact Gromov-Witten theory of (\mathbb{P}^2, E) .

Let

$$F_{g,n}^{K_{\mathbb{P}^2}} := \left(Q \frac{d}{dQ}\right)^n F_g^{K_{\mathbb{P}^2}}.$$

For $2g - 2 + n > 0$, we have [51] $F_{g,n}^{K_{\mathbb{P}^2}} \in \mathbb{Q}[S, X^\pm, I_{11}^{-1}]$, homogeneous of degree n with respect to I_{11}^{-1} .

We have the following holomorphic anomaly equation for local \mathbb{P}^2 , proved in various ways in [51], [17] and [21, 22, 27, 28]: for $2g - 2 + n > 0$,

$$\frac{X}{3 I_{11}^2} \frac{\partial}{\partial S} F_{g,n}^{K_{\mathbb{P}^2}} = \frac{1}{2} \sum_{\substack{g_1+g_2=g \\ n_1+n_2=n \\ 2g_i-2+n_i \geq 0}} \binom{n}{n_1} F_{g_1,n_1+1}^{K_{\mathbb{P}^2}} \cdot F_{g_2,n_2+1}^{K_{\mathbb{P}^2}} + \frac{1}{2} F_{g-1,n+2}^{K_{\mathbb{P}^2}}. \tag{21}$$

We prove a holomorphic anomaly equation for the series $F_g^{\mathbb{P}^2/E}$ of relative Gromov-Witten invariants of (\mathbb{P}^2, E) .

We denote

$$F_{g,n}^{\mathbb{P}^2/E} := \left(Q \frac{d}{dQ}\right)^n F_g^{\mathbb{P}^2/E}.$$

Theorem 1.7. For $2g - 2 + n > 0$,

$$F_{g,n}^{\mathbb{P}^2/E} \in \mathbb{Q}[S, X^\pm, I_{11}^{-1}],$$

homogeneous of degree n with respect to I_{11}^{-1} .

Theorem 1.8. For $2g - 2 + n > 0$, we have the following holomorphic anomaly equation

$$\frac{X}{3 I_{11}^2} \frac{\partial}{\partial S} F_{g,n}^{\mathbb{P}^2/E} = \frac{1}{2} \sum_{\substack{g_1+g_2=g, n_1+n_2=n \\ 2g_i-2+n_i \geq 0 \text{ for } i=1,2}} \binom{n}{n_1} F_{g_1,n_1+1}^{\mathbb{P}^2/E} \cdot F_{g_2,n_2+1}^{\mathbb{P}^2/E} \tag{22}$$

We note that the holomorphic anomaly equation for (\mathbb{P}^2, E) does not contain a loop term $F_{g-1,n+2}^{\mathbb{P}^2/E}$, unlike what happens for local \mathbb{P}^2 . A philosophical explanation for why the loop term is not expected for (\mathbb{P}^2, E) is that the invariants $N_{g,\beta}^{\mathbb{P}^2/E}$ are defined by insertion of the class λ_g , which is known to vanish in restriction to the locus of curves with one non-separating node (see Equation (5) of [24]).

We prove the holomorphic anomaly equation for (\mathbb{P}^2, E) using Theorem 1.2, the holomorphic anomaly equation for local \mathbb{P}^2 and the holomorphic anomaly equation for the elliptic curve recently proved by Oberdieck and Pixton [65].

We remark that Theorem 1.7 can be rewritten in terms of quasimodular forms (similar to Theorem 1.6) as

$$F_{g,n}^{\mathbb{P}^2/E} \in C^{-(2g-2+n)} \mathbb{Q}[A, B, C]_{6g-6+2n}$$

and the holomorphic anomaly equation of Theorem 1.8 can be rewritten as

$$\frac{\partial}{\partial B} F_{g,n}^{\mathbb{P}^2/E} = \frac{1}{4} \sum_{\substack{g_1+g_2=g, n_1+n_2=n \\ 2g_i-2+n_i \geq 0 \text{ for } i=1,2}} \binom{n}{n_1} F_{g_1,n_1+1}^{\mathbb{P}^2/E} \cdot F_{g_2,n_2+1}^{\mathbb{P}^2/E}. \tag{23}$$

1.6. Conifold gap conjecture

The conifold point is the point $q = -\frac{1}{27}$: that is, the cusp of the modular curve $Y_1(3)$ defined by the $\Gamma_1(3)$ -equivalence class of $\tau = 0$. Let $\tau_{\text{con}} := -\frac{1}{3\tau}$ be the modular coordinate in the neighborhood of the conifold point. For every $g \geq 2$, we define $F_{g,\text{con}}^{K_{\mathbb{P}^2}}$ (respectively, $F_{g,\text{con}}^{\mathbb{P}^2/E}$), functions of τ_{con} , by replacing in the expression of $F_g^{K_{\mathbb{P}^2}}$ (respectively, $F_g^{\mathbb{P}^2/E}$) in terms of A, B, C :

1. $A(\tau)$ by $A(\tau_{\text{con}})$,
2. $B(\tau)$ by $B(\tau_{\text{con}})$,
3. $C(\tau)$ by $C(\tau_{\text{con}})$.

Conceptually, $F_{g,\text{con}}^{K_{\mathbb{P}^2}}$ (respectively, $F_{g,\text{con}}^{\mathbb{P}^2/E}$) is the holomorphic part near the conifold point of the almost holomorphic modular function on $Y_1(3)$ whose holomorphic part is $F_g^{K_{\mathbb{P}^2}}$ (respectively, $F_g^{\mathbb{P}^2/E}$) near the large volume point $q = 0$. We refer to [4, 76] for details.

Let t_{con} be the flat coordinate near the conifold point defined as the unique solution of (11) such that $t_{\text{con}} = \frac{1}{\sqrt{3}}(1 + 27q) + O((1 + 27q)^2)$ near the conifold point. Both t_{con} and $e^{2i\pi\tau_{\text{con}}}$ are local coordinates near the conifold point. In particular, we can view $F_{g,\text{con}}^{K_{\mathbb{P}^2}}$ (respectively, $F_{g,\text{con}}^{\mathbb{P}^2/E}$) as functions of t_{con} . As C has a first-order pole and A, B are regular at the conifold point (see [76]), it follows from $F_g^{K_{\mathbb{P}^2}}$ (respectively, $F_g^{\mathbb{P}^2/E}$) $\in C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}$ that

$$F_{g,\text{con}}^{K_{\mathbb{P}^2}} = O\left(\frac{1}{t_{\text{con}}^{2g-2}}\right), \quad F_{g,\text{con}}^{\mathbb{P}^2/E} = O\left(\frac{1}{t_{\text{con}}^{2g-2}}\right) \tag{24}$$

near the conifold point $t_{\text{con}} = 0$.

According to the conifold gap conjecture [39, 41, 36], we should have, for every $g \geq 2$,

$$F_{g,\text{con}}^{K_{\mathbb{P}^2}} = \frac{B_{2g}}{2g(2g-2)} \frac{1}{t_{\text{con}}^{2g-2}} + O(1) \tag{25}$$

near the conifold point $t_{\text{con}} = 0$, where B_{2g} are the Bernoulli numbers. This conjecture can be checked explicitly in low genus (see section 10.8 [17]) but is still open in general.

The holomorphic anomaly equation fixes the B -dependence of $F_g^{K_{\mathbb{P}^2}}$ and so determines $F_g^{K_{\mathbb{P}^2}}$ up to some ambiguity living in the $(2g-1)$ -dimensional vector space $C^{-(2g-2)} \cdot \mathbb{Q}[A, C]_{6g-6}$. The coefficient of $1/t_{\text{con}}^{2g-2}$ and the $(2g-3)$ vanishings of the coefficients of $1/t_{\text{con}}^j$ for $1 \leq j \leq 2g-3$ predicted by (25), combined with the fact that the Q -expansion of $F_g^{K_{\mathbb{P}^2}}$ has no constant term, uniquely fix this ambiguity and so determine $F_g^{K_{\mathbb{P}^2}}$ completely.

We formulate a version of the conifold gap conjecture for the series $F_g^{\mathbb{P}^2/E}$.

Conjecture 1.9. *For every $g \geq 2$, we have*

$$F_{g,\text{con}}^{\mathbb{P}^2/E} = -\frac{2^{2g-1} - 1}{2^{2g-1}} \frac{|B_{2g}|}{2g(2g-1)(2g-2)} \frac{1}{t_{\text{con}}^{2g-2}} + O(1). \tag{26}$$

The holomorphic anomaly equation (23) fixes the B -dependence of $F_g^{\mathbb{P}^2/E}$ and so determines $F_g^{\mathbb{P}^2/E}$ up to some ambiguity living in the $(2g-1)$ -dimensional vector space $C^{-(2g-2)} \cdot \mathbb{Q}[A, C]_{6g-6}$. The coefficient of $1/t_{\text{con}}^{2g-2}$ and the $(2g-3)$ vanishings of the coefficients of $1/t_{\text{con}}^j$ for $1 \leq j \leq 2g-3$ predicted by (26), combined with the fact that the Q -expansion of $F_g^{\mathbb{P}^2/E}$ has no constant term, uniquely fix this ambiguity and so determine $F_g^{\mathbb{P}^2/E}$ completely.

As reviewed in the next section, $F_{g,\text{con}}^{\mathbb{P}^2/E}$ coincides with the Nekrasov-Shatashvili limit of the refined topological string on local \mathbb{P}^2 . Therefore, Conjecture 1.9 can be viewed as a special case of the conjectural universal behavior of the refined topological string near a conifold point predicted by physics arguments in [49] and [40, section 3.2].

It would be interesting to understand how (25) and (26) are related through Theorem 1.2. We leave this question open.

1.7. Nekrasov-Shatashvili limit of local \mathbb{P}^2

The results described in Sections 1.3–1.6 concern the series $F_g^{\mathbb{P}^2/E}$ of maximal contact relative Gromov-Witten invariants of (\mathbb{P}^2, E) with λ_g -insertion. However, they have exactly the form predicted in the string theory literature [40, 38] for the Nekrasov-Shatashvili limit of the refined topological string on local \mathbb{P}^2 . We explain below that this fact is not a coincidence and was one of the main motivations for our work. In combination with [8], we obtain a proof of a mathematically precise version of these physics conjectures.

According to the Gromov–Witten/stable pairs correspondence [60], the series $\bar{F}_g^{K_{\mathbb{P}^2}}$ can be described in terms of the stable pairs Pandharipande-Thomas invariants $P_{d,n}$ of local \mathbb{P}^2 [70]. More precisely, we have

$$1 + \sum_{d \geq 1} \sum_{n \in \mathbb{Z}} P_{d,n} (-x)^n Q^d = \exp \left(\sum_{g \geq 0} \bar{F}_g^{K_{\mathbb{P}^2}} u^{2g-2} \right) \tag{27}$$

where $x = e^{iu}$ and $\bar{F}_g^{K_{\mathbb{P}^2}} := \sum_{d \geq 1} N_{g,d}^{K_{\mathbb{P}^2}} Q^d$.

The stable pairs invariants $P_{d,n}$ are expected to admit a refinement $P_{d,n,j}$, such that $P_{d,n} = \sum_j P_{d,n,j}$. For local \mathbb{P}^2 , there are several approaches to the definition of $P_{d,n,j}$: cohomological/motivic [43, 48] or K-theoretic [64, 15], which conjecturally coincide. The refined topological string free energies $F_{g_1,g_2}^{K_{\mathbb{P}^2},\text{ref}}$ are then defined by the expansion

$$1 + \sum_{d \geq 1} \sum_{n,j \in \mathbb{Z}} P_{d,n,j} y^j (-x)^n Q^d = \exp \left(\sum_{(g_1,g_2) \geq 0} \bar{F}_{g_1,g_2}^{K_{\mathbb{P}^2},\text{ref}} (\epsilon_1 + \epsilon_2)^{2g_1} (-\epsilon_1 \epsilon_2)^{g_2-1} \right) \tag{28}$$

where $x = e^{i \frac{\epsilon_1 - \epsilon_2}{2}}$ and $y = e^{i \frac{\epsilon_1 + \epsilon_2}{2}}$. In the unrefined limit $\epsilon_1 = -\epsilon_2 = u, y = 1$, (28) reduces to (27), so

$$\bar{F}_g^{K_{\mathbb{P}^2}} = \bar{F}_{0,g}^{K_{\mathbb{P}^2},\text{ref}}.$$

From string theory arguments, the refined series $\bar{F}_{g_1,g_2}^{K_{\mathbb{P}^2},\text{ref}}$ are expected to behave in a way similar to the unrefined series $\bar{F}_g^{K_{\mathbb{P}^2}}$ and in particular should satisfy finite generation, quasimodularity and an appropriate generalization of the holomorphic anomaly equation [49, 40, 38]. These conjectures are widely open. The main issue is to get a geometric understanding of the change of variables $x = e^{i \frac{\epsilon_1 - \epsilon_2}{2}}$ and $y = e^{i \frac{\epsilon_1 + \epsilon_2}{2}}$. Even in the unrefined case, to prove the properties of the series $F_g^{K_{\mathbb{P}^2}}$ directly from the stable pairs – that is, using (27) as a definition and without using the Gromov-Witten interpretation – seems challenging.

However, it is possible to make progress in the Nekrasov-Shatashvili limit $\epsilon_2 \rightarrow 0$: that is, for the series $\bar{F}_{g,0}^{K_{\mathbb{P}^2},\text{ref}}$, for which we can use an alternative definition. Indeed, in the same way that the genus 0 series $\bar{F}_0^{K_{\mathbb{P}^2}}$ (more precisely, the genus 0 Gopakumar-Vafa invariants $n_{0,d}^{K_{\mathbb{P}^2}}$) can be described in terms of Euler characteristic of moduli spaces of one-dimensional semistable sheaves ([45], [12, Appendix

A)], the series $F_{g,0}^{K_{\mathbb{P}^2},\text{ref}}$ are conjecturally described in terms of Betti numbers of moduli spaces of one-dimensional semistable sheaves.

For every $d > 0$ and $\chi \in \mathbb{Z}$, let $M_{d,\chi}$ be the moduli space of one-dimensional Gieseker semistable sheaves on \mathbb{P}^2 , of degree d and Euler characteristic χ . We denote by $Ib_j(M_{d,\chi})$ the Betti numbers of the intersection cohomology of $M_{d,\chi}$. According to [9], the odd-degree part of the intersection cohomology of $M_{d,\chi}$ vanishes: $Ib_{2k+1}(M_{d,\chi}) = 0$. For every $d \in \mathbb{Z}_{>0}$ and $\chi \in \mathbb{Z}$, we define

$$\Omega_{d,\chi}^{\mathbb{P}^2}(y^{\frac{1}{2}}) := (-y^{\frac{1}{2}})^{-\dim M_{d,\chi}} \sum_{j=0}^{\dim M_{d,\chi}} Ib_{2j}(M_{d,\chi})y^j \in \mathbb{Z}[y^{\pm\frac{1}{2}}].$$

It is proved in [9] that the $\Omega_{d,\chi}^{\mathbb{P}^2}(y^{\frac{1}{2}})$ are the refined Donaldson–Thomas invariants for one-dimensional sheaves on $K_{\mathbb{P}^2}$. For $y^{\frac{1}{2}} = 1$, $\Omega_{d,\chi}^{\mathbb{P}^2}$ coincides with the genus 0 Gopakumar–Vafa invariant $n_{0,d}^{K_{\mathbb{P}^2}}$ ([45], [12, Appendix A]). Therefore, one should view $\Omega_{d,\chi}^{\mathbb{P}^2}(y^{\frac{1}{2}})$ as a refined genus 0 Gopakumar–Vafa invariant of local \mathbb{P}^2 .

Tensoring by $\mathcal{O}(1)$ gives an isomorphism between $M_{d,\chi}$ and $M_{d,\chi+d}$. Thus, we have $\Omega_{d,\chi}(y^{\frac{1}{2}}) = \Omega_{d,\chi'}(y^{\frac{1}{2}})$ if χ and χ' are equal modulo d . We define

$$\Omega_d^{\mathbb{P}^2}(y^{\frac{1}{2}}) := \frac{1}{d} \sum_{\chi \bmod d} \Omega_{d,\chi}(y^{\frac{1}{2}})$$

by averaging over the d possible values of χ modulo d . It is conjectured⁴ in [9] that $\Omega_{d,\chi}(y^{\frac{1}{2}})$ is in fact independent of χ . Assuming this conjecture, $\Omega_d^{\mathbb{P}^2}(y^{\frac{1}{2}})$ would be the common value of the $\Omega_{d,\chi}(y^{\frac{1}{2}})$.

Define

$$\bar{F}^{NS} := i \sum_{d \in \mathbb{Z}_{>0}} \sum_{k \in \mathbb{Z}_{>0}} \frac{1}{k} \frac{\Omega_d(y^{\frac{k}{2}})}{y^{\frac{k}{2}} - y^{-\frac{k}{2}}} Q^{kd} \in \mathbb{Q}(y^{\pm\frac{1}{2}})[[Q]]. \tag{29}$$

Using the change of variables $y = e^{i\hbar}$, we define series $\bar{F}_g^{NS} \in \mathbb{Q}[[Q]]$ by the expansion

$$\bar{F}^{NS} = \sum_{g \in \mathbb{Z}_{>0}} (-1)^g \bar{F}_g^{NS} \hbar^{2g-1}. \tag{30}$$

Conjecturally, we have $\bar{F}_g^{NS} = \bar{F}_{g,0}^{K_{\mathbb{P}^2},\text{ref}}$. At this point, it is unclear how the definition of \bar{F}^{NS} using one-dimensional sheaves is better than the definition of $\bar{F}_{g,0}^{K_{\mathbb{P}^2},\text{ref}}$ using stable pairs: one still needs to understand geometrically the change of variables $y = e^{i\hbar}$.

It is precisely such understanding that has been obtained in [9]. More precisely, one of the main results of [9], building on [11, 10, 8, 34], is the equality

$$\bar{F}_g^{NS} = \bar{F}_g^{\mathbb{P}^2/E}, \tag{31}$$

for every $g \geq 0$, where $\bar{F}_g^{\mathbb{P}^2/E} := \sum_{d \geq 1} \frac{(-1)^{d+g-1}}{3d} N_{g,d}^{\mathbb{P}^2/E} Q^d$. In other words, the series \bar{F}_g^{NS} have a Gromov–Witten interpretation, not as Gromov–Witten series of local \mathbb{P}^2 but as Gromov–Witten series of (\mathbb{P}^2, E) ! Therefore, all the results of Sections 1.3–1.5 for $\bar{F}_g^{\mathbb{P}^2/E}$ also hold for \bar{F}_g^{NS} , and they agree with the predictions of [40, 38].

⁴Note added in the final version (01/2021): this conjecture has now been proved by Maulik and Shen [61].

Theorem 1.10. Using (29)–(30) as definition of the Nekrasov-Shatashvili limit of the refined topological string on local \mathbb{P}^2 , the series \bar{F}_g^{NS} satisfy the finite generation, quasimodularity, holomorphic anomaly equation and low genus explicit formulae predicted by Huang and Klemm [40].

Independently from any motivation from physics, Theorem 1.10 provides a surprising way to construct quasimodular forms from Betti numbers of moduli spaces of one-dimensional semistable sheaves on \mathbb{P}^2 .

Finally, we remark that, given (31), we can view Theorem 1.2 as expressing the difference between the unrefined limit and the Nekrasov-Shatashvili limit of the refined topological string in terms of the Gromov-Witten theory of the elliptic curve. Such relation does not seem to have been predicted in the physics literature.

1.8. Outline of the paper

In Section 2, we prove Theorem 1.1: that is, the general form of the higher genus local-relative correspondence. In Section 3, we prove Theorem 1.2: that is, the explicit form of the higher genus local-relative correspondence for log Calabi-Yau surfaces. Starting with Section 4, we focus on the case of (\mathbb{P}^2, E) , and we prove the finite generation results (Theorems 1.3–1.6) and the low genus explicit formulae (Theorems 1.4–1.5). The holomorphic anomaly equation for (\mathbb{P}^2, E) (Theorem 1.8) and the S -degree bound of Theorem 1.3 are proven in Section 4. Appendix A describes a technical result used in Section 2.

2. Higher genus local-relative correspondence

2.1. Relative Gromov-Witten theory

Foundations of relative Gromov-Witten invariants were made by Li-Ruan [54], Ionel-Parker [42] and Eliashberg-Givental-Hofer [20] in symplectic geometry and Li [55, 56] in algebraic geometry. Our presentation is based on [55, 26].

Let X be a smooth projective variety and D a smooth divisor. The intersection number of a curve class β with a divisor D is denoted by $\beta \cdot D$.

A topological type Γ is a tuple $(g, n, \beta, \rho, \vec{\mu})$ where g, n are non-negative integers, $\beta \in H_2(X, \mathbb{Z})$ is a curve class and $\vec{\mu} = (\mu_1, \dots, \mu_\rho) \in \mathbb{Z}_{>0}^\rho$ is a partition of the number $\beta \cdot D$.

Let $\bar{\mathcal{M}}_\Gamma(X, D)$ be the moduli of relative stable maps with topological type Γ . There are evaluation maps

$$\begin{aligned} \text{ev}_X &= (\text{ev}_{X,1}, \dots, \text{ev}_{X,n}): \bar{\mathcal{M}}_\Gamma(X, D) \rightarrow X^n, \\ \text{ev}_D &= (\text{ev}_{D,1}, \dots, \text{ev}_{D,\rho}): \bar{\mathcal{M}}_\Gamma(X, D) \rightarrow D^\rho. \end{aligned}$$

The relative Gromov–Witten invariant with topological type Γ is defined to be

$$\langle \underline{\varepsilon} \mid \underline{\alpha} \rangle_\Gamma^{(X,D)} := \int_{[\bar{\mathcal{M}}_\Gamma(X,D)]^{\text{vir}}} \text{ev}_D^* \underline{\varepsilon} \cup \text{ev}_X^* \underline{\alpha},$$

where $\underline{\varepsilon} \in H^*(D)^{\otimes \rho}$, $\underline{\alpha} \in H^*(X)^{\otimes n}$,

$$\text{ev}_D^* \underline{\varepsilon} := \prod_{j=1}^\rho \text{ev}_{D,j}^* \varepsilon_j, \quad \text{ev}_X^* \underline{\alpha} := \prod_{i=1}^n \text{ev}_{X,i}^* \alpha_i.$$

We also allow disconnected domains. Let $\Gamma = \{\Gamma^\pi\}$ be a set of topological types. The relative

invariant with disconnected domain curves is defined by the product rule:

$$\langle \underline{\varepsilon} \mid \underline{\alpha} \rangle_{\Gamma}^{\bullet(X,D)} := \prod_{\pi} \langle \underline{\varepsilon}^{\pi} \mid \underline{\alpha}^{\pi} \rangle_{\Gamma^{\pi}}^{\bullet(X,D)}.$$

Here \bullet means possibly disconnected domains. We will call this Γ a *possibly disconnected topological type*. We now recall the definition of an *admissible graph*.

Definition 2.1 ([55, Definition 4.6]). *An admissible graph Γ is a graph without edges plus the following data:*

1. An ordered collection of legs
2. An ordered collection of weighted roots
3. A function $g : V(\Gamma) \rightarrow \mathbb{Z}_{\geq 0}$
4. A function $b : V(\Gamma) \rightarrow H_2(X, \mathbb{Z})$

Here, $V(\Gamma)$ denotes the set of vertices of Γ . Legs and roots are regarded as half-edges of the graph Γ . A relative stable morphism is associated with an admissible graph in the following way. Vertices in $V(\Gamma)$ correspond to the connected components of the domain curve. Roots and legs correspond to relative markings and interior markings, respectively. Weights on roots correspond to contact orders at the corresponding relative markings.

The functions g and b assign to a component its genus and degree, respectively. We do not spell out the formal definitions in order to avoid heavy notation, but we refer the readers to [55, Definition 4.7].

Remark 2.2. A (possibly disconnected) topological type and an admissible graph are equivalent concepts. Different terminologies emphasize different aspects. For example, admissible graphs will be glued at half-edges into actual graphs.

2.2. Statement of the local-relative correspondence

Let X be a smooth projective variety over \mathbb{C} and D be a smooth effective divisor on X . We further assume that D is nef: that is, $C \cdot D \geq 0$ for every curve C in X . Let β be a curve class on X such that $\beta \cdot D > 0$. Let $\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)$ be the moduli space of stable maps to the total space of $\mathcal{O}_X(-D)$. Here the domain curves contain no markings, and we omit the number of markings in the notation $\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)$ for simplicity. The moduli stack $\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)$ is isomorphic to $\overline{\mathcal{M}}_g(X, \beta)$ thanks to the condition $\beta \cdot D > 0$. In this case, it was proved by van Garrel, Graber and Ruddat that

Theorem 2.3 ([29]).

$$[\overline{\mathcal{M}}_0(\mathcal{O}_X(-D), \beta)]^{\text{vir}} = \frac{(-1)^{(\beta \cdot D) - 1}}{\beta \cdot D} F_* \left([\overline{\mathcal{M}}_{\Gamma}(X, D)]^{\text{vir}} \right)$$

where $\Gamma = (0, 0, \beta, 1, (\beta \cdot D))$ and $F : \overline{\mathcal{M}}_{\Gamma}(X, D) \rightarrow \overline{\mathcal{M}}_{0,0}(X, \beta)$ is the natural stabilization map, which also forgets the unique relative marking.

Note that the topological type $\Gamma = (0, 0, \beta, 1, (\beta \cdot D))$ corresponds to genus-0 relative stable maps with maximal contact at D . For convenience, whenever $\Gamma = (g, 0, \beta, 1, (\beta \cdot D))$ (genus- g maximal contact), we always denote the above relative moduli space (with only one relative marking) as $\overline{\mathcal{M}}_g(X/D, \beta)$.

Remark 2.4. In the treatment of [29], the log geometric definition of relative Gromov–Witten theory is used. But in this paper, we choose the expanded degeneration version because at the moment of writing, the references for the localization of the moduli of stable log maps are not available. Besides, since we only care about the pushforward of the relative virtual cycles to the moduli of stable maps, [1] tells us that the two choices of definitions do not affect the goal of this paper.

We generalize the main result of [29] to higher genera. More precisely, we show that $[\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)]^{\text{vir}}$ is of the form

$$\frac{(-1)^{\beta \cdot D - 1}}{\beta \cdot D} F_* \left((-1)^g \lambda_g \cap [\overline{\mathcal{M}}_g(X/D, \beta)]^{\text{vir}} \right) + \dots$$

where λ_g is the g -th Chern class of the Hodge bundle, \cap is the cap product of between cycles and Chow cohomology and ‘ \dots ’ consists of correction terms, which will be made explicit later. Before we explicitly describe the correction terms, we need the following preparation.

Definition 2.5. A graph of star type \mathcal{G} is a tuple (V, E, g, b) such that

1. The set V of vertices admits a disjoint union decomposition $V = \{v\} \coprod V_1$ such that the set E of edges contains exactly one edge between v and v_i for every $v_i \in V_1$ and no other edges.
2. A map $g: V \rightarrow \mathbb{Z}_{\geq 0}$.
3. A map $b: \{v\} \cup V_1 \rightarrow H_2(D, \mathbb{Z}) \cup H_2(X, \mathbb{Z})$ such that b maps v into $H_2(D, \mathbb{Z})$ and maps V_1 into $H_2(X, \mathbb{Z})$.

The automorphism group of \mathcal{G} consists of automorphisms of the graph (V, E) that commute with g, b . We denote it as $\text{Aut}(\mathcal{G})$.

Definition 2.6. The topological type of a graph of star type is a tuple (g, β) such that

1. g is the summation of all genera (the values of g).
2. $\beta = \iota_* b(v) + \sum_{v_i \in V_1} b(v_i) \in H_2(X, \mathbb{Z})$, where $\iota: D \hookrightarrow X$ is the natural inclusion.

We denote by $G_{g, \beta}$ the set of graphs with star type whose topological type is (g, β) . For each \mathcal{G} , we define

$$\overline{\mathcal{M}}_{\mathcal{G}} := \left(\prod_{v_i \in V_1} \overline{\mathcal{M}}_{g(v_i)}(X/D, b(v_i)) \right) \times_{D^{|E|}} \overline{\mathcal{M}}_{g(v), |E|}(D, b(v)) \tag{32}$$

where $\times_{D^{|E|}}$ is the fiber product identifying evaluation maps according to edges. The evaluation map from $\prod_{v_i \in V_1} \overline{\mathcal{M}}_{g(v_i)}(X/D, b(v_i))$ to $D^{|E|}$ is determined by the relative markings.

We define a virtual fundamental class on $\overline{\mathcal{M}}_{\mathcal{G}}$ by ‘gluing’ the virtual fundamental classes on the factors (for example, it was also introduced in [33, Equation (4)])

$$[\overline{\mathcal{M}}_{\mathcal{G}}]^{\text{vir}} := \Delta^! \left[\prod_{v_i \in V_1} \overline{\mathcal{M}}_{g(v_i)}(X/D, b(v_i)) \times \overline{\mathcal{M}}_{g(v), |E|}(D, b(v)) \right]^{\text{vir}},$$

where $\Delta: D^{|E|} \rightarrow D^{|E|} \times D^{|E|}$ is the diagonal map and $\Delta^!$ is the Gysin map.

There is a natural stabilization map

$$\mathfrak{s}: \prod_{v_i \in V_1} \overline{\mathcal{M}}_{g(v_i)}(X/D, b(v_i)) \longrightarrow \prod_{v_i \in V_1} \left(\overline{\mathcal{M}}_{g(v_i), 1}(X, b(v_i)) \times_X D \right)$$

where \times_X is the fiber product identifying the unique evaluation map and the inclusion map $D \hookrightarrow X$.

There is also a natural gluing map

$$\left(\prod_{v_i \in V_1} \left(\overline{\mathcal{M}}_{g(v_i), 1}(X, b(v_i)) \times_X D \right) \right) \times_{D^{|E|}} \overline{\mathcal{M}}_{g(v), |E|}(D, b(v)) \longrightarrow \overline{\mathcal{M}}_g(X, \beta).$$

By composition of the stabilization and gluing maps, we get a map

$$\tau_{\mathcal{G}}: \overline{\mathcal{M}}_{\mathcal{G}} \longrightarrow \overline{\mathcal{M}}_g(X, \beta).$$

Let $N_{D/X}$ be the normal bundle of D in X , and let $N_{D/X}^\vee$ be its dual. We consider the rank 2 vector bundle over D given by

$$N = N_{D/X} \oplus N_{D/X}^\vee. \tag{33}$$

There is a natural *anti-diagonal* scaling action of \mathbb{C}^* on N with weight 1 on $N_{D/X}$ and weight -1 on $N_{D/X}^\vee$.

We have a universal diagram

$$\begin{array}{ccc} U & \xrightarrow{f} & X \\ \downarrow \pi & & \\ \overline{\mathcal{M}}_{g(v),|E|}(D, b(v)) & & \end{array}$$

where π is the universal domain curve and f is the universal stable map. We view $-R^\bullet \pi_* f^* N$ as an element of the K -theory of $\overline{\mathcal{M}}_{g(v),|E|}(D, b(v))$, and we consider its equivariant Euler class⁵

$$e_{\mathbb{C}^*}(-R^\bullet \pi_* f^* N) \in A^* \left(\overline{\mathcal{M}}_{g(v),|E|}(D, b(v)) \right) (t)$$

where t is the equivariant parameter. Now let⁶

$$C_v := e_{\mathbb{C}^*}(-R^\bullet \pi_* f^* N) \prod_{v_i \in V_1} \frac{(t + \text{ev}_i^* c_1(N_{D/X}))(-1)^{d_i} d_i}{t + \text{ev}_i^* c_1(N_{D/X}) - d_i \psi_i} \tag{34}$$

where

$$\text{ev}_i : \overline{\mathcal{M}}_{g(v),|E|}(D, b(v)) \longrightarrow D \tag{35}$$

is the evaluation map for the i -th marking, $d_i := b(v_i) \cdot D$ and ψ_i is the psi-class of the i -th marking. We have

$$C_v \in A^* \left(\overline{\mathcal{M}}_{g(v),|E|}(D, b(v)) \right) (t).$$

For each $v_i \in V_1$, we define

$$C_{v_i} := \frac{t}{t + d_i \bar{\psi} + \bar{\text{ev}}^* c_1(N_{D/X})} (-1)^{g(v_i)} \lambda_{g(v_i)} \tag{36}$$

where

$$\bar{\text{ev}} : \overline{\mathcal{M}}_{g(v_i)}(X/D, b(v_i)) \longrightarrow D \tag{37}$$

is the evaluation map associated with the unique relative marking, and $\bar{\psi}$ is the psi-class associated with the unique relative marking. We have

$$C_{v_i} \in A^* \left(\overline{\mathcal{M}}_{g(v_i)}(X/D, b(v_i)) \right) (t).$$

Since there is only one relative marking, we have $\bar{\psi} = \mathfrak{s}_i^* \psi_1$, where

$$\mathfrak{s}_i : \overline{\mathcal{M}}_{g(v_i)}(X/D, b(v_i)) \longrightarrow \overline{\mathcal{M}}_{g(v_i),1}(X, b(v_i))$$

⁵It is a bivariant class because the equivariant Euler class can be expressed as a combination of Chern classes.

⁶The denominators of Equation (34) and (36) need to be expanded as a power series in t^{-1} . The expansions terminate at a finite degree because the classes in the denominators other than t are nilpotent.

is the natural stabilization map and ψ_1 is the psi-class associated with the unique marking of $\overline{\mathcal{M}}_{g(v_i),1}(X, b(v_i))$. So we may also treat C_{v_i} as a pullback class via \mathfrak{s}_i . We define

$$C_{\mathcal{G}} := \left[p_v^* C_v \prod_{v_i \in V_1} p_{v_i}^* C_{v_i} \right]_{t^0}$$

where p_v, p_{v_i} are projections from $\overline{\mathcal{M}}_{\mathcal{G}}$ to the corresponding factors and $[\dots]_{t^0}$ means that we take the constant term.

Now we are ready to state our higher genus local-relative correspondence.

Theorem 2.7 (=Theorem 1.1). *The following relation holds in $A_*(\overline{\mathcal{M}}_g(X, \beta))$.*

$$\begin{aligned} [\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)]^{\text{vir}} &= \frac{(-1)^{\beta \cdot D - 1}}{\beta \cdot D} F_* \left((-1)^g \lambda_g \cap [\overline{\mathcal{M}}_g(X/D, \beta)]^{\text{vir}} \right) \\ &+ \sum_{\mathcal{G} \in \mathcal{G}_{g,\beta}} \frac{1}{|\text{Aut}(\mathcal{G})|} (\tau_{\mathcal{G}})_* \left(C_{\mathcal{G}} \cap [\overline{\mathcal{M}}_{\mathcal{G}}]^{\text{vir}} \right). \end{aligned}$$

The proof of Theorem 2.7 is given in Sections 2.3–2.4 and uses the degeneration and localization formulae.

Remark 2.8. For a graph of star type \mathcal{G} , if $g(v) = 0$ then it follows from the Riemann-Roch theorem that $e_{C^*}(-R^* \pi_* f^* N)$ only contains negative powers of t . Therefore, $p_v^* C_v \prod_{v_i \in V_1} p_{v_i}^* C_{v_i}$ must also only contain negative powers of t , so $C_{\mathcal{G}} = 0$. Thus, a non-vanishing contribution of \mathcal{G} is only possible if $g(v) > 0$, and in particular if $g(v_i) < g$ for each $v_i \in V_1$. That also explains the absence of correction terms if $g = 0$.

2.3. Degeneration

As the first step to prove Theorem 2.7, we apply the degeneration formula to the degeneration to the normal cone of the embedding $D \hookrightarrow X$. This step is identical to the corresponding step in the proof of the main theorem of [29]. We simply recall the degeneration formula and some key lemmas in [29] for the sake of completeness.

Let \mathcal{X} be the blow-up of $X \times \mathbb{A}^1$ along $D \times \{0\}$. The space \mathcal{X} still admits a projection to \mathbb{A}^1 ; and \mathcal{X}_0 , the fiber over 0, is a union of the \mathbb{P}^1 bundle $\mathbb{P}_D(N_{D/X} \oplus \mathcal{O})$ and X glued along the section $D \cong \mathbb{P}_D(N_{D/X}) \subset \mathbb{P}_D(N_{D/X} \oplus \mathcal{O})$ and the hypersurface $D \subset X$. For convenience, we introduce the following notation:

- Denote the \mathbb{P}^1 bundle by P_0 and the component of \mathcal{X}_0 isomorphic to X by X_0 .
- Denote the section $\mathbb{P}_D(N_{D/X}) \subset \mathbb{P}_D(N_{D/X} \oplus \mathcal{O})$ by D_∞ and the section $\mathbb{P}_D(\mathcal{O}) \subset \mathbb{P}_D(N_{D/X} \oplus \mathcal{O})$ by D_0 .

Let \mathcal{L} be the line bundle on \mathcal{X} associated with the divisor of the strict transform of $D \times \mathbb{A}^1 \subset X \times \mathbb{A}^1$. Since we want to relate the local theory with the relative theory, we need to consider the total space $\text{Tot}(\mathcal{L})$ in the degeneration. The general fiber of $\text{Tot}(\mathcal{L})$ over \mathbb{A}^1 is isomorphic to the total space of $\mathcal{O}_X(-D)$, which is the target space of our local theory. The special fiber at $0 \in \mathbb{A}^1$ is a union of $\mathcal{L}|_{P_0} = \mathcal{O}_{P_0}(-D_0)$ and $\mathcal{L}|_{X_0} = \mathcal{O}_{X_0}$ glued along the corresponding divisors, both of which are isomorphic to $D \times \mathbb{A}^1$. Note that the line bundle $\mathcal{L}|_{X_0}$ is isomorphic to the trivial bundle on X .

The degeneration formula expresses the virtual cycle of $\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)$ in terms of the ones of $\overline{\mathcal{M}}_{\Gamma_1}(\mathcal{L}|_{P_0}, \mathcal{L}|_{D_\infty})$ and $\overline{\mathcal{M}}_{\Gamma_2}(X \times \mathbb{A}^1, D \times \mathbb{A}^1)$ summing over all splittings $\mathbf{i} = (\Gamma_1, \Gamma_2)$ of the genus- g degree- β curve type. The splitting forms a bipartite graph and must have matching contact orders at the corresponding relative markings. We follow [56, Definition 4.11] for treating the splitting of topological types, but we note that the more recent and general interpretation of the splitting is in terms of tropical

curves and can be found in [46], for example. Since the moduli stack of stable maps to the stack of expanded degeneration admits a morphism to $\overline{\mathcal{M}}_g(X, \beta)$ induced by the projection of the target $\mathcal{X} \rightarrow X$, we have the following version of the degeneration formula,

$$\begin{aligned}
 & [\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)]^{\text{vir}} \\
 &= \sum_{\mathbf{i}=(\Gamma_1, \Gamma_2)} \frac{\prod_{i=1}^{m(\mathbf{i})} d_i}{\text{Aut}(\mathbf{i})} \tau_* \Delta^! [\overline{\mathcal{M}}_{\Gamma_1}(\mathcal{L}|_{P_0}, \mathcal{L}|_{D_\infty}) \times \overline{\mathcal{M}}_{\Gamma_2}(X \times \mathbb{A}^1, D \times \mathbb{A}^1)]^{\text{vir}}, \tag{38}
 \end{aligned}$$

where there are $m(\mathbf{i})$ roots (relative markings) on each Γ_1 and Γ_2 , d_i are the weights (contact order) of the corresponding roots, $\Delta^!$ is the Gysin pullback induced by the diagonal $\Delta : (D \times \mathbb{A}^1)^{m(\mathbf{i})} \rightarrow (D \times \mathbb{A}^1 \times D \times \mathbb{A}^1)^{m(\mathbf{i})}$ and τ is the forgetful map from the fiber product $\overline{\mathcal{M}}_{\mathbf{i}}$ to $\overline{\mathcal{M}}_g(X, \beta)$.

First of all, there is a distinguished term in the degeneration formula, where Γ_1 consists of one vertex of genus 0, curve class $\beta \cdot D$ times of fiber class with one root of weight $\beta \cdot D$, and Γ_2 consists of one vertex of genus g , curve class β and a weight- $(\beta \cdot D)$ root. Thanks to [1], [29, Proposition 2.4] can be stated in our setting in order to understand this term.

Lemma 2.9. [29, Proposition 2.4] *Let Γ_1 be a topological type of genus 0, curve class $\beta \cdot D$ times the fiber class, and with one root of weight $\beta \cdot D$. Then*

$$\text{ev}_* [\overline{\mathcal{M}}_{\Gamma_1}(\mathcal{L}|_{P_0}, \mathcal{L}|_{D_\infty})]^{\text{vir}} = \frac{(-1)^{\beta \cdot D - 1}}{(\beta \cdot D)^2} [D],$$

where ev is the evaluation map of the (unique) relative marking.

The lemma implies that this distinguished term equals

$$\frac{(-1)^{\beta \cdot D - 1}}{\beta \cdot D} F_* \left((-1)^g \lambda_g \cap [\overline{\mathcal{M}}_g(X/D, \beta)]^{\text{vir}} \right)$$

in Theorem 2.7.

To understand the rest of the terms, we need [29, Lemma 3.1]. The idea is that in the relative theory of $(X \times \mathbb{A}^1, D \times \mathbb{A}^1)$, we rule out topological types Γ_2 with multiple relative contacts.

Lemma 2.10. [29, Lemma 3.1] *Let $\mathbf{i} = (\Gamma_1, \Gamma_2)$ be a splitting such that there exists a vertex v in Γ_2 having more than one root (roots correspond to relative markings). Then*

$$\Delta^! [\overline{\mathcal{M}}_{\Gamma_1}(\mathcal{L}|_{P_0}, \mathcal{L}|_{D_\infty}) \times \overline{\mathcal{M}}_{\Gamma_2}(X \times \mathbb{A}^1, D \times \mathbb{A}^1)]^{\text{vir}} = 0.$$

Although we use the expanded degeneration definition of relative Gromov–Witten theory, the proof of [29, Lemma 3.1] still applies here word-for-word because the gist of the proof is an intersection theoretic computation on the target spaces, and the geometry of moduli spaces plays a minor role.

A splitting $\mathbf{i} = (\Gamma_1, \Gamma_2)$ gives rise to a bipartite graph by gluing the corresponding roots. Lemma 2.10 tells us that every vertex of Γ_2 consists of only one root. This suggests that the bipartite graphs are comb-shaped and match the underlying graphs of Definition 2.5. Thus the degeneration formula is already giving us a general form that looks like Theorem 2.7 except that the pushforward of virtual cycles of $\overline{\mathcal{M}}_{\Gamma_1}(\mathcal{L}|_{P_0}, \mathcal{L}|_{D_\infty})$ are not understood in general.

2.4. Localization

We use the relative virtual localization formula [33] to understand the pushforward of $[\overline{\mathcal{M}}_{\Gamma_1}(\mathcal{L}|_{P_0}, \mathcal{L}|_{D_\infty})]^{\text{vir}}$. First recall that P_0 has two sections: D_0 and D_∞ . The normal bundle of D_0 is $\mathcal{O}_D(D)$ and the one of D_∞ is $\mathcal{O}_D(-D)$. The line bundle $\mathcal{L}|_{P_0}$ is naturally isomorphic to $\mathcal{O}_{P_0}(-D_0)$. Let \mathbb{C}^* act on $\mathcal{L}|_{P_0}$ as follows:

- \mathbb{C}^* acts fiberwise on P_0 so that the weight of fibers of the normal bundle of D_0 is 1.
- D_0 becomes an invariant divisor under the above \mathbb{C}^* -action on P_0 , and let \mathbb{C}^* act on $\mathcal{L}|_{P_0}$ in such a way that $\mathcal{L}|_{P_0} \cong \mathcal{O}_{P_0}(-D_0)$ as equivariant line bundles.

Under this construction, the \mathbb{C}^* acts fiberwise on $\mathcal{L}|_{D_0}$ with weight -1 , and it acts on $\mathcal{L}|_{D_\infty}$ trivially. The fixed loci of $\mathcal{L}|_{P_0}$ consists of D_0 and the whole total space $\mathcal{L}|_{D_\infty}$. The \mathbb{C}^* -action induces an action on $\overline{\mathcal{M}}_{\Gamma_1}^\bullet(\mathcal{L}|_{P_0}, \mathcal{L}|_{D_\infty})$. Invariant curves decompose into components mapped into D_0 , components mapped into rubbers over D_∞ and components⁷ mapped to fibers of P_0 . The localization formula can be summarized as follows:

$$\begin{aligned}
 & [\overline{\mathcal{M}}_{\Gamma_1}^\bullet(\mathcal{L}|_{P_0}, \mathcal{L}|_{D_\infty})]^{\text{eq, vir}} \\
 &= \frac{[\overline{\mathcal{M}}_{\Gamma_1}^\bullet(\mathcal{L}|_{P_0}, \mathcal{L}|_{D_\infty})^{\text{simple}}]^{\text{vir}}}{e_{\mathbb{C}^*}(N^{\text{vir}})} + \sum_{\eta=(\Gamma_1^{(0)}, \Gamma_1^{(\infty)})} \frac{[\overline{\mathcal{M}}_\eta]^{\text{vir}}}{e_{\mathbb{C}^*}(N_\eta^{\text{vir}})}, \tag{39}
 \end{aligned}$$

with the notation explained as follows. The first term corresponds to the fixed locus where the target does not degenerate (simple fixed locus according to [33]), and the second summand corresponds to the rest of fixed loci. We denote by N^{vir} and N_η^{vir} the corresponding virtual normal bundles. The index of the summation η is a splitting of the topological type Γ_1 into $\Gamma_1^{(0)}$ and $\Gamma_1^{(\infty)}$ such that:

- $\Gamma_1^{(0)}$ consists of a disjoint union of vertices with k_η ordered half-edges (corresponding to markings) and the decoration of genus and curve class attached to each vertex. We denote by $\overline{\mathcal{M}}_{\Gamma_1^{(0)}}^\bullet(D)$ the corresponding moduli of stable maps to D with possibly disconnected domains.
- $\Gamma_1^{(\infty)}$ is a possibly disconnected rubber graph (a possibly disconnected graph with assignments of genera, curve classes, inner markings, relative markings to both boundary divisors and their contact orders as detailed in [26, Definition 2.4]) with k_η ordered 0-root (corresponding to relative markings over D_0). The corresponding moduli space is $\overline{\mathcal{M}}_{\Gamma_1^{(\infty)}}^\sim(\mathcal{L}|_{D_\infty})$: that is, the moduli space of relative stable maps to the rubber target over $\mathcal{L}|_{D_\infty} \cong D \times \mathbb{A}^1$ (a description can be found in, for example, [33, Section 2.4]).

Gluing $\Gamma_1^{(0)}$ and $\Gamma_1^{(\infty)}$ along these ordered k_η half-edges forms a bipartite graph, and $\overline{\mathcal{M}}_\eta$ is the fixed locus corresponding to η . It is standard to describe an étale cover of $\overline{\mathcal{M}}_\eta$ as a fiber product of vertex moduli and edge moduli:

$$\overline{\mathcal{M}}_{\Gamma_1^{(0)}}^\bullet(D_0) \times_{D^{k_\eta}} D^{k_\eta} \times_{(D \times \mathbb{A}^1)^{k_\eta}} \overline{\mathcal{M}}_{\Gamma_1^{(\infty)}}^\sim(\mathcal{L}|_{D_\infty}) \rightarrow \overline{\mathcal{M}}_\eta,$$

where the fiber products are over evaluation maps and the inclusion of $D^{k_\eta} \rightarrow (D \times \mathbb{A}^1)^{k_\eta}$ sends D^{k_η} to $D^{k_\eta} \times \{0\}$.

The combinatorics of graph splittings and the precise formulae of $e_{\mathbb{C}^*}(N_\eta^{\text{vir}})$ are a priori very complicated. But similar to Lemma 2.10, we have the following result to cut down the number of graph types on $\Gamma_1^{(\infty)}$.

Lemma 2.11. *If any vertex of $\Gamma_1^{(\infty)}$ has more than one 0-roots, then we have*

$$[\overline{\mathcal{M}}_\eta]^{\text{vir}} = 0.$$

Note that $[\overline{\mathcal{M}}_\eta]^{\text{vir}}$ is a multiple of the pushforward of

$$[\overline{\mathcal{M}}_{\Gamma_1^{(0)}}^\bullet(D_0) \times_{D^{k_\eta}} D^{k_\eta} \times_{(D \times \mathbb{A}^1)^{k_\eta}} \overline{\mathcal{M}}_{\Gamma_1^{(\infty)}}^\sim(\mathcal{L}|_{D_\infty})]^{\text{vir}}.$$

⁷We call those \mathbb{C}^* -invariant components mapped to fibers of P_0 edge components.

The argument is the same as Lemma 2.10, and we omit the details. The conclusion of this lemma is that we can assume that each $\Gamma_1^{(0)}$ in the summation only consists of one single vertex. Thus, there is a unique edge between the vertex in $\Gamma_1^{(0)}$ and a vertex in $\Gamma_1^{(\infty)}$. Comparing with Definition 2.5 and Equation (32), we see that this shares the shape of graphs of star type, except that the vertices in V_1 represent relative stable maps to X instead of relative stable maps to rubbers over D , which appears in our localization formula (39). But combining the virtual localization formula (39) and the degeneration formula (38), we will be able to match the graph sum with Theorem 2.7.

Let us recap the whole process. Starting with the topological type Γ , we split it into three parts $\Gamma_1^{(0)}, \Gamma_1^{(\infty)}, \Gamma_2$ and sum over cycles

$$\begin{aligned} & [\overline{\mathcal{M}}_{\Gamma_1^{(0)}}^{\bullet}(D_0) \times_{D^{k_\eta}} D^{k_\eta} \times_{(D \times \mathbb{A}^1)^{k_\eta}} \overline{\mathcal{M}}_{\Gamma_1^{(\infty)}}^{\bullet}(\mathcal{L}|_{D_\infty}) \\ & \times_{(D \times \mathbb{A}^1)^{m(i)}} \overline{\mathcal{M}}_{\Gamma_2}^{\bullet}(X \times \mathbb{A}^1, D \times \mathbb{A}^1)]^{\text{vir}} \end{aligned}$$

capped with different cohomology classes. To simplify the situation, notice that the sum over $\Gamma_1^{(\infty)}$ and Γ_2 can be combined into a divisor in

$$\overline{\mathcal{M}}_{\Gamma'_2}^{\bullet}(X \times \mathbb{A}^1, D \times \mathbb{A}^1) = \overline{\mathcal{M}}_{\Gamma'_2}^{\bullet}(X, D) \times (\mathbb{A}^1)^{|V(\Gamma'_2)|}$$

corresponding to the locus where the target degenerates at least once (denoted by δ). Here Γ'_2 is the gluing of $\Gamma_1^{(\infty)}$ and Γ_2 along the corresponding roots, and $V(\Gamma'_2)$ is the set of vertices of Γ'_2 . In fact, δ can be written as another divisor that commonly appears in the virtual localization formula as follows.

According to [33, Section 2.5], $\overline{\mathcal{M}}_{\Gamma'_2}^{\bullet}(X, D)$ admits a map to \mathcal{T} (or [55, Chapter 4] as \mathfrak{X}^{rel}), the Artin stack of expanded degenerations of (X, D) . Also, by [33, Section 2.5], \mathcal{T} is isomorphic to an open substack of $\mathfrak{M}_{0,3}$ (the Artin stack of prestable 3-pointed rational curve) consisting of chains of curves that separate ∞ from 0, 1 (in [33]’s notation). There is a divisor corresponding to cotangent lines at ∞ , and we denote by Ψ its pullback to $\overline{\mathcal{M}}_{\Gamma'_2}^{\bullet}(X, D)$.

Lemma 2.12. *Ψ is linearly equivalent to δ .*

Proof. When identifying \mathcal{T} as an open substack of $\mathfrak{M}_{0,3}$, the relation $\Psi = \delta$ is simply the pullback of the relation $\psi_1 = D_{1|23}$ on $A^2(\mathfrak{M}_{0,3})$. □

Now that Ψ represents the decomposition of Γ'_2 into different $\Gamma_1^{(\infty)}$ and Γ_2 , we can simplify the localization formula combined with the degeneration formula into the following form:

$$\begin{aligned} & [\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)]^{\text{vir}} \\ & = \left[\sum_{\mathbf{i}=(\Gamma'_1, \Gamma'_2)} \frac{1}{\text{Aut}(\mathbf{i})} (\tau_{\mathbf{i}})_* \left(p_1^* \left(\frac{e_{\mathbb{C}^*}(-R^\bullet(\pi_1)_* f_1^* N)}{\prod_{e_i \in \text{HE}(\Gamma'_1)} \left(\frac{t + e^{v_i} c_1(N_{D/X})}{d_i} - \psi_i \right)} \right) \right. \right. \\ & \left. \left. p_2^* \left(\left(1 + \frac{\Psi}{-t - \Psi} \right) \cdot e(R^1(\pi_2)_* f_2^* \mathcal{O}) \right) \text{Edge}(\mathbf{i}) \cap [\overline{\mathcal{M}}_{\Gamma_1}^{\bullet}(D_0) \times_{D^{k_i}} \overline{\mathcal{M}}_{\Gamma_2}^{\bullet}(X, D)]^{\text{vir}} \right) \right]_{t^0}, \quad (40) \end{aligned}$$

where


- \mathbf{i} is a splitting such that each vertex of Γ'_2 has one unique root (corresponding to relative marking);
- Γ'_1 consists of a single vertex, which is allowed to be a degree-zero genus-zero unstable vertex;
- $\tau_{\mathbf{i}}$ is the natural gluing map

$$\tau_{\mathbf{i}} : \overline{\mathcal{M}}_{\Gamma_1}^{\bullet}(D_0) \times_{D^{k_i}} \overline{\mathcal{M}}_{\Gamma_2}^{\bullet}(X, D) \rightarrow \overline{\mathcal{M}}_g(X, \beta) = \overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta);$$

- Edge(i) is a certain bivariant class depending on edges in the bipartite graph splitting i (corresponding to the edge contribution in the localization);
- p_1 and p_2 are the projections from $\overline{\mathcal{M}}_{\Gamma_1}(D_0) \times_{D^{k_1}} \overline{\mathcal{M}}_{\Gamma_2}^\bullet(X, D)$ to the first and second factors, respectively;
- $\text{HE}(\Gamma'_1)$ refers to the set of half-edges of Γ'_1 and k_i is the number of half-edges of Γ'_1 , which is the same as the number of roots (or vertices) of Γ'_2 ;
- N and ev_i are defined by (33) and (35) in Section 2.2, and d_i is the weight⁸ of the root glued to the half-edge e_i .
- π_1, π_2 and f_1, f_2 are, respectively, the universal curves and universal maps for $\overline{\mathcal{M}}_{\Gamma_1}(D_0)$ and $\overline{\mathcal{M}}_{\Gamma_2}^\bullet(X, D)$ (similar to π and f used in Section 2.2).

We make the convention that $\overline{\mathcal{M}}_{\Gamma_1}(D_0) = D$ when Γ'_1 consists of an unstable vertex. In this case, the whole $p_1^*(\dots)$ factor degenerates into 1. We highlight a few key points regarding how to deduce this formula:

1. Combining the degeneration formula and the localization formula, we have three levels $\Gamma_1^{(0)}, \Gamma_1^{(\infty)}$ and Γ_2 on the graph. To deduce this formula, we turn $\Gamma_1^{(0)}$ into Γ'_1 and combine $\Gamma_1^{(\infty)}, \Gamma_2$ into Γ'_2 . An illustration of an example can be the following⁹:



By Lemma 2.12, the sum over different decomposition of Γ'_2 corresponds to the divisor Ψ written in the numerator in the factor

$$p_2^* \left(\left(1 + \frac{\Psi}{-t - \Psi} \right) \cdot e(R^1(\pi_2)_* f_2^* \mathcal{O}) \right).$$

Once a decomposition of Γ'_2 is given, the Ψ in the denominator becomes the corresponding target psi-class on the rubber moduli. Note that the extra summand ‘1’ corresponds to the simple fixed locus in the localization formula.

2. Recall that $N = N_{D/X} \oplus N_{D/X}^\vee$ under the anti-diagonal scaling action. The factor $e_{\mathbb{C}^*}(-R^*(\pi_1)_* f_1^* N)$ comes from the moving part computation on vertices over D_0 . $N_{D/X}$ comes from the log-tangent bundle of P_0 , and $N_{D/X}^\vee$ comes from the restriction of \mathcal{L} to D_0 .
3. The denominators

$$\prod_{e_i \in \text{HE}(\Gamma'_1)} \left(\frac{t + ev_i^* c_1(N_{D/X})}{d_i} - \psi_i \right), \quad -t - \Psi$$

correspond to the smoothing of nodes and the smoothing of the target, respectively.

4. In the formula, the edges should contribute an automorphism factor $\frac{1}{\prod_{i=1}^{k_\eta} d_i}$. But the gluing of edge components with the relative markings on rubber components also contribute $\prod_{i=1}^{k_\eta} d_i$. These two factors cancel each other. Note that in the case of the simple fixed locus, there are no rubber components, but there is a factor $\prod_{i=1}^{k_\eta} d_i$ coming from the degeneration formula (38).

⁸The weight of a root corresponds to the contact order of a relative marking.

⁹Each integer indicates the genus attached to the vertex. The red and blue vertices show clearly how the genera change after we combine $\Gamma_1^{(\infty)}, \Gamma_2$ into Γ'_2 .

One notices that the splitting i already resembles the graphs of star type defined in Definition 2.5. We are left to identify the formulae. It boils down to two things: matching suitable factors with the C_v term defined in Equation (34), and matching the rest with the C_{v_i} terms defined in Equation (36).

2.4.1. Matching with C_v

By explicitly computing $\text{Edge}(i)$, one can combine it with

$$p_1^* \left(\frac{e_{\mathbb{C}^*}(-R^\bullet(\pi_1)_* f_1^* N)}{\prod_{e_i \in \text{HE}(\Gamma'_1)} \left(\frac{t + \text{ev}_i^* c_1(N_{D/X})}{d_i} - \psi_i \right)} \right)$$

to obtain C_v . Some details about computing $\text{Edge}(i)$ are presented in the following.

$\text{Edge}(i)$ encodes the deformation and obstruction contributions of edge components.¹⁰ Recall that edge components are those \mathbb{C}^* -invariant components that are mapped to fibers of P_0 . For each half-edge e_i , we can construct an edge moduli space $\overline{\mathcal{M}}_{e_i}$, which consists of \mathbb{C}^* -invariant maps from a rational curve to a fiber of P_0 with multiplicity d_i . Obviously, $\overline{\mathcal{M}}_{e_i}$ can be parameterized by D . A priori, we should write $\overline{\mathcal{M}}_{\Gamma'_1}(D_0) \times_{D^{k_i}} \overline{\mathcal{M}}_{\Gamma'_2}^\bullet(X, D)$ as $\overline{\mathcal{M}}_{\Gamma'_1}(D_0) \times_{D^{k_i}} \prod_{e_i} \overline{\mathcal{M}}_{e_i} \times_{D^{k_i}} \overline{\mathcal{M}}_{\Gamma'_2}^\bullet(X, D)$. Then $\text{Edge}(i)$ can be written as a product of contributions pulling back from different $\overline{\mathcal{M}}_{e_i}$ (using the projections p_{e_i}). Next, let us determine each individual contribution.

Let π_{e_i} be the projection from the universal curves of $\overline{\mathcal{M}}_{e_i}$ and f_{e_i} be the universal map. First, suppose that the unique vertex of Γ'_1 is stable. In this case, the contribution of $\overline{\mathcal{M}}_{e_i}$ to $\text{Edge}(i)$ can be written as a product of three factors. The equivariant Euler class of the moving part¹¹ of $R^\bullet(\pi_{e_i})_* f_{e_i}^* TP_0(-\log D_\infty)$ gives the first factor

$$\frac{d_i^{d_i}}{(d_i)!(t + \text{ev}_i^* c_1(N_{D/X}))^{d_i}}. \tag{41}$$

The equivariant Euler class of the moving part of $R^\bullet(\pi_{e_i})_* f_{e_i}^*(\mathcal{L}|_{P_0})$ contributes another factor:

$$\frac{(d_i - 1)!(-t - \text{ev}_i^* c_1(N_{D/X}))^{d_i - 1}}{d_i^{d_i - 1}}. \tag{42}$$

Finally, there is a factor $-(t + \text{ev}_i^* c_1(N_{D/X}))^2$ coming from the gluing of the edge component and the component over D_0 . Putting them together, we know that the contribution of $\overline{\mathcal{M}}_{e_i}$ to $\text{Edge}(i)$ is $(-1)^{d_i} (t + \text{ev}_i^* c_1(N_{D/X}))$. So we obtain

$$\text{Edge}(i) = \prod_{e_i \in \text{HE}(\Gamma'_1)} (-1)^{d_i} (t + \text{ev}_i^* c_1(N_{D/X})).$$

Next, suppose that the vertex over D_0 is unstable (for example, the leading term in Theorem 2.7). The valence-2 case (the unstable vertex has two half-edges) will contribute 0 to the right-hand side of (40) for the dimensional reason.¹² When the unstable vertex has valence 1, there is only one half-edge e_i in $\text{HE}(\Gamma'_1)$. In this case, the gluing factor $-(t + \text{ev}_1^* c_1(N_{D/X}))^2$ disappears because the component over D_0 degenerates. But there is an extra factor $(t + \text{ev}_1^* c_1(N_{D/X}))/d_1$ coming from the moving part

¹⁰ It can be evaluated using the method of [32, Section 4]. See also [73, Appendix B].

¹¹ For a more precise definition, one should check [32, Section 1] for the definition of the virtual normal bundle, and look for the term ‘moving part’ after the proof of [32, Proposition 1].

¹² In this case, one can check that the dimension of

$$p_2^* \left(e(R^1(\pi_2)_* f_2^* \mathcal{O}) \right) \cap [\overline{\mathcal{M}}_{\Gamma'_1}(D_0) \times_{D^{k_i}} \overline{\mathcal{M}}_{\Gamma'_2}^\bullet(X, D)]^{\text{vir}}$$

is smaller than the dimension of $[\overline{\mathcal{M}}_g(\mathcal{O}_X(-D), \beta)]^{\text{vir}}$.

of the space of infinitesimal automorphisms of domain curves. Combining with (41), (42) (setting $i = 1$ there), we may conclude that in this case

$$\text{Edge}(i) = \frac{(-1)^{d_1-1}}{d_1}.$$

This gives another explanation of the coefficient $\frac{(-1)^{\beta \cdot D-1}}{\beta \cdot D}$ appearing in the leading term of Theorem 2.7 (comparing with Lemma 2.9).

2.4.2. Matching with C_{v_i}

First, each factor $(-1)^{g(v)} \lambda_{g(v)}$ of C_{v_i} in (36) comes from $e(R^1(\pi_2)_* f_2^* \mathcal{O})$. To match the other factor in C_{v_i} requires slightly more effort. For a vertex $v \in V(\Gamma)$, denote by γ_v the graph consisting of a single vertex v plus all the decorations on v . The goal now is to rewrite the pushforward of

$$\left(1 + \frac{\Psi}{-t - \Psi}\right) \cap [\overline{\mathcal{M}}_{\Gamma_2}^\bullet(X, D)]^{\text{vir}}$$

in terms of pushforward classes from the product $\prod_{v \in V(\Gamma_2)} \overline{\mathcal{M}}_{\gamma_v}(X, D)$ and match it with the remaining factor of C_{v_i} . More precisely, we need the following lemma.

Lemma 2.13. *We have the following identity:*

$$\begin{aligned} & \tau_* \left(\frac{t}{t + \Psi} \cap [\overline{\mathcal{M}}_{\Gamma_2}^\bullet(X, D)]^{\text{vir}} \right) \\ &= \tau'_* \left(\prod_{v \in V(\Gamma_2)} p_v^* \left(\frac{t}{t + \Psi} \right) \cap \left[\prod_{v \in V(\Gamma_2)} \overline{\mathcal{M}}_{\gamma_v}(X, D) \right]^{\text{vir}} \right), \end{aligned}$$

where p_v is the projection to the factor corresponding to v , and the two Ψ on both sides are the target psi-classes on their corresponding moduli spaces, and τ, τ' are the corresponding stabilization maps to $\prod_{v \in V(\Gamma_2)} \overline{\mathcal{M}}_{g(v),1}(X, b(v)) \times_X D$.

Lemma 2.13 is a special case of Lemma Appendix A.4. Each factor on the right-hand side of Lemma 2.13 matches with the $\frac{t}{t + d_i \bar{\psi} + e \bar{v}^* c_1(N_{D/X})}$ part of C_{v_i} because of the following two lemmas.

Lemma 2.14. *Let e be a root with multiplicity d . Then, we have the following identity on the moduli of relative stable maps $\overline{\mathcal{M}}_\Gamma(X, D)$:*

$$\Psi = d\psi_e + e \bar{v}_e^* c_1(N_{D/X}),$$

where ψ_e is the psi-class of the relative marking e defined using the universal curve over $\overline{\mathcal{M}}_\Gamma(X, D)$.

For a proof, see [44, Theorem 5.13.1].

Lemma 2.15. *If Γ is a connected admissible graph with only one root, let ψ be the psi-class of the relative marking and recall that $\bar{\psi}$ is the pullback of the psi-class from the moduli of stable maps to X . Then we have $\psi = \bar{\psi}$.*

Proof. Because Γ has only one root and D is nef, when the target expands, there can only be one component in the rubber. Due to the stability condition on the rubber, the relative marking cannot lie on an unstable component (a multiple cover of a fiber of a rubber target with only two relative markings). Thus, the stabilization does not contract components containing the relative marking. As a result, the psi-class and the pullback psi-class are the same. \square

3. The case of log Calabi-Yau surfaces

Let S be a smooth projective surface over \mathbb{C} and E a smooth effective anticanonical divisor on S , which is nef. By the adjunction formula, E is a genus 1 curve. In this section, we prove Theorem 1.2 expressing the local series $F_g^{K_S}$ in terms of the relative series $F_g^{S/E}$ and the stationary series $F_{g,a}^E$ of the elliptic curve E . We systematically use the notation introduced in Section 1.2.

3.1. Specializing Theorem 2.7 to the pair (S, E)

Using Theorem 2.7, we obtain a relation between the local invariants $N_{g,\beta}^{K_S}$ and the relative invariants $N_{g,\beta}^{S/E}$.

Let $N_{E/S}$ be the normal bundle to E in S . We consider the rank 2 vector bundle $N := N_{E/S} \oplus N_{E/S}^\vee$ over E and the anti-diagonal scaling action of \mathbb{C}^* on N with weight 1 on $N_{E/S}$ and weight -1 on $N_{E/S}^\vee$. We denote by t the corresponding equivariant parameter. For every $d_E \geq 0$ and $\mathbf{d} = (d_1, \dots, d_n)$, we set

$$N_{h,d_E}^{E,tw}(\mathbf{d}) := \int_{[\overline{\mathcal{M}}_{h,n}(E,d_E)]^{vir}} \left(\prod_{j=1}^n \frac{t \operatorname{ev}_j^* \omega}{t - d_j \psi_j} \right) e_{\mathbb{C}^*}(-R^* \pi_* f^* N),$$

where ω is a point class for E . Note that the equivariant classes

$$\prod_{j=1}^n \frac{t \operatorname{ev}_j^* \omega}{t - d_j \psi_j} \quad \text{and} \quad e_{\mathbb{C}^*}(-R^* \pi_* f^* N)$$

have equivariant degrees n and $2h - 2$,¹³ respectively. The summation of their degrees equals to the virtual dimension of $[\overline{\mathcal{M}}_{h,n}(E, d_E)]^{vir}$. So the degree of $N_{h,d_E}^{E,tw}(\mathbf{d}) \in \mathbb{Q}(t)$ must be zero. Note that the equivariant parameter t has degree 1. It further implies that $N_{h,d_E}^{E,tw}(\mathbf{d}) \in \mathbb{Q}$: that is, does not depend on t .

Proposition 3.1. *For every $\beta \in H_2(S, \mathbb{Z})$ such that $\beta \cdot E > 0$ we have*

$$N_{g,\beta}^{K_S} = \frac{(-1)^{\beta \cdot E - 1}}{\beta \cdot E} N_{g,\beta}^{S/E} + \sum_{n \geq 0} \sum_{\substack{g=h+g_1+\dots+g_n \\ \beta=d_E[E]+\beta_1+\dots+\beta_n \\ d_E \geq 0, \beta_j \cdot E > 0}} \frac{N_{h,d_E}^{E,tw}(\beta \cdot E)}{|\operatorname{Aut}(\beta, \mathbf{g})|} \prod_{j=1}^n \left((-1)^{\beta_j \cdot E} (\beta_j \cdot E) N_{g_j,\beta_j}^{S/E} \right)$$

where

$$\beta \cdot E = (\beta_1 \cdot E, \dots, \beta_n \cdot E),$$

$$\frac{1}{|\operatorname{Aut}(\beta, \mathbf{g})|} := \frac{1}{|\operatorname{Aut}((\beta_1, g_1), \dots, (\beta_n, g_n))|}.$$

Proof. We apply Theorem 2.7. Let $\mathcal{G} \in G_{g,\beta}$ with $|V_1| = n$, $g = h + \sum_{j=1}^n g_j$, $\beta = d_E[E] + \sum_{j=1}^n \beta_j$. We denote $d_j = \beta_j \cdot E$. The contribution of \mathcal{G} is

$$(\tau_{\mathcal{G}})_* \left[p_v^* C_v \prod_{v_j \in V_1} p_{v_j}^* C_{v_j} \right]$$

¹³ The equivariant degree of $e_{\mathbb{C}^*}(-R^* \pi_* f^* N)$ can be computed by the Riemann-Roch formula.

where $\tau_{\mathfrak{G}}$ is the gluing map

$$\left(\prod_{v_j \in V_1} \overline{\mathcal{M}}_{g_j}(S/E, \beta_j) \right) \times_{E^n} \overline{\mathcal{M}}_{h,n}(E, d_E) \longrightarrow \overline{\mathcal{M}}_{\mathfrak{G}}(S, \beta).$$

According to (36), we have

$$C_{v_j} = \frac{t}{t + d_j \bar{\psi} + \bar{e}v^* c_1(N_{E/S})} (-1)^{g_j} \lambda_{g_j}.$$

The key point is that $\dim[\overline{\mathcal{M}}_{g_j}(S/E, \beta_j)]^{\text{vir}} = g_j$. Therefore the insertion of $(-1)^{g_j} \lambda_{g_j}$ already eats up all the dimension of $[\overline{\mathcal{M}}_{g_j}(S/E, \beta_j)]^{\text{vir}}$, so C_{v_j} reduces to $(-1)^{g_j} \lambda_{g_j}$ and the only possibly non-vanishing contribution of the class of the diagonal $E \hookrightarrow E \times E$ defining the gluing $\tau_{\mathfrak{G}}$ is $p_{v_j}^* 1 \times p_v^* \omega$ on $\overline{\mathcal{M}}_{g_j}(S/E, \beta_j) \times \overline{\mathcal{M}}_{h,n}(E, d_E)$, where $1 \in H^0(E)$ and $\omega \in H^2(E)$. Thus, the contribution of $\overline{\mathcal{M}}_{g_j}(S/E, \beta_j)$ is

$$\int_{[\overline{\mathcal{M}}_{g_j}(S/E, \beta_j)]^{\text{vir}}} (-1)^{g_j} \lambda_j = N_{g_j, \beta_j}^{S/E}$$

and the contribution of $\overline{\mathcal{M}}_{h,n}(E, d_E)$ is

$$\int_{[\overline{\mathcal{M}}_{h,n}(E, d_E)]^{\text{vir}}} C_v \prod_{j=1}^n \text{ev}_j^* \omega.$$

According to (34), we have

$$C_v = e_{\mathbb{C}^*}(-R^\bullet \pi_* f^* N) \prod_{j=1}^n \frac{(t + \text{ev}_j^* c_1(N_{E/S}))(-1)^{d_j} d_j}{t + \text{ev}_j^* c_1(N_{E/S}) - d_j \psi_j}.$$

As $c_1(N_{E/S}) \cup \omega = 0$ in $A^\bullet(E)$, we have

$$C_v \prod_{j=1}^n \text{ev}_j^* \omega = e_{\mathbb{C}^*}(-R^\bullet \pi_* f^* N) \prod_{j=1}^n \frac{t(-1)^{d_j} d_j}{t - d_j \psi_j} \text{ev}_j^* \omega,$$

so the contribution of $\overline{\mathcal{M}}_{h,n}(E, d_E)$ is indeed $N_{h,d_E}^{E, \text{tw}}(\mathbf{d}) \prod_{j=1}^n (-1)^{d_j} d_j$. □

Expanding the denominator¹⁴ in the formula for $N_{h,d_E}^{E, \text{tw}}(\mathbf{d})$, and using that $\dim[\overline{\mathcal{M}}_{h,n}(E, d_E)]^{\text{vir}} = 2h - 2 + n$, we get

$$N_{h,d_E}^{E, \text{tw}}(\mathbf{d}) = \sum_{\substack{\mathbf{a}=(a_1, \dots, a_n) \\ a_j \geq 0, \sum_j a_j \leq 2h-2}} N_{h,d_E, \mathbf{a}}^{E, \text{tw}} \prod_{j=1}^n d_j^{a_j},$$

where, for every $\mathbf{a} = (a_1, \dots, a_n)$,

$$N_{h,d_E, \mathbf{a}}^{E, \text{tw}} := \frac{1}{t^{\sum_{j=1}^n a_j}} \int_{[\overline{\mathcal{M}}_{h,n}(E, d_E)]^{\text{vir}}} \left(\prod_{j=1}^n \text{ev}_j^*(\omega) \psi_j^{a_j} \right) e_{\mathbb{C}^*}(-R^\bullet \pi_* f^* N).$$

¹⁴ The first few terms of the expansion are

$$\prod_{j=1}^n \frac{t}{t - d_j \psi_j} = 1 + \frac{1}{t} \sum_{j=1}^n d_j \psi_j + \frac{1}{t^2} \left(\sum_{j=1}^n d_j^2 \psi_j^2 + \sum_{1 \leq i < j \leq n} d_i d_j \psi_i \psi_j \right) + \dots$$

Therefore, we have

$$N_{g,\beta}^{K_S} = \frac{(-1)^{\beta \cdot E - 1}}{\beta \cdot E} N_{g,\beta}^{S/E} + \sum_{n \geq 0} \sum_{\substack{g=h+g_1+\dots+g_n \\ \beta=d_E[E]+\beta_1+\dots+\beta_n \\ d_E \geq 0, \beta_j \cdot E > 0}} \sum_{\substack{\mathbf{a}=(a_1,\dots,a_n) \\ a_j \geq 0, \sum_j a_j \leq 2h-2}} \frac{N_{h,d_E,\mathbf{a}}^{E,tw}}{|\text{Aut}(\boldsymbol{\beta}, \mathbf{g})|} \prod_{j=1}^n \left((-1)^{\beta_j \cdot E} (\beta_j \cdot E)^{a_j+1} N_{g_j,\beta_j}^{S/E} \right).$$

Using generating series, we package the above recursive formula as follows. Let

$$\bar{F}_g^{K_S} := \sum_{\substack{\beta \\ \beta \cdot E > 0}} N_{g,\beta}^{K_S} Q^\beta, \tag{43}$$

$$\bar{F}_g^{S/E} := \sum_{\substack{\beta \\ \beta \cdot E > 0}} \frac{(-1)^{\beta \cdot E + g - 1}}{\beta \cdot E} N_{g,\beta}^{S/E} Q^\beta, \tag{44}$$

$$F_{h,\mathbf{a}}^{E,tw} := -\frac{\delta_{h,1} \delta_{n,0}}{24} \log((-1)^{E \cdot E} \tilde{Q}) + \sum_{d_E \geq 0} N_{h,d_E,\mathbf{a}}^{E,tw} ((-1)^{E \cdot E} \tilde{Q})^{d_E}, \tag{45}$$

where the variable \tilde{Q} in $F_{h,\mathbf{a}}^{E,tw}$ is related to the variable Q in $\bar{F}_g^{S/E}$ and $\bar{F}_g^{K_S}$ by the formula (5).

Proposition 3.2.

$$\bar{F}_g^{K_S} = (-1)^g \bar{F}_g^{S/E} + \frac{\delta_{g,1}}{24} \log Q^E + \sum_{n \geq 0} \sum_{\substack{g=h+g_1+\dots+g_n, \\ \mathbf{a}=(a_1,\dots,a_n) \in \mathbb{Z}_{\geq 0}^n \\ (a_j, g_j) \neq (0,0), \sum_{j=1}^n a_j \leq 2h-2}} \frac{F_{h,\mathbf{a}}^{E,tw}}{|\text{Aut}(\mathbf{a}, \mathbf{g})|} \prod_{j=1}^n (-1)^{g_j-1} D^{a_j+2} \bar{F}_{g_j}^{S/E}.$$

Proof. Given $\mathbf{g} = (g_1, \dots, g_n)$ and $\mathbf{a} = (a_1, \dots, a_n)$, we have a disjoint sum decomposition

$$\{1, \dots, n\} = I_{\mathbf{g},\mathbf{a}} \bigsqcup J_{\mathbf{g},\mathbf{a}},$$

where $I_{\mathbf{g},\mathbf{a}}$ is the subset of j such $(g_j, a_j) \neq (0, 0)$ and $J_{\mathbf{g},\mathbf{a}}$ is the subset of j such that $(g_j, a_j) = (0, 0)$. Denote $\mathbf{a}' = (a_j)_{j \in I_{\mathbf{g},\mathbf{a}}}$. If $a_j = 0$, then there is no insertion of ψ_j in $N_{h,d_E,\mathbf{a}}^{E,tw}$, we can remove $\text{ev}_j^*(\omega)$ from the integral using the divisor equation, so we have $N_{h,d_E,\mathbf{a}}^{E,tw} = (\prod_{j \in J_{\mathbf{g},\mathbf{a}}} d_E) N_{h,d_E,\mathbf{a}'}^{E,tw}$. There is one exception: we cannot apply the divisor equation if $d_E = 0, n = 1, h = 0, g_1 = 0$ and $a_1 = 0$, in which case

$$N_{1,0,(0)}^{E,tw} = \int_{[\overline{\mathcal{M}}_{1,1}(E,0)]^{\text{vir}}} \text{ev}^*(\omega) = -\frac{1}{24}.$$

It follows that the correct general relation is

$$N_{h,d_E,\mathbf{a}}^{E,tw} = -\frac{\delta_{h,1} \delta_{n,1} \delta_{\mathbf{a},(0)}}{24} + N_{h,d_E,\mathbf{a}'}^{E,tw} \prod_{j \in J_{\mathbf{g},\mathbf{a}}} d_E. \tag{46}$$

It follows that we can replace the sum over \mathbf{a} by a sum over \mathbf{a}' .

After summing over β to form generating series, the factors indexed by $j \in J_{g,\mathbf{a}}$ are absorbed in $F_{h,\mathbf{a}}^{E,\text{tw}}$ via the change of variables $Q \mapsto \tilde{Q}$. Indeed, according to formula 7, we have

$$\tilde{Q} = (-1)^{E \cdot E} \exp(-D^2 F_0^{S/E}) = (-1)^{E \cdot E} Q^E \exp(-D^2 \bar{F}_0^{S/E}),$$

so

$$((-1)^{E \cdot E} \tilde{Q})^{d_E} = Q^{d_E} \sum_{l \geq 0} \frac{1}{l!} (-1)^l (d_E D^2 \bar{F}_0^{S/E})^l.$$

□

Recall from (2) and (3) that

$$F_g^{K_S} = -\frac{1 - \delta_{(E \cdot E),0}}{3!(E \cdot E)^2} (\log Q^E)^3 \delta_{g,0} + \left(\frac{1 - \delta_{(E \cdot E),0}}{(E \cdot E)} \frac{\chi(S)}{24} - \frac{1}{24} \right) (\log Q^E) \delta_{g,1} + \bar{F}_g^{K_S}$$

and

$$F_g^{S/E} = -\frac{1 - \delta_{(E \cdot E),0}}{3!(E \cdot E)^2} (\log Q^E)^3 \delta_{g,0} - \frac{1 - \delta_{(E \cdot E),0}}{(E \cdot E)} \frac{\chi(S)}{24} (\log Q^E) \delta_{g,1} + \bar{F}_g^{S/E}.$$

Proposition 3.3.

$$F_g^{K_S} = (-1)^g F_g^{S/E} +$$

$$\sum_{n \geq 0} \sum_{\substack{g=h+g_1+\dots+g_n, \\ \mathbf{a}=(a_1,\dots,a_n) \in \mathbb{Z}_{\geq 0}^n \\ (a_j, g_j) \neq (0,0), \sum_{j=1}^n a_j \leq 2h-2}} \frac{F_{h,\mathbf{a}}^{E,\text{tw}}}{|\text{Aut}(\mathbf{a}, \mathbf{g})|} \prod_{j=1}^n (-1)^{g_j-1} \left(D^{a_j+2} F_{g_j}^{S/E} + (E \cdot E) \delta_{g_j,0} \delta_{a_j,1} \right).$$

Proof. We rewrite Proposition 3.2 in terms of the series $F_g^{K_S}$ and $F_g^{S/E}$. One needs to use that $F_1^{K_S} - \bar{F}_1^{K_S} = -(F_1^{S/E} - \bar{F}_1^{S/E}) - \frac{1}{24} \log Q^E$ and $D^{a+2} \bar{F}_0^{S/E} = D^{a+2} F_0^{S/E} + (E \cdot E) \delta_{a,1}$ for $a \geq 1$. □

3.2. Twisted Gromov-Witten theory of the elliptic curve

In this section, we compute the twisted Gromov-Witten series $F_{g,\mathbf{a}}^{E,\text{tw}}$ of the elliptic curve in terms of the untwisted Gromov-Witten series $F_{g,\mathbf{a}}^E$. Recall from (45) that

$$F_{g,\mathbf{a}}^{E,\text{tw}} = -\frac{\delta_{g,1} \delta_{n,0}}{24} \log((-1)^{E \cdot E} \tilde{Q}) + \bar{F}_{g,\mathbf{a}}^{E,\text{tw}}$$

where

$$\bar{F}_{g,\mathbf{a}}^{E,\text{tw}} := \sum_{d_E \geq 0} \frac{((-1)^{E \cdot E} \tilde{Q})^{d_E}}{t^{\sum_{j=1}^n a_j}} \int_{[\overline{\mathcal{M}}_{g,n}(E, d_E)]^{\text{vir}}} \left(\prod_{j=1}^n \text{ev}_j^*(\omega) \psi_j^{a_j} \right) e_{\mathbb{C}^*}(-R^* \pi_* f^* N),$$

and from (4) that

$$F_{g,\mathbf{a}}^E = -\frac{\delta_{g,1} \delta_{n,0}}{24} \log((-1)^{E \cdot E} \tilde{Q}) + \bar{F}_{g,\mathbf{a}}^E$$

where

$$\bar{F}_{g,\mathbf{a}}^E := \sum_{d_E \geq 0} \tilde{Q}^{d_E} \int_{[\overline{\mathcal{M}}_{g,n}(E, d_E)]^{\text{vir}}} \prod_{j=1}^n \text{ev}_j^*(\omega) \psi_j^{a_j}.$$

Proposition 3.4. For every $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n$ such that $\sum_{j=1}^n a_j \leq 2g - 2$, the twisted Gromov-Witten theory of the elliptic curve is related with the untwisted one via

$$\bar{F}_{g,\mathbf{a}}^{E,\text{tw}} = (-1)^{g-1} \frac{(E \cdot E)^m}{m!} \bar{F}_{g,(\mathbf{a}, 1^m)}^E$$

where $m := 2g - 2 - \sum_{j=1}^n a_j$ and $(\mathbf{a}, 1^m) = (a_1, \dots, a_n, \underbrace{1, \dots, 1}_m)$.

The proof of Proposition 3.4 takes the remainder of this section.

Let W be a rank r vector bundle over a projective variety X . Let ρ_i be the Chern roots of W . We fix a fiberwise action of \mathbb{C}^* on W , and we denote by λ_i the corresponding equivariant parameters.

Coates and Givental [16] have expressed in terms of the Givental’s quantization formalism the computation of the W -twisted Gromov-Witten theory in terms of the untwisted one obtained by applying the Grothendieck-Riemann-Roch theorem to the universal curve over moduli spaces of stable maps [63, 23].

More precisely, according to the main result¹⁵ of [16], (the stable part of) the generating function of the W -twisted Gromov-Witten invariants

$$\int_{[\overline{\mathcal{M}}_{g,n}(X,d)]^{\text{vir}}} \left(\prod_{j=1}^n \text{ev}_j^*(\tau_j) \psi_j^{a_j} \right) e_{\mathbb{C}^*}(-R^* \pi_* f^* W)$$

can be computed via the quantization of the symplectic operator

$$\Delta(z) := \prod_{i=1}^r \sqrt{\lambda_i + \rho_i} \exp \left(- \sum_{m>0} \frac{B_{2m}}{2m(2m-1)} \frac{z^{2m-1}}{(\lambda_i + \rho_i)^{2m-1}} \right)$$

together with the quantization of the symplectic operator

$$\Psi(z) := \prod_{i=1}^r \exp \left(- \frac{\rho_i \ln \lambda_i}{z} + \frac{1}{z} \sum_{k>0} \frac{(-1)^{k-1} \rho_i^{k+1}}{k(k+1) \lambda_i^k} \right)$$

acting on the untwisted Gromov-Witten potential.

For the case relevant to Proposition 3.4, W is the rank 2 vector bundle $N = N_{E/S} \oplus N_{E/S}^\vee$, and we have

$$\lambda_1 = t, \quad \rho_1 = (E \cdot E) \omega, \quad \lambda_2 = -t, \quad \rho_2 = -(E \cdot E) \omega.$$

In this case, the expressions for the symplectic operators become much simpler. Indeed, most of the terms are odd under $\rho_i \mapsto -\rho_i, \lambda_i \mapsto -\lambda_i$, so most of the terms with $i = 1$ cancel pairwise with the terms with $i = 2$. We have

$$\Psi(z) = \exp \left(\frac{\log(-1)(E \cdot E) \omega}{z} \right),$$

and $\Delta(z)$ does not depend on z :

$$\Delta(z) = \sqrt{-1} (t + (E \cdot E) \omega).$$

Using the divisor equation, we see that the operator Ψ acts like the change of variables (see also Remarks under Theorem 1’ in [16])

$$\tilde{\mathcal{Q}} \mapsto \tilde{\mathcal{Q}} e^{\log(-1)(E \cdot E) \int_E \omega} = (-1)^{(E \cdot E)} \tilde{\mathcal{Q}}.$$

¹⁵ Theorem 1 of [16] computes the theory twisted by an arbitrary multiplicative characteristic class $\exp(\sum s_k \text{ch}_k)$, and Corollary 1 of [16] describes the twist by $e_{\mathbb{C}^*}(R^* \pi_* f^* W)$. In order to get the twist by $e_{\mathbb{C}^*}(-R^* \pi_* f^* W) = 1/e_{\mathbb{C}^*}(R^* \pi_* f^* W)$, we have to flip the sign of the coefficients s_k .

The action of $\Delta(z)$ can be viewed as an R -matrix action on the CohFT defined via the Gromov-Witten theory of elliptic curve.

We recall the definition of the R -matrix (quantization) action (see also [31], [69]; see also Section 2 of [13]). Suppose we have two symplectic vector spaces V, V' of the same dimension, with units $1, 1'$, and a symplectic transformation $R(z) \in \text{Hom}(V, V')[[z]]$. Here, by the symplectic condition $R(z)R^*(-z) = I$, we see that $R(z)$ is invertible and

$$R^{-1}(z) = R^*(-z).$$

For each $2g - 2 + n > 0$, let $F_{g,n}(-) = \langle - \rangle_{g,n}$ be the (given) correlation functions with insertions in $V[[z]]$. We define $\hat{R}F_{g,n}$ by the following graph sum

$$\hat{R}F_{g,n}(\tau_1 \psi_1^{k_1}, \dots, \tau_n \psi_n^{k_n}) := \sum_{\Gamma \in G_{g,n}} \frac{1}{\text{Aut}(\Gamma)} \text{Cont}_{\Gamma},$$

where $G_{g,n}$ is the set of stable graphs with genus g and n legs, and the contributions Cont_{Γ} are defined via the following construction:

- At each leg l of Γ , we place an insertion

$$R^{-1}(\psi_l) \tau_l \psi_l^{k_l};$$

- At each edge $e = (v_1, v_2)$ of Γ , we place a bi-vector as a two-direction insertion

$$\frac{\sum_{\alpha} e_{\alpha} \otimes e^{\alpha} - \sum_{\alpha} R^{-1}(\psi_{(e,v_1)}) e'_{\alpha} \otimes R^{-1}(\psi_{(e,v_2)}) e'^{\alpha}}{\psi_{(e,v_1)} + \psi_{(e,v_2)}},$$

where $\{e_{\alpha}\}, \{e'_{\alpha}\}$ are arbitrary bases of V, V' , with the dual basis $\{e^{\alpha}\}$ and $\{e'^{\alpha}\}$, respectively.

- At each vertex v of Γ , we place the map

$$\tau'_1 \psi_1^{l_1} \otimes \dots \otimes \tau'_{n_v} \psi_{n_v}^{l_{n_v}} \mapsto \sum_{k \geq 0} \frac{1}{k!} F_{g_v, n_v+k}(\tau'_1 \psi_1^{l_1}, \dots, \tau'_{n_v} \psi_{n_v}^{l_{n_v}}, T(\psi_{n_v+1}), \dots, T(\psi_{n_v+k})),$$

where $(\tau'_i, l_i) \in \{(\tau_1, k_1), \dots, (\tau_n, k_n)\}$, and $T(z) := z\mathbf{1} - zR^{-1}(z)\mathbf{1}'$.

In our case, $V = H^{\bullet}(E)(t)$ with pairing $(\alpha, \beta) = \int_E \alpha \wedge \beta$, $e_{\mathbb{C}^*}(N) = (\lambda_1 + \rho_1)(\lambda_2 + \rho_2) = -(t + (E \cdot E)\omega)^2$, and $V' = H^{\bullet}(E)(t)$ with pairing $(\alpha, \beta)' = \int_E \alpha \wedge \beta \wedge e_{\mathbb{C}^*}^{-1}(N) = \int_E \frac{\alpha}{\sqrt{-1}(t+(E \cdot E)\omega)} \wedge \frac{\beta}{\sqrt{-1}(t+(E \cdot E)\omega)}$. The units $\mathbf{1}, \mathbf{1}'$ are both equal to $1 \in H^0(E)$. The R -matrix $R(z) = \Delta$ is independent of z , so the edge bi-vector is simply zero. Hence the contribution is non-vanishing only if the graph is a single vertex. Namely

$$\langle \tau_1 \psi_1^{a_1}, \dots, \tau_n \psi_n^{a_n} \rangle_{g,n}^{E, \text{tw}} = \sum_{k \geq 0} \frac{1}{k!} \langle R^{-1}(\psi_1) \tau_1 \psi_1^{a_1}, \dots, R^{-1}(\psi_n) \tau_n \psi_n^{a_n}, T(\psi_{n+1}), \dots, T(\psi_{n+k}) \rangle_{g, n+k}^E \Big|_{\tilde{Q} \mapsto (-1)^{(E \cdot E)} \tilde{Q}}$$

where $\tau_j \in H^\bullet(E)$,

$$\langle \tau_1 \psi_1^{a_1}, \dots, \tau_n \psi_n^{a_n} \rangle_{g,n}^{E,tw} := \sum_{d_E \geq 0} \tilde{Q}^{d_E} \int_{[\overline{\mathcal{M}}_{g,n}(E,d_E)]^{\text{vir}}} \left(\prod_{j=1}^n \psi_j^{a_j} \text{ev}_j^*(\tau_j) \right) e_{\mathbb{C}^*}(-R^* \pi_* f^* N),$$

$$\langle \tau_1 \psi_1^{a_1}, \dots, \tau_n \psi_n^{a_n} \rangle_{g,n}^E := \sum_{d_E \geq 0} \tilde{Q}^{d_E} \int_{[\overline{\mathcal{M}}_{g,n}(E,d)]^{\text{vir}}} \prod_{j=1}^n \psi_j^{a_j} \text{ev}_j^*(\tau_j).$$

We have

$$R(z) = \Delta(z) = \sqrt{-1}(t + (E \cdot E)\omega),$$

$$R^{-1}(z) = \frac{1}{\sqrt{-1}t} - \frac{(E \cdot E)\omega}{\sqrt{-1}t^2},$$

$$T(z) = z \left(1 - \frac{1}{\sqrt{-1}t} + \frac{(E \cdot E)\omega}{\sqrt{-1}t^2} \right).$$

We focus on the case where all the insertions τ_i are point classes ω . We have

$$\langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n}^{E,tw} = \sum_{k \geq 0} \frac{1}{k!} \langle R^{-1}(\psi_1)\omega \psi_1^{a_1}, \dots, R^{-1}(\psi_n)\omega \psi_n^{a_n}, T(\psi_{n+1}), \dots, T(\psi_{n+k}) \rangle_{g,n+k}^E |_{\tilde{Q} \mapsto (-1)^{(E \cdot E)} \tilde{Q}}.$$

For every j , we have $R^{-1}(\psi_j)\omega = \frac{1}{\sqrt{-1}t}\omega$, so

$$\langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n}^{E,tw} = \frac{1}{(\sqrt{-1}t)^n} \sum_{k \geq 0} \frac{1}{k!} \langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n}, T(\psi_{n+1}), \dots, T(\psi_{n+k}) \rangle_{g,n+k}^E |_{\tilde{Q} \mapsto (-1)^{(E \cdot E)} \tilde{Q}}.$$

Writing

$$T(\psi_j) = \left(1 - \frac{1}{\sqrt{-1}t} \right) \psi_j + \frac{(E \cdot E)}{\sqrt{-1}t^2} \psi_j \omega$$

and expanding, we obtain

$$\langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n}^{E,tw} = \frac{1}{(\sqrt{-1}t)^n} \sum_{m,l \geq 0} \frac{1}{m!l!} \left(\frac{(E \cdot E)}{\sqrt{-1}t^2} \right)^m \left(1 - \frac{1}{\sqrt{-1}t} \right)^l \times \langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n}, \omega \psi_{n+m}, \psi_{n+m+1}, \dots, \psi_{n+m+l} \rangle_{g,n+m+l}^E |_{\tilde{Q} \mapsto (-1)^{(E \cdot E)} \tilde{Q}}.$$

The sum over l can be evaluated using the dilaton equation

$$\begin{aligned} & \sum_{l \geq 0} \frac{1}{l!} \left(1 - \frac{1}{\sqrt{-1}t}\right)^l \langle -, \psi_{n+m+1}, \dots, \psi_{m+m+1} \rangle_{g,n+m+1}^E \\ &= \sum_{l \geq 0} \binom{-(2g-2+n+m)}{l} \left(\frac{1}{\sqrt{-1}t} - 1\right)^l \langle - \rangle_{g,n+m}^E \\ &= \left(\frac{1}{\sqrt{-1}t}\right)^{-(2g-2+n+m)} \langle - \rangle_{g,n+m}^E . \end{aligned}$$

Notice that in the second equality we have used the following generalized Newton’s binomial theorem:

$$\sum_{k > 0} \binom{\alpha}{k} x^k = (1+x)^\alpha .$$

Hence, collecting the powers of $\sqrt{-1}$ and t , we obtain

$$\begin{aligned} \langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n}^{E,tw} &= (\sqrt{-1})^{2g-2} \times \\ & \sum_{m \geq 0} \frac{(E \cdot E)^m}{m!} t^{2g-2-m} \langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n}, \omega \psi_{n+1}, \dots, \omega \psi_{n+m} \rangle_{g,n+m}^E |_{\tilde{Q} \mapsto (-1)^{(E \cdot E)} \tilde{Q}} . \end{aligned}$$

By dimension constraint, the correlator in the sum is non-vanishing only if $m = 2g - 2 - \sum_{j=1}^n a_j$. Therefore, we have

$$\begin{aligned} \langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n}^{E,tw} &= (-1)^{g-1} t^{\sum_{j=1}^n a_j} \frac{(E \cdot E)^m}{m!} \times \\ & \langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n}, \omega \psi_{n+1}, \dots, \omega \psi_{n+m} \rangle_{g,n+m}^E |_{\tilde{Q} \mapsto (-1)^{(E \cdot E)} \tilde{Q}} \end{aligned}$$

where $m = 2g - 2 - \sum_{j=1}^n a_j$. Note that the term $t^{\sum_{j=1}^n a_j}$ cancels with the same term in the denominator of $\bar{F}_{g,\mathbf{a}}^{E,tw}$. This concludes the proof of Proposition 3.4.

3.3. Conclusion of the proof of Theorem 1.2

By Proposition 3.3, we have

$$F_g^{KS} = (-1)^g F_g^{S/E} +$$

$$\sum_{\substack{n \geq 0 \\ (a_j, g_j) \neq (0,0), \sum_{j=1}^n a_j \leq 2h-2}} \sum_{\substack{g=h+g_1+\dots+g_n, \\ \mathbf{a}=(a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n}} \frac{1}{|\text{Aut}(\mathbf{a}, \mathbf{g})|} F_{h,\mathbf{a}}^{E,tw} \prod_{j=1}^n (-1)^{g_j-1} \left(D^{a_j+2} \bar{F}_{g_j}^{S/E} + (E \cdot E) \delta_{g_j,0} \delta_{a_j,1} \right) .$$

For every $\mathbf{g} = (g_j)_j$ and $\mathbf{a} = (a_j)_j$ such that $(a_j, g_j) \neq (0, 0)$ for every j and $\sum_j a_j \leq 2h - 2$, we denote $m = 2h - 2 - \sum_{j=1}^n a_j$, and we define

$$\tilde{\mathbf{g}} := (g_1, \dots, g_n, 0, \dots, 0)$$

and

$$\bar{\mathbf{a}} := (a_1, \dots, a_n, \underbrace{1, \dots, 1}_m).$$

We have $(\bar{g}_j, \bar{a}_j) \neq (0, 0)$ for every j and $\sum_j \bar{a}_j = 2h - 2$.

According to Proposition 3.4, we have

$$F_{h,\mathbf{a}}^{E,\text{tw}} = (-1)^{h-1} \frac{(E \cdot E)^m}{m!} F_{h,\bar{\mathbf{a}}}^E.$$

Therefore, we can rewrite the sum over $\mathbf{a} = (a_j)_j$ with $\sum_j a_j \leq 2h - 2$ as a sum over $\mathbf{a} = (\bar{a}_j)_j$ with $\sum_j \bar{a}_j = 2h - 2$. More precisely, let $\bar{\mathbf{g}} = (\bar{g}_j)_j$ and $\bar{\mathbf{a}} = (\bar{a}_j)_j$ be of the form

$$\begin{aligned} \bar{\mathbf{g}} &= (\bar{g}_1, \dots, \bar{g}_k, \underbrace{0, \dots, 0}_l) \\ \bar{\mathbf{a}} &= (\bar{a}_1, \dots, \bar{a}_k, \underbrace{1, \dots, 1}_l) \end{aligned}$$

with $(\bar{a}_j, \bar{g}_j) \neq (0, 0), (1, 0)$ for $1 \leq j \leq k, g = h + \sum_{j=1}^k \bar{g}_j$ and $\sum_{j=1}^{k+l} \bar{a}_j = 2h - 2$. The total contribution of $(\bar{\mathbf{a}}, \bar{\mathbf{g}})$ in the rewritten formula is the sum of contributions of (\mathbf{a}, \mathbf{g}) with $(\bar{\mathbf{a}}, \bar{\mathbf{g}}) = (\bar{\mathbf{a}}, \bar{\mathbf{g}})$. Such (\mathbf{a}, \mathbf{g}) is of the form

$$\begin{aligned} \mathbf{g} &= (\bar{g}_1, \dots, \bar{g}_k, \underbrace{0, \dots, 0}_{l-m}) \\ \mathbf{a} &= (\bar{a}_1, \dots, \bar{a}_k, \underbrace{1, \dots, 1}_{l-m}) \end{aligned}$$

with $0 \leq m \leq l$. Using that

$$|\text{Aut}(\mathbf{a}, \mathbf{g})| = |\text{Aut}(\{(\bar{a}_1, \bar{g}_1), \dots, (\bar{a}_k, \bar{g}_k)\})|(l - m)!,$$

we get that the total contribution of $(\bar{\mathbf{a}}, \bar{\mathbf{g}})$ is given by

$$\begin{aligned} &\frac{(-1)^{h-1} F_{h,\bar{\mathbf{a}}}^E}{|\text{Aut}(\{(\bar{a}_1, \bar{g}_1), \dots, (\bar{a}_k, \bar{g}_k)\})|} \left(\prod_{j=1}^k (-1)^{\bar{g}_j-1} D^{\bar{a}_j+1} F_{\bar{g}_j}^{S/E} \right) \times \\ &\sum_{m=0}^l (-1)^{l-m} \frac{(E \cdot E)^m (D^3 F_0^{S/E} + (E \cdot E))^{l-m}}{m!(l-m)!}. \end{aligned}$$

Using the binomial theorem, this can be rewritten as

$$\frac{(-1)^{h-1} F_{h,\bar{\mathbf{a}}}^E}{|\text{Aut}(\{(\bar{a}_1, \bar{g}_1), \dots, (\bar{a}_k, \bar{g}_k)\})|} \left(\prod_{j=1}^k (-1)^{\bar{g}_j-1} D^{\bar{a}_j+1} F_{\bar{g}_j}^{S/E} \right) (-1)^l \frac{(D^3 F_0^{S/E})^l}{l!}.$$

As $|\text{Aut}(\bar{\mathbf{a}}, \bar{\mathbf{g}})| = |\text{Aut}(\{(\bar{a}_1, \bar{g}_1), \dots, (\bar{a}_k, \bar{g}_k)\})|l!$, we finally obtain

$$\frac{(-1)^{h-1} F_{h,\bar{\mathbf{a}}}^E}{|\text{Aut}(\bar{\mathbf{a}}, \bar{\mathbf{g}})|} \left(\prod_{j=1}^{k+l} (-1)^{\bar{g}_j-1} D^{\bar{a}_j+1} F_{\bar{g}_j}^{S/E} \right).$$

This ends the proof of Theorem 1.2.

3.4. Stationary Gromov-Witten theory of the elliptic curve

According to Theorem 1.2, the local series $F_g^{K_S}$ and the relative series $F_g^{S/E}$ determine each other through the stationary series $F_{g,\mathbf{a}}^E$ of the elliptic curve. In this section, we review the computation by Okounkov and Pandharipande [67, §5] of the stationary Gromov-Witten theory of the elliptic curve.

Recall that, for every $(a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n$, we denote by

$$\bar{F}_{g,\mathbf{a}}^E := \sum_{d_E \geq 0} \tilde{Q}^{d_E} \int_{[\overline{\mathcal{M}}_{g,n}(E,d_E)]^{\text{vir}}} \prod_{j=1}^n \text{ev}_j^*(\omega) \psi_j^{a_j}$$

the generating series of stationary Gromov-Witten invariants of the elliptic curve E .

For every $k \geq 1$, we consider the Eisenstein series

$$E_{2k}(\tilde{\tau}) = 1 - \frac{4k}{B_{2k}} \sum_{n=1}^{\infty} \frac{n^{2k-1} \tilde{Q}^n}{1 - \tilde{Q}^n},$$

where $\tilde{Q} = e^{2i\pi\tilde{\tau}}$ and the Bernoulli numbers B_{2k} are defined by $\frac{t}{e^t-1} = \sum_{n \geq 0} B_n \frac{t^n}{n!}$. As functions on the upper half-plane $\{\tilde{\tau} \in \mathbb{C} \mid \text{Im } \tilde{\tau} > 0\}$, E_{2k} is modular of weight $2k$ for $SL(2, \mathbb{Z})$ for every $k \geq 2$, and E_2 is quasimodular of weight 2 for $SL(2, \mathbb{Z})$. The ring $\mathbb{C}[E_2, E_4, E_6]$, graded by the weight, is exactly the graded ring $\text{QMod}(SL(2, \mathbb{Z}))$ of quasimodular forms for $SL(2, \mathbb{Z})$ [74]. For every $k \geq 0$, we denote by $\mathbb{C}[E_2, E_4, E_6]_k$ the weight k subspace of $\mathbb{C}[E_2, E_4, E_6]$.

Theorem 3.5 (Okounkov-Pandharipande [67]). *For every $g \geq 0$, $n \geq 1$ and $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n$, we have*

$$\bar{F}_{g,\mathbf{a}}^E \in \mathbb{Q}[E_2, E_4, E_6]_{\sum_{j=1}^n (a_j+2)}.$$

In fact, Okounkov and Pandharipande give an explicit formula computing $\bar{F}_{g,\mathbf{a}}^E$ as a polynomial in E_2, E_4, E_6 .

For every $n \geq 1$, let

$$F(z_1, \dots, z_n) := \delta_{1,n} z_1^{-1} + \sum_{\substack{\mathbf{a}=(a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n, \\ 2g-2=\sum_{j=1}^n a_j}} \bar{F}_{g,\mathbf{a}}^E \cdot z_1^{a_1+1} \cdots z_n^{a_n+1},$$

and let $F^\bullet(z_1, \dots, z_n)$ be the disconnected generating function defined by¹⁶

$$F^\bullet(z_1, \dots, z_n) := \sum_{k > 0} \sum_{\substack{I_1 \sqcup \dots \sqcup I_k = \{1, \dots, n\} \\ I_j \neq \emptyset}} \frac{1}{|\text{Aut}(I_1, \dots, I_k)|} \prod_{j=1}^k F(z_{I_j}).$$

This relation can be inverted to compute the connected series $F(z_1, \dots, z_n)$ in terms of the disconnected ones.

We introduce the odd theta function

$$\vartheta(\tilde{\tau}, z) := \vartheta_{\frac{1}{2}, \frac{1}{2}}(\tilde{\tau}, z) = \sum_{k=-\infty}^{\infty} (-1)^k e^{(k+\frac{1}{2})z} e^{\pi i \tilde{\tau} (k+\frac{1}{2})^2},$$

¹⁶ The condition $I_j \neq \emptyset$ excludes genus-1 unmarked connected components, so only series $F(z_1, \dots, z_n)$ with $n \geq 1$ enter the formula. For $n = 0$, the series $-\frac{1}{24} + \bar{F}_{1,0}^E = -\log \eta$ is not quasimodular.

and following Bloch and Okounkov [7], we denote

$$\begin{aligned} \Theta(z) &:= \eta(\tilde{\tau})^{-3} \vartheta(\tilde{\tau}, z) \\ &= z \exp\left(\sum_{k=1}^{\infty} \frac{B_{2k}}{2k(2k)!} E_{2k}(\tilde{\tau}) z^{2k}\right). \end{aligned}$$

Theorem 3.6 (Okounkov-Pandharipande [67]). *For every $n \geq 1$, we have*

$$F^\bullet(z_1, \dots, z_n) = \sum_{\substack{\sigma \text{ permutation} \\ \text{of } z_1, \dots, z_n}} \frac{\det \left[\frac{\Theta^{(j-i+1)}(z_{\sigma(1)} + \dots + z_{\sigma(n-j)})}{(j-i+1)!} \right]_{i,j=1}^n}{\Theta(z_{\sigma(1)}) \Theta(z_{\sigma(1)} + z_{\sigma(2)}) \dots \Theta(z_{\sigma(1)} + \dots + z_{\sigma(n)})},$$

where $\Theta^{(k)}$ is the k -th derivative of Θ with respect to z and $\frac{\Theta^{(j-i+1)}}{(j-i+1)!}$ is interpreted as 0 if $j - i + 1 < 0$.

For example, using that $\Theta'(0) = 1$ and $\Theta''(0) = 0$, we get

$$\begin{aligned} F(z) = F^\bullet(z) &= \frac{1}{\Theta(z)} = z^{-1} - \frac{E_2}{24} z + \left(\frac{E_4}{2880} + \frac{E_2^2}{1152}\right) z^3 \\ &\quad - \left(\frac{E_6}{181440} + \frac{E_2 E_4}{69120} + \frac{E_2^3}{82944}\right) z^5 + \dots \\ F(z_1, z_2) &= F^\bullet(z_1, z_2) - F^\bullet(z_1) F^\bullet(z_2) \\ &= \frac{1}{\Theta(z_1 + z_2)} \left(\frac{\Theta'(z_1)}{\Theta(z_1)} + \frac{\Theta'(z_2)}{\Theta(z_2)}\right) - \frac{1}{\Theta(z_1)\Theta(z_2)} \\ &= -\frac{(E_2^2 - E_4) z_1 z_2}{288} + \frac{(5 E_2^3 - 3 E_2 E_4 - 2 E_6) z_1^2 z_2^2}{25920} \\ &\quad + \frac{(5 E_2^3 - E_2 E_4 - 4 E_6) (z_1 z_2^3 + z_1^3 z_2)}{34560} + \dots \end{aligned}$$

Hence, we obtain

$$F_{1,(0)}^E = -\frac{E_2}{24}, \tag{47}$$

$$F_{1,(0^2)} = -\frac{1}{12 \cdot 24} (E_2^2 - E_4), \tag{48}$$

$$F_{2,(1^2)}^E = -\frac{1}{9^2 \cdot 320} (2E_6 + 3E_2 E_4 - 5E_2^3), \tag{49}$$

$$F_{2,(2)}^E = \frac{1}{9 \cdot 640} (2E_4 + 5E_2^2). \tag{50}$$

4. Finite generation and quasimodularity for (\mathbb{P}^2, E)

4.1. Quasimodular forms for $\Gamma_1(3)$

We refer to [18, 74] for basics on modular and quasimodular forms. For every Γ congruence subgroup of $SL(2, \mathbb{Z})$, we denote by $\text{Mod}(\Gamma) = \bigoplus_{k \geq 0} \text{Mod}(\Gamma)_k$ the ring of modular forms for Γ , graded by the weight, and by $\text{QMod}(\Gamma) = \bigoplus_{k \geq 0} \text{QMod}(\Gamma)_k$ the ring of quasimodular forms for Γ , also graded by the weight.

We focus on the congruence subgroups $\Gamma_1(3)$ and $\Gamma_0(3)$ defined by

$$\Gamma_1(3) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}) \mid \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix} \pmod{3} \right\},$$

$$\Gamma_0(3) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}) \mid \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \pmod{3} \right\}.$$

We have $\Gamma_1(3) \subset \Gamma_0(3)$, subgroup of index 2, with $-I \in \Gamma_0(3)$ and $-I \notin \Gamma_1(3)$. It follows that the modular (respectively, quasimodular) forms for $\Gamma_0(3)$ are exactly the modular (respectively, quasimodular) forms for $\Gamma_1(3)$ of even weight.

Let

$$A(\tau) := \left(\frac{\eta(\tau)^9}{\eta(3\tau)^3} + 27 \frac{\eta(3\tau)^9}{\eta(\tau)^3} \right)^{\frac{1}{3}} = 1 + 6Q + 6Q^3 + 6Q^4 + 12Q^7 + 6Q^9 + \dots$$

$$C(\tau) := \frac{\eta(\tau)^9}{\eta(3\tau)^3} = 1 - 9Q + 27Q^2 - 9Q^3 - 117Q^4 + 216Q^5 + 27Q^6 - 450Q^7 + \dots$$

where $\eta(\tau) := Q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - Q^n)$ is the Dedekind eta function and $Q = e^{2i\pi\tau}$. It follows from the known modular properties of the eta function that A and C are modular forms for $\Gamma_1(3)$ respectively of weight 1 and 3. We refer to section 3 of [59] for details.¹⁷ The combination

$$\frac{A^3 - C}{27} = \frac{\eta(3\tau)^9}{\eta(\tau)^3} = Q + 3Q^2 + 9Q^3 + 13Q^4 + 24Q^5 + 27Q^6 + 50Q^7 + \dots$$

is a cusp form of weight 3 for $\Gamma_1(3)$.

Lemma 4.1. *The functions A and C are algebraically independent over \mathbb{C} .*

Proof. If there exists a non-trivial polynomial relation between A and C , then there exists a non-trivial polynomial relation between A and C , which is homogeneous for the weight and so of the form $\sum_{n+3m=k} a_{nm} A^n C^m$ for some k . If such a non-trivial relation existed, then, dividing by A^k , we would get that C/A^3 is the solution of a non-trivial polynomial equation with complex coefficients; so C/A^3 would be constant, which is not the case as $C/A^3 = 1 - 27Q + O(Q^2)$. \square

According to [18, Figure 3.4],¹⁸ the dimension of $\text{Mod}(\Gamma_1(3))_k$ is $\lfloor \frac{k}{3} \rfloor + 1$. Therefore, Lemma 4.1 implies¹⁹ that

$$\text{Mod}(\Gamma_1(3)) = \mathbb{C}[A, C].$$

By Proposition 20 of [74], the Eisenstein series $E_2(\tau)$ for $SL(2, \mathbb{Z})$ is algebraically independent of $\text{Mod}(\Gamma_1(3))$ over \mathbb{C} and

$$\text{QMod}(\Gamma_1(3)) = \text{Mod}(\Gamma_1(3))[E_2].$$

¹⁷ In terms of the notation \mathcal{A}_3 and \mathcal{B}_3 of [59], we have $A = \mathcal{A}_3$ and $C = \mathcal{B}_3^3$.

¹⁸ In [18], the formula is a priori valid only for $k \geq 2$. We get that $\dim \text{Mod}(\Gamma_1(3))_1 = 1$ because it is > 0 by the existence of A , and ≤ 1 as $\dim \text{Mod}(\Gamma_1(3))_2 = 1$.

¹⁹ Therefore, A is the unique weight 1 modular form for $\Gamma_1(3)$ with constant term 1 and can be described as the weight 1 Eisenstein series $E_1^{\psi,1} = 1 + 6 \sum_{d \geq 1} \sum_{d|n} \psi(d) Q^n$, where ψ is the non-trivial character of $(\mathbb{Z}/3)^* \cong \{\pm 1\}$ (extended by $\psi(d) = 0$ if $3|d$), or as the theta series of the hexagonal lattice $\mathbb{Z}[\sqrt{-3}]$. In particular, the coefficient of Q^n in A is the number of $(x, y) \in \mathbb{Z}^2$ such that $|x + e^{\frac{2i\pi}{3}} y|^2 = x^2 - xy + y^2 = n$ (see [18, Exercice 4.11.5]).

The difference $3E_2(3\tau) - E_2(\tau)$ is modular for $\Gamma_1(3)$. Using that the space of weight 2 modular forms for $\Gamma_1(3)$ is of dimension 1, we find that $3E_2(3\tau) - E_2(\tau) = 2A^2$. Therefore, we can use

$$B(\tau) := \frac{1}{4}(E_2(\tau) + 3E_2(3\tau)) = 1 - 6Q - 18Q^2 - 42Q^3 - 42Q^2 + \dots$$

instead of E_2 as depth 1 weight 2 generator, and we have

$$\text{QMod}(\Gamma_1(3)) = \mathbb{C}[A, B, C].$$

The ring $\mathbb{Q}[A, B, C]$ is closed under the differential operator

$$\partial_\tau := \frac{1}{2\pi i} \frac{d}{d\tau} = Q \frac{d}{dQ}.$$

Indeed, ∂_τ maps weight k quasimodular forms to weight $k + 2$ quasimodular forms by a straightforward calculation from the definition of quasimodularity, and in particular for $\text{QMod}(\Gamma_1(3)) = \mathbb{C}[A, B, C]$ this allows to show easily the Ramanujan-type identities

$$\begin{aligned} \partial_\tau A &= \frac{1}{6}A(B + A^2) - \frac{C}{3}, \\ \partial_\tau B &= \frac{1}{6}(B^2 - A^4), \quad \partial_\tau C = \frac{1}{2}C(B - A^2), \end{aligned} \tag{51}$$

because the quasi-modular properties of both sides are known, so it is enough to identify finitely many terms of the Q -expansions.

4.2. *Mirror of local \mathbb{P}^2 and quasimodular forms*

We review the relation between the mirror family of local \mathbb{P}^2 and modular forms, following [4, 76, 17].

Let $\mathbb{H} = \{\tau \in \mathbb{C} \mid \text{Im } \tau > 0\}$ be the upper half-plane. We consider the modular curve $Y_1(3) = [\mathbb{H}/\Gamma_1(3)]$. It is a smooth orbifold, whose coarse moduli space can be identified with $\{q \in \mathbb{C} \mid q \neq -\frac{1}{27}, 0\} \cup \{\infty\}$, and with a single $\mathbb{Z}/3$ -orbifold point at $q = \infty$. The modular curve $Y_1(3)$ has two cusps, given by $q = 0$ and $q = -\frac{1}{27}$ (corresponding, respectively, to the $\Gamma_1(3)$ -equivalence classes of $\tau = i\infty$ and $\tau = 0$). In the context of mirror symmetry, where $Y_1(3)$ is viewed as the stringy Kähler moduli space of local \mathbb{P}^2 , the point $q = 0$ is the large volume point, $q = -\frac{1}{27}$ is the conifold point and $q = \infty$ is the orbifold point.

The coordinate q on $Y_1(3)$ is expressed in terms of τ [58]²⁰ by

$$\frac{1}{1 + 27q} = 1 + 27 \frac{\eta(3\tau)^{12}}{\eta(\tau)^{12}}, \tag{52}$$

that is, denoting $X := (1 + 27q)^{-1}$, by

$$X = \frac{A^3}{C}. \tag{53}$$

The periods of the universal family of elliptic curves with $\Gamma_1(3)$ -level structure are solution of the Picard-Fuchs equation [59]

$$\left[\left(q \frac{d}{dq} \right)^2 - 3q \left(3 \left(q \frac{d}{dq} \right) + 1 \right) \left(3 \left(q \frac{d}{dq} \right) + 2 \right) \right] \Pi = 0. \tag{54}$$

²⁰ The description of the Hauptmodul of the genus 0 modular curve $Y_1(3)$ in terms of eta functions goes back to Klein and Fricke at the end of the 19th century.

In Section 1.3, we defined the functions $S, I_{11} = q \frac{dI_1}{dq}, I_{12} = q \frac{dI_2}{dq}$ in terms of solutions I_1 and I_2 of the differential equation (11) describing genus 0 mirror symmetry for local \mathbb{P}^2 . As the differential equation (11) is obtained from (54) by applying $q \frac{d}{dq}$ and flipping the sign of q , we deduce that τ , viewed as a multivalued function of q , is given by

$$\tau = \frac{1}{2} + \frac{1}{2\pi i} \frac{I_{12}(q)}{I_{11}(q)}, \tag{55}$$

that is,

$$\mathcal{Q} = e^{2\pi i \tau} = -\exp\left(\frac{I_{12}(q)}{I_{11}(q)}\right). \tag{56}$$

One should not confuse the variables q, \mathcal{Q}, Q :

- q is such that $(1 + 27q)^{-1}$ is a globally defined coordinate on $Y_1(3)$,
- $\mathcal{Q} = e^{2i\pi\tau}$ is the flat modular coordinate for the family of elliptic curves parametrized by $Y_1(3)$,
- Q is the flat coordinate determined by the mirror of local \mathbb{P}^2 .

The variables q and \mathcal{Q} are related by (52)–(56). The variables q and Q are related by the mirror transformation (12).

According to [4, 58, 59, 76], the functions X, I_{11}, S and the quasimodular forms A, B, C determine each other through the identities

$$A = I_{11}, \quad B = \frac{I_{11}^2}{X}(X + 6S), \quad C = \frac{I_{11}^3}{X}, \tag{57}$$

of inverse

$$X = \frac{A^3}{C}, \quad I_{11} = A, \quad S = \frac{1}{6} \frac{AB - A^3}{C}. \tag{58}$$

In particular, viewed as functions of τ , X is a modular function of weight 0 for $\Gamma_1(3)$, I_{11} is a modular form of weight 1 for $\Gamma_1(3)$ and S is a quasimodular function of weight 0 for $\Gamma_1(3)$.

Lemma 4.2. ²¹ *The functions S, X, I_{11} are algebraically independent over \mathbb{C} .*

Proof. It is a direct corollary of the algebraic independence of the quasimodular forms A, B, C reviewed in Section 4.1. □

By Lemma 4.2, the ring of functions generated by S and X is the polynomial ring

$$\mathbf{R} := \mathbb{Q}[X, S].$$

We define a grading on \mathbf{R} by $\deg X = \deg S = 1$ and denote by $\mathbf{R}_{\leq k}$ the subspace of polynomials with degree no more than k .

We consider the vector space $[X^{-(g-1)} \cdot \mathbf{R}_{\leq 3g-3}]^{\text{reg}}$, where $[-]^{\text{reg}}$ (the ‘orbifold regularity’ condition) is defined by

$$[-]^{\text{reg}} := \{f(X, S) : 3 \deg_X f + \deg_S f \geq 0\}. \tag{59}$$

Proposition 4.3. *The expression of S and X in terms of the quasimodular forms A, B, C induces an identification*

$$[X^{-(g-1)} \cdot \mathbf{R}_{\leq 3g-3}]^{\text{reg}} = C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}$$

for every $g \geq 2$.

²¹ In [51], the algebraic independence of S, X, I_{11} is not considered as known, and explicit lifts of functions to the ring $\mathbb{Q}[S, X]$ are constructed. This extra work is not necessary by Lemma 4.2.

Proof. We first prove that $[X^{-(g-1)} \cdot \mathbf{R}_{\leq 3g-3}]^{\text{reg}} \subset C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}$. Let $X^{-(g-1)} \cdot X^j S^k \in [X^{-(g-1)} \cdot \mathbf{R}_{\leq 3g-3}]^{\text{reg}}$. According to (58), the C -degree of $X^{-(g-1)} \cdot X^j S^k$ is $-j-k+(g-1) \geq -(2g-2)$, using that $j+k \leq 3g-3$ by definition of $\mathbf{R}_{\leq 3g-3}$. The B -degree is $k \geq 0$. The A -degree of each term is $\geq -(3g-3)+3j+k$, which is ≥ 0 by definition of $[-]^{\text{reg}}$. Therefore, $X^{-(g-1)} \cdot X^j S^k = C^{-(2g-2)} f(A, B, C)$ for some $f \in \mathbb{Q}[A, B, C]$. As $X^{-(g-1)} \cdot X^j S^k$ is of weight 0 and C of weight 3, f is of weight $6g-6$.

We prove that conversely, $C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6} \subset [X^{-(g-1)} \cdot \mathbf{R}_{\leq 3g-3}]^{\text{reg}}$. Let $C^{-(2g-2)} \cdot A^m B^n C^l \in C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}$. According to (57), the power of I_{11} is equal to the weight. As $C^{-(2g-2)} \cdot A^m B^n C^l$ is of weight 0, $C^{-(2g-2)} \cdot A^m B^n C^l$ is independent of I_{11} and is only a function of S and X . The S -degree of $C^{-(2g-2)} \cdot A^m B^n C^l$ is $n \geq 0$. The X -degree of each term in $C^{-(2g-2)} \cdot A^m B^n C^l$ is $\geq (2g-2) - n - l$ and $\leq (2g-2) - l$. Therefore, $\deg_X + \deg_S \leq (2g-2) - l \leq 2g-2$. On the other hand, $3 \deg_X + \deg_S \geq (6g-6) - 3n - 3l + n = (6g-6) - 2n - 3l$, which is ≥ 0 as $A^m B^n C^l$ is of weight $6g-6$, B is of weight 2 and C is of weight 3. □

Remark 4.4. It follows from the proof of Proposition 4.3 that the constraint defined by $[-]^{\text{reg}}$ is equivalent to the absence of negative powers of A . According to [59], A has a zero at the orbifold point $q = \infty$. Therefore, $[-]^{\text{reg}}$ is indeed the condition imposed by the regularity at the orbifold point.

We consider the differential operator

$$D = 3Q \frac{d}{dQ} .$$

Lemma 4.5.

$$D = 3I_{11}^{-1} \cdot q \frac{d}{dq} \tag{60}$$

$$= 3C^{-1} \cdot Q \frac{d}{dQ} . \tag{61}$$

Proof. The equality (60) is clear as $Q = e^{I_1(q)}$ by (12) and (15).

In order to prove (61), we use (56):

$$Q \frac{d}{dQ} = I_{22}^{-1} q \frac{d}{dq}$$

where

$$I_{22} := q \frac{d}{dq} \left(\frac{I_{12}(q)}{I_{11}(q)} \right) .$$

Theorem 2 in [75] shows that $I_{22} = \frac{X}{I_{11}^2}$. We conclude using that $C = \frac{I_{11}^3}{X}$ according to (57). □

Lemma 4.6. For every $n, k \geq 0$, we have

$$D(C^{-n} \cdot \mathbb{Q}[A, B, C]_k) \subset C^{-(n+1)} \cdot \mathbb{Q}[A, B, C]_{k+2} .$$

Proof. This follows from (61) and (51). The important point is that $\partial_\tau C$ is divisible by C . □

Lemma 4.7. We have

$$q \frac{d}{dq} S = -S^2 + \frac{X-1}{3} S - \frac{X(X-1)}{9} , \quad q \frac{d}{dq} X = X(X-1) . \tag{62}$$

Proof. The formula for X is clear as $X = (1 + 27q)^{-1}$ by definition.

The formula for S follows from the differential equation (11). Alternatively, one can use the expression (58) in terms of quasimodular forms and (51)–(61). □

4.3. Quasimodular forms: from $SL(2, \mathbb{Z})$ to $\Gamma_1(3)$

As reviewed in Section 3.4, the ring of quasimodular forms for $SL(2, \mathbb{Z})$ is generated by the Eisenstein series $E_2(\tilde{\tau}), E_4(\tilde{\tau}), E_6(\tilde{\tau})$. On the other hand, we have seen in Section 4.1 that the ring of quasimodular forms for $\Gamma_1(3)$ is generated by the functions $A(\tau), B(\tau), C(\tau)$.

The following result shows that after the change of variables $\tilde{\tau} = 3\tau$, the ring of quasimodular forms for $SL(2, \mathbb{Z})$ embeds in the ring of quasimodular form for $\Gamma_1(3)$.

Proposition 4.8. *We have the embedding of graded rings*

$$\mathbb{Q}[E_2(3\tau), E_4(3\tau), E_6(3\tau)] \subset \mathbb{Q}[A(\tau), B(\tau), C(\tau)].$$

Explicitly, we have the identities

$$\begin{aligned} 3 E_2(3\tau) &= 2B(\tau) + A(\tau)^2 = \frac{3 I_{11}^2}{X}(X + 4S), \\ 9 E_4(3\tau) &= A(\tau)^4 + 8 A(\tau)C(\tau) = \frac{I_{11}^4}{X}(X + 8), \\ 27 E_6(3\tau) &= -A(\tau)^6 + 20 A(\tau)^3 C(\tau) + 8 C(\tau)^2 = \frac{-I_{11}^6}{X^2}(X^2 - 20X - 8). \end{aligned} \tag{63}$$

Proof. For every $n \geq 1$, if $f(\tau)$ is a modular (respectively, quasimodular) form for $SL(2, \mathbb{Z})$ of weight k , then $f(n\tau)$ is a modular (respectively, quasimodular) form for $\Gamma_0(n)$ of weight k .

Once we know the modularity properties of each side, the identities (63) are easy to prove: it is enough to match finitely many terms of the \mathbb{Q} -expansions. Expressions in terms of X, I_{11} and S follow from (57). □

4.4. Genus 0 invariants of (\mathbb{P}^2, E)

Applying Theorem 1.2 for $g = 0$ (which reduces in this case to the genus 0 local-relative correspondence of [29]), we get

$$F_0^{\mathbb{P}^2/E} = F_0^{K_{\mathbb{P}^2}}.$$

Lemma 4.9.

$$DF_0^{K_{\mathbb{P}^2}} = -I_2 \tag{64}$$

$$D^2 F_0^{K_{\mathbb{P}^2}} = -3 \frac{I_{12}}{I_{11}} \tag{65}$$

$$D^3 F_0^{K_{\mathbb{P}^2}} = -9C^{-1} = -\frac{9X}{I_{11}^3}. \tag{66}$$

Proof. The formula (64) is the genus 0 mirror theorem for $K_{\mathbb{P}^2}$ [30, 57, 14]. Formula (65) follows directly from (64) and (60).

Taking the derivative of (65), we obtain $D^3 F_0^{K_{\mathbb{P}^2}} = -\frac{9I_{22}}{I_{11}}$ where

$$I_{22} = q \frac{d}{dq} \left(\frac{I_{12}(q)}{I_{11}(q)} \right).$$

Theorem 2 in [75] shows that $I_{22} = \frac{X}{I_{11}^2}$. We end the proof of (66) using that $C = \frac{I_{11}^3}{X}$ according to (57). □

Proposition 4.10. For every $n \geq 1$, we have

$$D^{n+2}F_0^{K_{\mathbb{P}^2}} \in C^{-n}\mathbb{Q}[A, B, C]_{2n-2}.$$

Proof. The case $n = 1$ is clear by (66). The general case follows by induction on n from Lemma 4.6. \square

For example, using (60)–(62) or (61)–(51), we get

$$D^4F_0^{K_{\mathbb{P}^2}} = \frac{81SX}{I_{11}^4} = \frac{27}{2}C^{-2}(B - A^2). \tag{67}$$

4.5. Genus 1 invariants of (\mathbb{P}^2, E)

Theorem 4.11 (=Theorem 1.4). We have

$$\begin{aligned} F_1^{\mathbb{P}^2/E} &= -\frac{1}{24} \log(-\mathcal{Q}) + \frac{1}{2} \sum_{n \geq 1} \log(1 - \mathcal{Q}^n) - \frac{1}{2} \sum_{n \geq 1} \log(1 - \mathcal{Q}^{3n}) \\ &= -\frac{1}{24} \log q + \frac{1}{24} \log(1 + 27q). \end{aligned}$$

Proof. According to Theorem 1.2, we have

$$F_1^{K_{\mathbb{P}^2}} = -F_{1,\emptyset}^{\mathbb{P}^2/E} + F_{1,\emptyset}^E.$$

By formulae (A.3) and (A.15) of [37], we have

$$\begin{aligned} F_1^{K_{\mathbb{P}^2}} &= -\frac{1}{12} \log q - \frac{1}{2} \log I_{11} - \frac{1}{12} \log(1 + 27q) \\ &= -\frac{1}{12} \log(-\mathcal{Q}) - \frac{1}{2} \sum_{n \geq 1} \log(1 - \mathcal{Q}^{3n}) - \frac{1}{2} \sum_{n \geq 1} \log(1 - \mathcal{Q}^n). \end{aligned}$$

On the other hand, by (4) we have

$$F_{1,\emptyset}^E = -\frac{1}{24}(-\tilde{\mathcal{Q}}) + \bar{F}_{1,\emptyset}^E$$

and it is well-known [19] that

$$\bar{F}_{1,\emptyset}^E = -\sum_{n \geq 1} \log(1 - \tilde{\mathcal{Q}}^n).$$

As $\tilde{\mathcal{Q}} = \mathcal{Q}^3$, we get

$$F_{1,\emptyset}^E = -\frac{1}{8} \log(-\mathcal{Q}) - \sum_{n \geq 1} \log(1 - \mathcal{Q}^{3n}),$$

so

$$\begin{aligned} F_1^{\mathbb{P}^2/E} &= F_{1,\emptyset}^E - F_1^{K_{\mathbb{P}^2}} \\ &= -\frac{1}{8} \log(-\mathcal{Q}) - \sum_{n \geq 1} \log(1 - \mathcal{Q}^{3n}) \\ &\quad + \frac{1}{12} \log(-\mathcal{Q}) + \frac{1}{2} \sum_{n \geq 1} \log(1 - \mathcal{Q}^{3n}) + \frac{1}{2} \sum_{n \geq 1} \log(1 - \mathcal{Q}^n) \\ &= -\frac{1}{24} \log(-\mathcal{Q}) + \frac{1}{2} \sum_{n \geq 1} \log(1 - \mathcal{Q}^n) - \frac{1}{2} \sum_{n \geq 1} \log(1 - \mathcal{Q}^{3n}), \end{aligned}$$

which equals

$$-\frac{1}{24} \log q + \frac{1}{24} \log(1 + 27q)$$

by formula (A.14) of [37]. □

Remark 4.12. Expanding the right-hand side of Theorem 4.11, we get

$$F_1^{\mathbb{P}^2/E} = -\frac{1}{24} \log Q + \frac{7Q}{8} - \frac{129Q^2}{16} + \frac{589Q^3}{6} - \frac{43009Q^4}{32} + \frac{392691Q^5}{20} + \dots$$

Lemma 4.13.

$$DF_1^{\mathbb{P}^2/E} = -\frac{1}{8} \frac{X}{I_{11}} = -\frac{1}{8} \frac{A^2}{C}. \tag{68}$$

Proof. Using Theorem 4.11, we obtain

$$\begin{aligned} Q \frac{dF_1^{\mathbb{P}^2/E}}{dQ} &= -\frac{1}{24} \frac{1}{I_{11}} q \frac{d}{dq} (\log q - \log(1 + 27q)) = -\frac{1}{24} \frac{1}{I_{11}} \left(1 - \frac{27q}{1 + 27q}\right) \\ &= -\frac{1}{24} \frac{(1 + 27q)^{-1}}{I_{11}} = -\frac{1}{24} \frac{X}{I_{11}}. \end{aligned}$$

We get the expression in terms of quasimodular forms using (58). □

Proposition 4.14. For every $n \geq 1$, we have

$$D^n F_1^{\mathbb{P}^2/E} \in C^{-n} \mathbb{Q}[A, B, C]_{2n}.$$

Proof. The case $n = 1$ is clear by (68). The general case follows by induction on n from Lemma 4.6. □

For example, using (60)–(62) or (51)–(61), we get

$$D^2 F_1^{\mathbb{P}^2/E} = \frac{3X}{8I_{11}^2} \left(S - \frac{2}{3}(X - 1) \right) = \frac{A}{16C^2} (-5A^3 + AB + 4C). \tag{69}$$

Remark 4.15. We have $\frac{\partial}{\partial S}(DF_1^{\mathbb{P}^2/E}) = 0$, as predicted by the holomorphic anomaly equation (22).

4.6. Proof of finite generation and quasimodularity

We prove the finite generation statements of Theorems 1.3 and 1.6. By Proposition 4.3, it is enough to prove the finite generation part of Theorem 1.3.

The finite generation property for local \mathbb{P}^2 is known by [51, 17]: we have

$$F_g^{K_{\mathbb{P}^2}} \in C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6} \tag{70}$$

for every $g \geq 2$. More precisely, the result proved in [51] is slightly weaker, using the generator $L = X^{1/3}$ instead of X , not getting the optimal degree bound on X , and not mentioning the ‘orbifold regularity’. However, using the R -matrix techniques used in [52] and its appendix, it is possible to prove that $F_g^{K_{\mathbb{P}^2}} \in [X^{-(g-1)} \cdot \mathbf{R}_{\leq 3g-3}]^{\text{reg}}$ for every $g \geq 2$. Such refinement of the R -matrix is described in [35] for the proof of a ‘graded finite generation’ for the quintic 3-fold. Once we know that $F_g^{K_{\mathbb{P}^2}} \in [X^{-(g-1)} \cdot \mathbf{R}_{\leq 3g-3}]^{\text{reg}}$, we get that $F_g^{K_{\mathbb{P}^2}} \in C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}$ by Proposition 4.3. Alternatively, one can use [17], which proves directly the result in terms of quasimodular forms.

We show by induction on g that, for every $g \geq 2$, we have

$$F_g^{\mathbb{P}^2/E} \in C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}.$$

Let $g \geq 2$. According to Theorem 1.2, we have

$$(-1)^{g-1} F_g^{\mathbb{P}^2/E} = -F_g^{K_{\mathbb{P}^2}} + \sum_{n \geq 0} \sum_{\substack{g=h+g_1+\dots+g_n, \\ \mathbf{a}=(a_1, \dots, a_n) \in \mathbb{Z}_{\geq 0}^n \\ (a_j, g_j) \neq (0,0), \sum_{j=1}^n a_j = 2h-2}} \frac{(-1)^{h-1} F_{h,\mathbf{a}}^E}{|\text{Aut}(\mathbf{a}, \mathbf{g})|} \prod_{j=1}^n (-1)^{g_j-1} D^{a_j+2} F_{g_j}^{\mathbb{P}^2/E},$$

where the variable \tilde{Q} in the definition (4) of $F_{h,\mathbf{a}}^E$ is expressed in terms of the variable Q in the definitions (8) and (9) of $F_g^{K_{\mathbb{P}^2}}$ and $F_g^{\mathbb{P}^2/E}$ by (7):

$$\tilde{Q} = \exp\left(-D^2 F_0^{K_{\mathbb{P}^2}}\right).$$

By (70), we know that $F_g^{K_{\mathbb{P}^2}} \in C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}$. Therefore, it remains to show that each summand

$$\frac{(-1)^{h-1} F_{h,\mathbf{a}}^E}{|\text{Aut}(\mathbf{a}, \mathbf{g})|} \prod_{j=1}^n (-1)^{g_j-1} D^{a_j+2} F_{g_j}^{\mathbb{P}^2/E} \tag{71}$$

belongs to $C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}$.

Terms with $n = 0$ only arise for $h = 1$, so $g = 1$. Thus, for $g \geq 2$, only the terms with $n \geq 1$ contribute, and by Theorem 3.5, the series $F_{h,\mathbf{a}}^E$ are quasimodular as functions of $\tilde{\tau}$, where $\tilde{Q} = e^{2i\pi\tilde{\tau}}$. More precisely, we have

$$F_{h,\mathbf{a}}^E \in \mathbb{Q}[E_2(\tilde{\tau}), E_4(\tilde{\tau}), E_6(\tilde{\tau})]_{\sum_{j=1}^n (a_j+2)}.$$

Our aim is to show quasimodularity as functions of τ , which is given by (56) as

$$Q = e^{2i\pi\tau} = -\exp\left(\frac{I_{12}}{I_{11}}\right).$$

By (65), we have $D^2 F_0^{K_{\mathbb{P}^2}} = -3 \frac{I_{12}}{I_{11}}$, so $\tilde{\tau} = 3\tau$ and $\tilde{Q} = Q^3$. Using Proposition 4.8, we deduce that

$$F_{h,\mathbf{a}}^E \in \mathbb{Q}[A, B, C]_{\sum_{j=1}^n (a_j+2)}.$$

By induction on the genus, we know that for every $g_j \geq 2$

$$F_{g_j}^{\mathbb{P}^2/E} \in C^{-(2g_j-2)} \cdot \mathbb{Q}[A, B, C]_{6g_j-6},$$

so

$$D^{a_j+2} F_{g_j}^{\mathbb{P}^2/E} \in C^{-(2g_j-2+a_j+2)} \cdot \mathbb{Q}[A, B, C]_{6g_j-6+2a_j+4}$$

using Lemma 4.6. By Propositions 4.10 and 4.14, this result also holds for $g_j = 0$ (in this case, $a_j \geq 1$ so $a_j + 2 \geq 3$) and $g_j = 1$.

Therefore,

$$\frac{(-1)^{h-1} F_{h,\mathbf{a}}^E}{|\text{Aut}(\mathbf{a}, \mathbf{g})|} \prod_{j=1}^n (-1)^{g_j-1} D^{a_j+2} F_{g_j}^{\mathbb{P}^2/E} \in C^{-(2g-2)} \cdot \mathbb{Q}[A, B, C]_{6g-6}$$

follows from

$$h + \sum_{j=1}^n g_j = g, \quad \sum_{j=1}^n a_j = 2h - 2.$$

4.7. Genus 2 invariants of (\mathbb{P}^2, E)

We prove Theorem 1.5.

By Theorem 1.2, we have

$$F_2^{K_{\mathbb{P}^2}} = F_2^{\mathbb{P}^2/E} + D^2 F_1^{\mathbb{P}^2/E} \cdot F_{1,(0)}^E - \frac{1}{2} (D^3 F_0^{\mathbb{P}^2/E})^2 \cdot F_{2,(1,1)}^E + D^4 F_0^{\mathbb{P}^2/E} \cdot F_{2,(2)}^E.$$

By [51], we have

$$F_2^{K_{\mathbb{P}^2}} = \frac{5S^3}{8X} + \frac{1}{8} S^2 + \frac{1}{96} SX + \frac{X^2}{4320} + \frac{X}{4320} - \frac{1}{2160}.$$

Using (69)-(47)-(63), we get

$$D^2 F_1^{\mathbb{P}^2/E} \cdot F_{1,(0)}^E = -\frac{S^2}{16} + \frac{5SX}{192} - \frac{S}{24} + \frac{X^2}{96} - \frac{X}{96}.$$

Using (66)-(49)-(63), we obtain

$$-\frac{1}{2} (D^3 F_0^{\mathbb{P}^2/E})^2 \cdot F_{2,(1,1)}^E = -\frac{S^3}{2X} - \frac{3S^2}{8} - \frac{11SX}{120} + \frac{S}{60} - \frac{X^2}{135} + \frac{7X}{1080} + \frac{1}{1080}.$$

Using (67)-(50)-(63), we have

$$D^4 F_0^{\mathbb{P}^2/E} \cdot F_{2,(2)}^E = \frac{9S^3}{8X} + \frac{9S^2}{16} + \frac{47SX}{640} + \frac{S}{40}.$$

Therefore, we find that in $F_2^{\mathbb{P}^2/E}$ the coefficient of $\frac{S^3}{X}$ is

$$\frac{5}{8} - \left(-\frac{1}{2} + \frac{9}{8}\right) = 0,$$

the coefficient of S^2 is

$$\frac{1}{8} - \left(-\frac{1}{16} - \frac{3}{8} + \frac{9}{16}\right) = 0,$$

the coefficient of SX is

$$\frac{1}{96} - \left(\frac{5}{192} - \frac{11}{120} + \frac{47}{640}\right) = \frac{1}{384},$$

the coefficient of X^2 is

$$\frac{1}{4320} - \left(\frac{1}{96} - \frac{1}{135}\right) = -\frac{1}{360},$$

the coefficient of X is

$$\frac{1}{4320} - \left(-\frac{1}{96} + \frac{7}{1080} \right) = \frac{1}{240},$$

and the constant term is

$$-\frac{1}{2160} - \frac{1}{1080} = -\frac{1}{720}.$$

This concludes the proof of Theorem 1.5.

Remark 4.16. Expanding the right-hand side of Theorem 1.5, we get

$$F_2^{\mathbb{P}^2/E} = \frac{29Q}{640} - \frac{207Q^2}{64} + \frac{18447Q^3}{160} - \frac{526859Q^4}{160} + \frac{5385429Q^5}{64} + \dots$$

Using (58), we can rewrite Theorem 1.5 as

$$F_2^{\mathbb{P}^2/E} = \frac{1}{11520C^2}(-37A^6 + 5A^4B + 48A^3C - 16C^2). \tag{72}$$

Taking the S -derivative of Theorem 1.5, we obtain

$$\frac{\partial}{\partial S} F_2^{\mathbb{P}^2/E} = \frac{X}{384},$$

and so, using (68),

$$\frac{3X}{I_{11}^2} \frac{\partial}{\partial S} F_2^{\mathbb{P}^2/E} = \frac{1}{2} (DF_1^{\mathbb{P}^2/E})^2,$$

as predicted by the holomorphic anomaly equation (22).

5. Holomorphic anomaly equation for (\mathbb{P}^2, E)

In this section, we prove Theorem 1.8: that is, the holomorphic anomaly equation for the series $F_{g,n}^{\mathbb{P}^2/E}$. We will use the following definitions.

Definition 5.1. A partition \mathbf{a} of length n is an ordered set (a_1, \dots, a_n) such that

$$a_1 \geq a_2 \geq \dots \geq a_n \geq 0.$$

Note that we allow the entries a_i to be zero, which is different from the ordinary definition of a partition.

Definition 5.2. For any two partitions \mathbf{a} and \mathbf{b} of length n_1 and n_2 , respectively, we define

$$\mathbf{a} \cup \mathbf{b}$$

to be the partition of length $n_1 + n_2$ with entries that are exactly the entries of \mathbf{a} and \mathbf{b} with decreasing ordering.

5.1. Holomorphic anomaly equation for the elliptic curve

We first review some known results for the elliptic curve. Recall that we denote

$$\left\langle \tau_1 \psi_1^{k_1}, \dots, \tau_n \psi_n^{k_n} \right\rangle^E := \sum_{\tilde{Q}^d} \int_{\tilde{Q}^d} \prod_{i=1}^n \psi_i^{k_i} \text{ev}_i^* \tau_i$$

the generating series of Gromov-Witten invariants of the elliptic curve E , with $\tau_i \in H^*(E)$. Recall also that we denote by $\omega \in H^2(E)$ the (Poincaré dual) class of a point and

$$F_{g,\mathbf{a}}^E := -\frac{\delta_{g,1}\delta_{n,0}}{24} \log \left((-1)^{E \cdot E} \tilde{Q} \right) + \sum_{d \geq 0} \tilde{Q}^d \langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n,d}^E,$$

where the stationary invariants $\langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n,d}^E$ have been introduced in (1).

Using the polynomiality of the double ramification cycle in the parts of the ramification profiles, Oberdieck and Pixton [65] proved the following holomorphic anomaly equation for the Gromov-Witten theory of the elliptic curve: for $2g - 2 + n > 0$, we have

$$\begin{aligned} -24 \frac{\partial}{\partial E_2} \langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n}^E = & \sum_{\substack{g_1+g_2=g, \\ \mathbf{a}' \cup \mathbf{a}'' = \mathbf{a}}} \langle \omega \psi_1^{a'_1}, \dots, \omega \psi_s^{a'_s}, 1 \rangle_{g_1,s+1}^E \langle \omega \psi_1^{a''_1}, \dots, \omega \psi_{n-s}^{a''_{n-s}}, 1 \rangle_{g_2,n-s+1}^E \\ & + \langle \omega \psi_1^{a_1}, \dots, \omega \psi_n^{a_n}, 1, 1 \rangle_{g-1,n+2}^E - 2 \sum_{j=1}^n \langle \omega \psi_1^{a_1}, \dots, \psi_j^{a_j+1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n}^E. \end{aligned}$$

By the Virasoro constraints proved by Okounkov and Pandharipande [68], we have

$$\begin{aligned} \sum_{j=1}^n \langle \omega \psi_1^{a_1}, \dots, \psi_j^{a_j+1}, \dots, \omega \psi_n^{a_n} \rangle_{g,n}^E & = \sum_{1 \leq i \neq j \leq n} \binom{a_i + a_j + 1}{a_i} \langle \omega \psi_1^{a_1}, \dots, \widehat{\omega \psi_i^{a_i}}, \dots, \widehat{\omega \psi_j^{a_j}}, \dots, \omega \psi_n^{a_n}, \omega \psi_i^{a_i+a_j} \rangle_{g,n-1}^E, \end{aligned}$$

where as usual the caret denotes omission.

Together with the string equation, we obtain the following form of the holomorphic anomaly equation for the series $F_{g,\mathbf{a}}^E$. Let \mathbf{a} be a partition of $2h - 2$, i.e. $\sum_{i=1}^n a_i = 2h - 2$; then, for $2h - 2 + n > 0$, we have

$$\begin{aligned} -24 \frac{\partial}{\partial E_2} F_{h,\mathbf{a}}^E = & \sum_{1 \leq i, j \leq n} F_{h-1, \mathbf{a} - \vec{e}_i - \vec{e}_j}^E + \sum_{\substack{h_1+h_2=h \\ \mathbf{a}' \cup \mathbf{a}'' = \mathbf{a}}} \sum_{\substack{1 \leq i \leq l(\mathbf{a}') \\ 1 \leq j \leq l(\mathbf{a}'')}} F_{h_1, \mathbf{a}' - \vec{e}_i}^E F_{h_2, \mathbf{a}'' - \vec{e}_j}^E \\ & - 2 \sum_{1 \leq i \neq j \leq n} \binom{a_i + a_j + 1}{a_i} F_{h, \mathcal{G}_{ij}(\mathbf{a})}^E \end{aligned} \tag{1}$$

Here, for a partition $\mathbf{a} = (a_1, \dots, a_n)$, we define the gluing operation by

$$\mathcal{G}_{ij}(\mathbf{a}) = (a_1, \dots, \hat{a}_i, \dots, \hat{a}_j, \dots, a_n) \cup (a_i + a_j),$$

and for the vectors \vec{e}_i ($i = 1, \dots, n$),

$$\vec{e}_i = (0, \dots, 1, \dots, 0) \text{ with } 1 \text{ lies in the } i\text{-th component,}$$

we define (see also Definition 5.2)

$$\begin{aligned} \mathbf{a} - \vec{e}_i & := (a_1, \dots, \hat{a}_i, \dots, a_n) \cup (a_i - 1), \\ \mathbf{a} - \vec{e}_i - \vec{e}_j & := \begin{cases} (a_1, \dots, \hat{a}_i, \dots, a_n) \cup (a_i - 2), & \text{if } i = j, \\ (a_1, \dots, \hat{a}_i, \dots, \hat{a}_j, \dots, a_n) \cup (a_i - 1, a_j - 1), & \text{otherwise.} \end{cases} \end{aligned}$$

Example 5.3. Using (47)–(50), one can check directly that

$$-12 \frac{\partial}{\partial E_2} F_{2,(1^2)}^E = F_{1,(0^2)}^E + (F_{1,(0)}^E)^2 - 6F_{2,(2)}^E,$$

$$-24 \frac{\partial}{\partial E_2} F_{2,(2)}^E = F_{1,(0)}^E.$$

5.2. Holomorphic anomaly equation for local \mathbb{P}^2

We denote by $F_{g,n}^{K_{\mathbb{P}^2}}$ the generating function for the local \mathbb{P}^2 theory with n insertions of the hyperplane classes. By the divisor equation,

$$F_{g,n}^{K_{\mathbb{P}^2}} = \left(Q \frac{d}{dQ}\right)^n F_g^{K_{\mathbb{P}^2}}.$$

We have the following holomorphic anomaly equation, which was proved using various techniques in [51], [17] and [21, 22, 27, 28]:

$$\frac{X}{3I_{11}^2} \cdot \frac{\partial}{\partial S} F_{g,n}^{K_{\mathbb{P}^2}} = \frac{1}{2} \sum_{\substack{g_1+g_2=g \\ n_1+n_2=n \\ 2g_i-2+n_i \geq 0}} \binom{n}{n_1} F_{g_1,n_1+1}^{K_{\mathbb{P}^2}} \cdot F_{g_2,n_2+1}^{K_{\mathbb{P}^2}} + \frac{1}{2} F_{g-1,n+2}^{K_{\mathbb{P}^2}}. \tag{2}$$

5.3. Proof of the holomorphic anomaly equation for (\mathbb{P}^2, E)

We prove the holomorphic anomaly equation (22) for $F_{g,n}^{\mathbb{P}^2/E}$ (Theorem 1.8) by induction on the genus g .

We have $F_{0,n}^{\mathbb{P}^2/E} = F_{0,n}^{K_{\mathbb{P}^2}}$, so (22) holds for $g = 0$ by (2).

Let g and n such that $2g - 2 + n > 0$. By taking derivatives of both sides of Theorem 1.2, we obtain

$$F_{g,n}^{K_{\mathbb{P}^2}} = (-1)^g F_{g,n}^{\mathbb{P}^2/E} + \sum_{\substack{0 < h \leq g \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g,n}(h)}} (-1)^{h-1} F_{h,\mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E} \tag{3}$$

$$= (-1)^g F_{g,n}^{\mathbb{P}^2/E} + 9 F_{1,(0)}^E \cdot (-1)^{g-2} F_{g-1,n+2}^{\mathbb{P}^2/E} + \sum_{\substack{0 < h \leq g \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g,n}^+(h)}} (-1)^{h-1} F_{h,\mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}$$

where we define

$$\mathcal{P}_{g,n}(h) := \left\{ (\mathbf{g}, \mathbf{a}, \mathbf{B}) = \{(g_i, a_i, B_i)\}_{i=1}^l \mid \begin{array}{l} h + \sum_i g_i = g, \sum_i a_i = 2h - 2, \\ \sum_i |B_i| = n, 2g_i - 2 + 2a_i + 2|B_i| \geq 0, \\ g_i, a_i \in \mathbb{Z}_{\geq 0}, \prod_i B_i = \{1, 2, \dots, n\} \end{array} \right\},$$

$$\mathcal{P}_{g,n}^+(h) := \mathcal{P}_{g,n}(h) \setminus \left\{ (g-1, 0, \{1, 2, \dots, n\}) \right\},$$

$$\text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E} := \frac{1}{|\text{Aut}(\mathbf{g}, \mathbf{a}, \mathbf{B})|} \prod_i (-1)^{g_i-1} 3^{a_i+2} F_{g_i, a_i + b_i + 2}^{\mathbb{P}^2/E}.$$

Note that \mathbf{B} corresponds to the assignment of insertions, and we allow B_i to be empty in the definition of $\mathcal{P}_{g,n}(h)$. Given an element $\{(g_i, a_i, B_i)\}_{i=1}^l \in \mathcal{P}_{g,n}(h)$, we set $b_i = |B_i|$ to be the number of elements in B_i , and set $\mathbf{g} = \{g_1, \dots, g_l\}$, $\mathbf{a} = \{a_1, \dots, a_l\}$, $\mathbf{B} = \{B_1, \dots, B_l\}$. We use $l(\mathbf{g}), l(\mathbf{a}), l(\mathbf{B})$ to denote the lengths of $\mathbf{g}, \mathbf{a}, \mathbf{B}$, respectively. Here all the lengths are equal to l . Finally, $\text{Aut}(\mathbf{g}, \mathbf{a}, \mathbf{B})$ is the symmetry group consisting of permutation symmetries of $(\mathbf{g}, \mathbf{a}, \mathbf{B})$.

We denote $\partial_S := \frac{X}{3I_1^2} \cdot \frac{\partial}{\partial S}$. By (63), we have

$$\partial_S = \frac{1}{18} \cdot 24 \frac{\partial}{\partial E_2}.$$

By applying the operator ∂_S on (3) and using (47), we obtain

$$\begin{aligned} \partial_S F_{g,n}^{K_{\mathbb{P}^2}} &= (-1)^g \partial_S F_{g,n}^{\mathbb{P}^2/E} + \frac{(-1)^{g-1}}{2} F_{g-1,n+2}^{\mathbb{P}^2/E} + \sum_{\substack{0 < h \leq g, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g,n}(h)}} (-1)^{h-1} F_{h,\mathbf{a}}^E \cdot \partial_S \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E} \\ &\quad + \sum_{\substack{0 < h \leq g, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g,n}^+(\mathbf{h})}} (-1)^{h-1} \partial_S F_{h,\mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E} \tag{4} \\ &= (-1)^g \partial_S F_{g,n}^{\mathbb{P}^2/E} + \frac{1}{2} (-1)^{g-1} F_{g-1,n+2}^{\mathbb{P}^2/E} + \mathcal{C}_1 + \mathcal{C}_2 + \mathcal{C}_3 + \mathcal{C}_4 \end{aligned}$$

where \mathcal{C}_1 is the contribution of the last term in the first line, and $\mathcal{C}_2, \mathcal{C}_3, \mathcal{C}_4$ are the three contributions obtained by applying the holomorphic anomaly equation (1) to the term in the second line:

$$\begin{aligned} \mathcal{C}_1 &:= \sum_{\substack{0 < h \leq g, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g,n}(h)}} (-1)^{h-1} F_{h,\mathbf{a}}^E \cdot \partial_S \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}, \\ \mathcal{C}_2 &:= \frac{1}{18} \sum_{\substack{0 < h \leq g, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g,n}^+(\mathbf{h})}} \sum_{1 \leq i, j \leq l(\mathbf{a})} (-1)^h F_{h-1, \mathbf{a} - \vec{e}_i - \vec{e}_j}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}, \\ \mathcal{C}_3 &:= \frac{1}{18} \sum_{\substack{0 < h \leq g, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g,n}^+(\mathbf{h})}} \sum_{\substack{h_1 + h_2 = h \\ \mathbf{a}' \sqcup \mathbf{a}'' = \mathbf{a}}} \sum_{\substack{1 \leq i \leq l(\mathbf{a}') \\ 1 \leq j \leq l(\mathbf{a}'')}} (-1)^{h_1-1} F_{h_1, \mathbf{a}' - \vec{e}_i}^E (-1)^{h_2-1} F_{h_2, \mathbf{a}'' - \vec{e}_j}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}, \\ \mathcal{C}_4 &:= \frac{1}{18} \sum_{\substack{0 < h \leq g, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g,n}^+(\mathbf{h})}} \sum_{1 \leq i \neq j \leq l(\mathbf{a})} 2 \binom{a_i + a_j + 1}{a_i} (-1)^{h-1} F_{h, \mathfrak{S}_{ij}(\mathbf{a})}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}. \end{aligned}$$

On the other hand, we can first apply the holomorphic anomaly equation (2) for local \mathbb{P}^2 . We then apply equation (3) to the right-hand side of (2). We have

$$\begin{aligned} \partial_S F_{g,n}^{K_{\mathbb{P}^2}} &= \frac{1}{2} \left((-1)^{g-1} F_{g-1,n+2}^{\mathbb{P}^2/E} + \sum_{\substack{0 < h \leq g-1, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g-1,n+2}(h)}} (-1)^{h-1} F_{h,\mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E} \right) \\ &\quad + \frac{1}{2} \sum_{\substack{g_1 + g_2 = g \\ n_1 + n_2 = n \\ 2g_i - 2 + n_i \geq 0}} \binom{n}{n_1} \prod_{i=1}^2 \left((-1)^{g_i} F_{g_i, n_i+1}^{\mathbb{P}^2/E} + \sum_{\substack{0 < h \leq g_i, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g_i, n_i+1}(h)}} (-1)^{h-1} F_{h,\mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E} \right) \\ &= \frac{(-1)^{g-1}}{2} F_{g-1,n+2}^{\mathbb{P}^2/E} + \mathcal{C}'_2 + \frac{(-1)^g}{2} \sum_{\substack{g_1 + g_2 = g \\ n_1 + n_2 = n \\ 2g_i - 2 + n_i \geq 0}} \binom{n}{n_1} (-1)^{g_1-1} F_{g_1, n_1+1}^{\mathbb{P}^2/E} (-1)^{g_2-1} F_{g_2, n_2+1}^{\mathbb{P}^2/E} \\ &\quad + \mathcal{C}'_1 + \mathcal{C}'_3, \end{aligned}$$

where \mathcal{C}'_2 is the contribution of the last term in the first line, and $\mathcal{C}'_1, \mathcal{C}'_3$ are the two types of contributions of the terms in the second line:

$$\begin{aligned} \mathcal{C}'_2 &= \frac{1}{2} \sum_{\substack{0 < h \leq g-1, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g-1, n+2}(h)}} (-1)^{h-1} F_{h, \mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}, \\ \mathcal{C}'_1 &= \sum_{\substack{g_1+g_2=g \\ n_1+n_2=n \\ 2g_i-2+n_i \geq 0}} \binom{n}{n_1} (-1)^{g_1} F_{g_1, n_1+1}^{\mathbb{P}^2/E} \sum_{\substack{0 < h \leq g_2, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g_2, n_2+1}(h)}} (-1)^{h-1} F_{h, \mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}, \\ \mathcal{C}'_3 &= \frac{1}{2} \sum_{\substack{g_1+g_2=g \\ n_1+n_2=n \\ 2g_i-2+n_i \geq 0}} \binom{n}{n_1} \sum_{\substack{0 < h \leq g_1, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g_1, n_1+1}(h)}} (-1)^{h-1} F_{h, \mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E} \sum_{\substack{0 < h \leq g_2, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g_2, n_2+1}(h)}} (-1)^{h-1} F_{h, \mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}. \end{aligned}$$

Suppose that the holomorphic anomaly equation for (\mathbb{P}^2, E) (Theorem 1.8) holds for $g' < g$. Namely

$$\partial_S F_{g', n}^{\mathbb{P}^2/E} = \frac{1}{2} \sum_{\substack{g_1+g_2=g' \\ n_1+n_2=n \\ 2g_i-2+n_i \geq 0}} \binom{n}{n_1} F_{g_1, n_1+1}^{\mathbb{P}^2/E} \cdot F_{g_2, n_2+1}^{\mathbb{P}^2/E} \quad \text{for } g' < g. \tag{5}$$

We prove the holomorphic anomaly equation for genus g case by showing that

$$\mathcal{C}_1 + \mathcal{C}_4 = \mathcal{C}'_1, \quad \mathcal{C}_2 = \mathcal{C}'_2, \quad \mathcal{C}_3 = \mathcal{C}'_3.$$

These three identities follow from the following three lemmas.

Lemma 5.4. *Suppose that the holomorphic anomaly equation for (\mathbb{P}^2, E) holds for all $g' < g$. Then we have*

$$\mathcal{C}_1 + \mathcal{C}_4 = \mathcal{C}'_1.$$

Proof. By using the fact that

$$\binom{a_i + a_j + 1}{a_i} + \binom{a_i + a_j + 1}{a_j} = \binom{a_i + a_j + 2}{a_i + 1},$$

we may write \mathcal{C}_4 as

$$\frac{1}{18} \sum_{\substack{0 < h \leq g, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g, n}(h)}} \sum_{1 \leq i \neq j \leq l(\mathbf{a})} \binom{a_i + a_j + 2}{a_i + 1} (-1)^{h-1} F_{h, \mathcal{G}_{ij}(\mathbf{a})}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}. \tag{6}$$

In the summation, we can replace $\mathcal{P}_{g, n}^+(h)$ by $\mathcal{P}_{g, n}(h)$ since $l(\mathbf{a}) \geq 2$.

Let us fix a $(\mathbf{g}, \mathbf{a}, \mathbf{B}) = \{(g_i, a_i, B_i)\}_{i=1}^l \in \mathcal{P}_{g, n}(h)$ and $1 \leq s \neq t \leq l$. If we sum over those $i \neq j$ in (6) such that

$$(g_i, a_i, B_i) = (g_s, a_s, B_s), \quad (g_j, a_j, B_j) = (g_t, a_t, B_t),$$

then the total contribution can be written as

$$\begin{aligned} & (-1)^{h-1} F_{h, \mathcal{G}_{ij}(\mathbf{a})}^E \frac{1}{|\text{Aut}(P_1)|} \text{Cont}_{P_1}^{\mathbb{P}^2/E} \cdot \\ & \left(-\frac{1}{2} (-1)^{g_s+g_t-1} 3^{a_s+a_t+2} \binom{a_s+a_t+2}{a_s+1} F_{g_s, a_s+b_s+2}^{\mathbb{P}^2/E} F_{g_t, a_t+b_t+2}^{\mathbb{P}^2/E} \right) \end{aligned} \tag{7}$$

where $P_1 = \{(g_k, a_k, B_k) \in (\mathbf{g}, \mathbf{a}, \mathbf{B}) \mid k \neq s, t\}$ and

$$\text{Cont}_{P_1}^{\mathbb{P}^2/E} = \prod_{k \neq s, t} (-1)^{g_k-1} 3^{a_k+2} F_{a_k+b_k+2}^{\mathbb{P}^2/E}.$$

Now let us vary $(\mathbf{g}, \mathbf{a}, \mathbf{B}) \in P_{g,n}(h)$. We sum over those (7) with

$$(\mathbf{g}, \mathbf{a}, \mathbf{B}) = P_1 \cup \{(g_s, a'_s, B'_s)\} \cup \{(g_t, a'_t, B'_t)\}$$

such that $a'_s + b'_s = a_s + b_s$ and $a'_t + b'_t = a_t + b_t$. Since $\sum a'_i = 2h - 2$ is fixed, we also deduce that $a'_s + a'_t = a_s + a_t$ and $b'_s + b'_t = b_s + b_t$. Then we get

$$(-1)^{h-1} F_{h, \mathcal{G}_{ij}(\mathbf{a})}^E \frac{\text{Cont}_{P_1}^{\mathbb{P}^2/E}}{|\text{Aut}(P_1)|} \cdot \left(-\frac{1}{2} (-1)^{g_s+g_t-1} 3^{a_s+a_t+2} F_{g_s, a_s+b_s+2}^{\mathbb{P}^2/E} F_{g_t, a_t+b_t+2}^{\mathbb{P}^2/E} \right) \sum_{\substack{a'_s+b'_s=a_s+b_s \\ 0 \leq a'_s \leq a_s+a_t \\ 0 \leq b'_s \leq b_s+b_t}} \binom{a_s+a_t+2}{a'_s+1} \binom{b_s+b_t}{b'_s}.$$

Using the Vandermonde's identity

$$\sum_{\substack{a'_s+b'_s=a_s+b_s \\ 0 \leq a'_s \leq a_s+a_t \\ 0 \leq b'_s \leq b_s+b_t}} \binom{a_s+a_t+2}{a'_s+1} \binom{b_s+b_t}{b'_s} = \binom{a_s+b_s+a_t+b_t+2}{a_s+b_s+1} - \binom{b_s+b_t}{a_s+b_s+1} - \binom{b_s+b_t}{a_t+b_t+1},$$

the above equation can be further written as

$$T_1 + T_2$$

where

$$T_1 = (-1)^{h-1} F_{h, \mathcal{G}_{ij}(\mathbf{a})}^E \frac{\text{Cont}_{P_1}^{\mathbb{P}^2/E}}{|\text{Aut}(P_1)|} \left(-\frac{1}{2} (-1)^{g_s+g_t-1} 3^{a_s+a_t+2} \binom{a_s+b_s+a_t+b_t+2}{a_s+b_s+1} F_{g_s, a_s+b_s+2}^{\mathbb{P}^2/E} F_{g_t, a_t+b_t+2}^{\mathbb{P}^2/E} \right),$$

$$T_2 = (-1)^{h-1} F_{h, \mathcal{G}_{ij}(\mathbf{a})}^E \frac{\text{Cont}_{P_1}^{\mathbb{P}^2/E}}{|\text{Aut}(P_1)|} \cdot \left(\binom{b_s+b_t}{a_s+b_s+1} + \binom{b_s+b_t}{a_t+b_t+1} \right) \frac{1}{2} (-1)^{g_s+g_t-1} 3^{a_s+a_t+2} F_{g_s, a_s+b_s+2}^{\mathbb{P}^2/E} F_{g_t, a_t+b_t+2}^{\mathbb{P}^2/E}.$$

Using the holomorphic anomaly equation (5) for $g' < g$, it is easy to see that the total contribution of those T_1 when we vary $(\mathbf{g}, \mathbf{a}, \mathbf{B})$ and s, t is

$$-\mathcal{C}_1 + \sum_{\substack{0 < h \leq g, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g,n}(h)}} (-1)^{h-1} F_{h,\mathbf{a}}^E \frac{1}{\text{Aut}(\mathbf{g}, \mathbf{a}, \mathbf{B})} \sum_{1 \leq i \leq l(\mathbf{a})} \prod_{j \neq i} (-1)^{g_j-1} 3^{a_j+2} F_{g_j, a_j+b_j+2}^{\mathbb{P}^2/E} \left(\sum_{g'_1+g'_2=g_i} (-1)^{g'_1} F_{g'_1,1}^{\mathbb{P}^2/E} (-1)^{g'_2-1} 3^{a_2+2} F_{g'_2, a_i+b_i+3}^{\mathbb{P}^2/E} \right)$$

It is easy to check that

$$\sum_{\substack{0 < h \leq g, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g,n}(h)}} (-1)^{h-1} F_{h,\mathbf{a}}^E \frac{1}{\text{Aut}(\mathbf{g}, \mathbf{a}, \mathbf{B})} \sum_{1 \leq i \leq l(\mathbf{a})} \prod_{j \neq i} (-1)^{g_j-1} 3^{a_j+2} F_{g_j, a_j+b_j+2}^{\mathbb{P}^2/E} \left(\sum_{g'_1+g'_2=g_i} (-1)^{g'_1} F_{g'_1,1}^{\mathbb{P}^2/E} (-1)^{g'_2-1} 3^{a_2+2} F_{g'_2, a_i+b_i+3}^{\mathbb{P}^2/E} \right)$$

equals

$$\sum_{g_1+g_2=g} (-1)^{g_1} F_{g_1,1}^{\mathbb{P}^2/E} \cdot \sum_{\substack{0 < h \leq g_2, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g_2, n+1}(h)}} (-1)^{h-1} F_{h,\mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}.$$

So it remains to check the total contribution of those T_2 to \mathcal{C}_1 when we vary $(\mathbf{g}, \mathbf{a}, \mathbf{B})$ and s, t is

$$\sum_{\substack{g_1+g_2=g \\ n_1+n_2=n, n_1 > 0}} \binom{n}{n_1} (-1)^{g_1} F_{g_1, n_1+1}^{\mathbb{P}^2/E} \sum_{\substack{0 < h \leq g_2, \\ (\mathbf{g}, \mathbf{a}, \mathbf{B}) \in \mathcal{P}_{g_2, n+1}(h)}} (-1)^{h-1} F_{h,\mathbf{a}}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}$$

which is obvious. □

Lemma 5.5.

$$\mathcal{C}_2 = \mathcal{C}'_2$$

Proof. Given a term

$$\frac{1}{18} (-1)^h F_{h-1, \mathbf{a}-\vec{e}_i-\vec{e}_j}^E \cdot \text{Cont}_{(\mathbf{g}, \mathbf{a}, \mathbf{B})}^{\mathbb{P}^2/E}$$

in \mathcal{C}_2 , it can be rewritten as

$$\frac{|\text{Aut}(\mathbf{g}', \mathbf{a}', \mathbf{B}')|}{|\text{Aut}(\mathbf{g}, \mathbf{a}, \mathbf{B})|} \frac{1}{2} (-1)^{h'-1} F_{h', \mathbf{a}'}^E \cdot \text{Cont}_{(\mathbf{g}', \mathbf{a}', \mathbf{B}')}^{\mathbb{P}^2/E}$$

where $h' = h - 1$, $\mathbf{g}' = \mathbf{g}$, $\mathbf{a}' = \mathbf{a} - \vec{e}_i - \vec{e}_j$ and \mathbf{B}' is a partition of the set $\{1, 2, \dots, n + 2\}$, which can be determined from \mathbf{B} by adding $n + 1$ to the set B_i and adding $n + 2$ to the set B_j . Obviously, $(\mathbf{g}', \mathbf{a}', \mathbf{B}') \in P_{g-1, n+2}(h')$. So

$$\frac{1}{2} (-1)^{h'-1} F_{h', \mathbf{a}'}^E \cdot \text{Cont}_{(\mathbf{g}', \mathbf{a}', \mathbf{B}')}^{\mathbb{P}^2/E}$$

becomes one summand in \mathcal{C}'_2 . Now Lemma 5.5 follows from the fact that for a fixed $(\mathbf{g}', \mathbf{a}', \mathbf{B}') \in P_{g-1, n+2}(h')$, there are exactly $\frac{|\text{Aut}(\mathbf{g}, \mathbf{a}, \mathbf{B})|}{|\text{Aut}(\mathbf{g}', \mathbf{a}', \mathbf{B}')|}$ choices of $(\mathbf{g}, \mathbf{a}, \mathbf{B}), i, j$, which gives $(\mathbf{g}', \mathbf{a}', \mathbf{B}')$ via the above procedure. Actually, we see that $(\mathbf{g}, \mathbf{a}, \mathbf{B})$ can be determined from $(\mathbf{g}', \mathbf{a}', \mathbf{B}')$. The only flexibility comes from the choices of i and j . □

Lemma 5.6.

$$\mathcal{C}_3 = \mathcal{C}'_3$$

Proof. The proof is similar to the proof of Lemma 5.5. We omit the details here. □

5.4. *S*-degree bound on $F_g^{\mathbb{P}^2/E}$

We prove the *S*-degree bound of Theorem 1.3 (equivalently, the *B*-degree bound of Theorem 1.6): that is, for every $g \geq 2$,

$$\deg_S F_g^{\mathbb{P}^2/E} \leq 2g - 3. \tag{8}$$

As we only have

$$\deg_S F_g^{K_{\mathbb{P}^2}} \leq 3g - 3$$

in general, the bound (8) is not an obvious consequence of Theorem 1.2 and requires a non-trivial cancellation of higher-degree terms. For example, we have observed such cancellation in the genus 2 computation of Section 4.7 (vanishing of the terms in S^3/X and S^2). Rather than trying to prove directly this cancellation in general, we show that (8) follows from the holomorphic anomaly equation (22).

We prove (8) by induction on g . Using (60), we rewrite the holomorphic anomaly equation (22) as

$$\frac{X}{3} \frac{\partial}{\partial S} F_g^{\mathbb{P}^2/E} = \frac{1}{4} \sum_{\substack{g_1+g_2=g, \\ g_i > 0 \text{ for } i=1,2}} \left(q \frac{dF_{g_1}^{\mathbb{P}^2/E}}{dq} \right) \cdot \left(q \frac{dF_{g_2}^{\mathbb{P}^2/E}}{dq} \right). \tag{9}$$

By induction, we have for $g_j \geq 2$

$$\deg_S F_{g_j}^{\mathbb{P}^2/E} \leq 2g_j - 3,$$

so using (62),

$$\deg_S \left(q \frac{dF_{g_j}^{\mathbb{P}^2/E}}{dq} \right) \leq 2g_j - 2.$$

By (68), this bound also holds for $g_j = 1$. Therefore, the *S*-degree of the right-hand side of (9) is $\leq (2g_1 - 2) + (2g_2 - 2) = 2g - 4$, so the *S*-degree of $F_g^{\mathbb{P}^2/E}$ is $\leq 2g - 3$.

Appendix A. Product formulae for the relative theory

In this section, we mainly want to prove a product formula relating the relative theory of possibly disconnected domains and the one with connected domains (see Lemma Appendix A.4). It implies Lemma 2.13 and so completes the last step in the localization calculation of Section 2.4. Before proving Lemma Appendix A.4, we need a product formula relating the rubber theory of possibly disconnected domains and the one with connected domains.

Let D be a smooth projective variety and L a line bundle on D . We consider a rubber target over D obtained as a chain of $\mathbb{P}_D(L \oplus \mathcal{O})$. Let $\widetilde{\mathcal{M}}_{\Gamma}^{\sim}(D)$ be a moduli space of relative stable maps to the rubber target, where Γ is a possibly disconnected rubber graph (see [26, Definition 2.4]). The rubber theory is relative to the two ends of the rubber target. One of the ends of the rubber target has normal bundle L^\vee , and we denote target psi-class associated with this end by Ψ_∞ (see also [33, Section 2.5]). The target psi-class associated with the other end will be denoted by Ψ_0 .

For a vertex $v \in V(\Gamma)$, recall that we denote by γ_v the graph consisting of a single vertex v plus all the decorations on v , and by $g(v)$, $n(v)$, $b(v)$ the genus, number of markings and curve class of the

vertex v . We say that a vertex v is *unstable* if the curve class of v is pushed forward to 0 on D , v has only two relative markings and no absolute markings. We say that a vertex v is *stable* if it is not unstable. Let $V^s(\Gamma)$ be the set of stable vertices and $V^{us}(\Gamma)$ the set of unstable vertices. In the following theorem, we need to consider the stabilization map from $\overline{\mathcal{M}}_{\Gamma}^{\bullet\sim}(D)$ to the moduli of stable maps of D . Note that stabilization does not make sense on components corresponding to unstable vertices. Our convention for the stabilization map

$$\tau : \overline{\mathcal{M}}_{\Gamma}^{\bullet\sim}(D) \rightarrow \prod_{v \in V^s(\Gamma)} \overline{\mathcal{M}}_{g(v),n(v)}(D, b(v)) \times D^{|V^{us}(\Gamma)|}$$

is that we first stabilize stable components and send unstable components to the corresponding points in D . If v is an unstable vertex, the two relative markings have the same multiplicity, which we denote by d_v .

Theorem Appendix A.1. *We have the following identity.*

$$\begin{aligned} & \tau_* \left(\frac{1}{t - \Psi_{\infty}} \cap [\overline{\mathcal{M}}_{\Gamma}^{\bullet\sim}(D)]^{\text{vir}} \right) \\ &= \frac{1}{\prod_{v \in V^{us}(\Gamma)} d_v} \tau'_* \left(\prod_{v \in V^s(\Gamma)} p_v^* \left(\frac{1}{t - \Psi_{\infty}} \right) \cap \left[\prod_{v \in V^s(\Gamma)} \overline{\mathcal{M}}_{\gamma_v}^{\sim}(D) \times D^{|V^{us}(\Gamma)|} \right]^{\text{vir}} \right), \end{aligned}$$

where each Ψ_{∞} on the left-hand side and right-hand side are the target psi-classes on their corresponding moduli, p_v is the projection of the product of rubber moduli to $\overline{\mathcal{M}}_{\gamma_v}^{\sim}(D)$, and τ, τ' are the corresponding stabilization maps to $\prod_{v \in V^s(\Gamma)} \overline{\mathcal{M}}_{g(v),n(v)}(D, b(v)) \times D^{|V^{us}(\Gamma)|}$.

Proof. The theorem is an application of [25, Theorem 4.1]. Note that [25, Theorem 4.1] is stated for connected domains. But our situation requires a disconnected domain, and it is straightforward to check that the same proof works for disconnected domains. We omit the details here.

Let us recall the content of the theorem. In [25, Theorem 4.1], we consider $P_{D_0,r}$, which is the r th root stack of $P := \mathbb{P}_D(L \oplus \mathcal{O})$. After the root stack construction, there are two invariant substacks $\mathcal{D}_0, \mathcal{D}_{\infty}$ under the fiberwise \mathbb{C}^* action. \mathcal{D}_0 is the one isomorphic to the root gerbe $\sqrt[r]{D/L}$, and \mathcal{D}_{∞} is the one isomorphic to D .

The result [25, Theorem 4.1] compares rubber theory with the orbifold Gromov–Witten theory of the gerbe \mathcal{D}_0 . The topological data Γ imposes contact orders on the two ends of the rubber target. In [25], the end with normal bundle L is called the 0-side, and the other end with normal bundle L^{\vee} is called the ∞ -side. For the gerbe theory over \mathcal{D}_0 , we still use Γ to represent the topological data where the weights of roots (contact order conditions) are replaced by suitable ages. A relative marking on the 0-side of order μ corresponds to an orbifold marking of age μ/r , whereas a relative marking on the ∞ -side of order μ corresponds to an orbifold marking of age $(r - \mu)/r$. Under this convention, denote by $\overline{\mathcal{M}}_{\Gamma}^{\bullet}(\mathcal{D}_0)$ the corresponding moduli of twisted stable maps to the gerbe \mathcal{D}_0 .

For simplicity, we assume all vertices on Γ are stable. Denote the forgetful maps by the following:

$$\begin{aligned} \tau_1 : \overline{\mathcal{M}}_{\Gamma}^{\bullet}(\mathcal{D}_0) &\rightarrow \prod_{v \in V(\Gamma)} \overline{\mathcal{M}}_{g(v),n(v)}(D, b(v)), \\ \tau_2 : \overline{\mathcal{M}}_{\Gamma}^{\bullet\sim}(D) &\rightarrow \prod_{v \in V(\Gamma)} \overline{\mathcal{M}}_{g(v),n(v)}(D, b(v)). \end{aligned}$$

On the root gerbe \mathcal{D}_0 , there is a universal line bundle L_r . Let $\pi : \mathcal{C} \rightarrow \overline{\mathcal{M}}_{\Gamma}^{\bullet}(\mathcal{D}_0)$ be the universal curve and $f : \mathcal{C} \rightarrow \mathcal{D}_0$ the map to the target. Let $\mathcal{L}_r = f^*L_r$ and

$$-R^* \pi_* \mathcal{L}_r := R^1 \pi_* \mathcal{L}_r - R^0 \pi_* \mathcal{L}_r \in K^0(\overline{\mathcal{M}}_{\Gamma}^{\bullet}(\mathcal{D}_0)).$$

As mentioned above, [25, Theorem 4.1] can be generalized to moduli with a disconnected domain using the same proof. Let g be the arithmetic genus of the disconnected curve corresponding to the graph Γ . The disconnected version of the theorem implies that the following identity

$$\begin{aligned}
 & (\tau_2)_* \left(\frac{1}{t - \Psi_\infty} \cap [\overline{\mathcal{M}}_\Gamma^\bullet(D)]^{\text{vir}} \right) \\
 &= \frac{\left[(\tau_1)_* \left(\sum_{i=0}^\infty \left(\frac{t}{r} \right)^{g-i-1} c_i(-R^* \pi_* \mathcal{L}_r) \cap [\overline{\mathcal{M}}_\Gamma^\bullet(D_0)]^{\text{vir}} \right) \right]_{r^0}}{\prod_{i=1}^{\rho_\infty} \left(1 + \frac{e v_i^* c_1(L) - v_i \psi_i}{t} \right)}
 \end{aligned}$$

holds in $A_* \left(\prod_{v \in V(\Gamma)} \overline{\mathcal{M}}_{g(v), n(v)}(D, b(v)) \right) [t, t^{-1}]$ as Laurent polynomials in the formal variable t .

The right-hand side decomposes into a product of such expressions according to the connected component of domain curves, because the disconnected moduli in orbifold theory is a product of connected moduli, and $-R^* \pi_* \mathcal{L}_r$ decomposes into a sum accordingly. Applying the original connected version of [25, Theorem 4.1] to each factor, we conclude that the right-hand side is nothing but the pushforward of a product of $\left(\frac{1}{t - \Psi_\infty} \right) \cap [\overline{\mathcal{M}}_{\gamma_v}^\bullet(D)]^{\text{vir}}$, where Ψ_∞ should be treated as the target psi-class of $\overline{\mathcal{M}}_{\gamma_v}^\bullet(D)$.

If there are unstable vertices in Γ , the unstable vertices in Γ correspond to unstable vertices in the localization of the root stack $P_{D_0, r}$. It is straightforward to add factors of D in the statement of the theorem and match with the statement that we want to prove. \square

Theorem Appendix A.1 has interesting corollaries. If we take the $1/t$ coefficient of the theorem, we find the following corollary.

Corollary Appendix A.2. *If Γ has more than two stable vertices, we have that*

$$\tau_*([\overline{\mathcal{M}}_\Gamma^\bullet(D)]^{\text{vir}}) = 0.$$

If we take the coefficient of $1/t^{|V^s(\Gamma)|}$, we obtain the following corollary.

Corollary Appendix A.3.

$$\begin{aligned}
 & \tau_*(\Psi_\infty^{(|V^s(\Gamma)|-1)} \cap [\overline{\mathcal{M}}_\Gamma^\bullet(D)]^{\text{vir}}) \\
 &= \frac{1}{\prod_{v \in V^{\text{us}}(\Gamma)} d_v} \tau'_* \left(\left[\prod_{v \in V^s(\Gamma)} \overline{\mathcal{M}}_{\gamma_v}^\bullet(D) \times D^{|V^{\text{us}}(\Gamma)|} \right]^{\text{vir}} \right).
 \end{aligned}$$

More importantly, we can use Theorem Appendix A.1 to deduce Lemma 2.13. Using the notations of Section 2, we repeat below the statement of Lemma 2.13 for convenience.

Lemma Appendix A.4. *We have the following identity:*

$$\begin{aligned}
 & \tau_* \left(\frac{t}{t + \Psi} \cap [\overline{\mathcal{M}}_{\Gamma_2}^\bullet(X, D)]^{\text{vir}} \right) \\
 &= \tau'_* \left(\prod_{v \in V(\Gamma_2)} p_v^* \left(\frac{t}{t + \Psi} \right) \cap \left[\prod_{v \in V(\Gamma_2)} \overline{\mathcal{M}}_{\gamma_v}^\bullet(X, D) \right]^{\text{vir}} \right),
 \end{aligned}$$

where p_v is the projection to the factor corresponding to v , and τ, τ' are the corresponding stabilization maps to

$$\prod_{v \in V(\Gamma'_2)} \overline{\mathcal{M}}_{g(v), n(v)}(X, b(v)) \times_{X^{r(v)}} D^{r(v)}$$

with $n(v)$ the number of half-edges on v and $r(v)$ the number of roots on v .

Lemma 2.13 is a special case of Lemma Appendix A.4 with $n(v) = r(v) = 1$.

Proof. First, we have a weaker product rule without involving psi-classes:

$$\tau_*[\overline{\mathcal{M}}_{\Gamma'_2}^\bullet(X, D)]^{\text{vir}} = \tau'_* \left(\left[\prod_{v \in V(\Gamma'_2)} \overline{\mathcal{M}}_{\gamma_v}(X, D) \right]^{\text{vir}} \right). \tag{A.1}$$

A proof can be formulated by passing to moduli of stable log maps using [1] and working out the perfect obstruction theory and virtual cycles (for example, see [46, Sections 9.2, 9.3]).

Next, observe that $\frac{\Psi}{-t - \Psi}$ is in fact $\frac{\delta}{-t - \Psi}$, where δ is the divisor corresponding to the locus where the target degenerates. Expanding δ , we have

$$\begin{aligned} & \left(\frac{\delta}{-t - \Psi} \right) \cap [\overline{\mathcal{M}}_{\Gamma'_2}^\bullet(X, D)]^{\text{vir}} \\ &= \sum_{i=(\Gamma'_2)_1, (\Gamma'_2)_2} \frac{\prod_{i=1}^{m(i)} d_i}{\text{Aut}(i)} (\tau_i)_* \left(\left(\frac{1}{-t - \Psi_0} \right) \cap [\overline{\mathcal{M}}_{(\Gamma'_2)_1}^\sim(D) \times_{D^{k_i}} \overline{\mathcal{M}}_{(\Gamma'_2)_2}^\bullet(X, D)]^{\text{vir}} \right), \end{aligned} \tag{A.2}$$

where it is easy to see that the restriction of Ψ to $\overline{\mathcal{M}}_{(\Gamma'_2)_1}^\sim(D)$ becomes Ψ_0 . The splitting of Γ'_2 into $(\Gamma'_2)_1, (\Gamma'_2)_2$ is determined by the splitting of each component of Γ'_2 . Thus, the next step is to split the virtual classes according to the components of $(\Gamma'_2)_1$ and $(\Gamma'_2)_2$. First, Equation (A.1) (simply replace Γ'_2 by $(\Gamma'_2)_2$) tells us that the pushforward of $[\overline{\mathcal{M}}_{(\Gamma'_2)_2}^\bullet(X, D)]^{\text{vir}}$ splits into a product of cycles according to each component (product rule). As to the rubber moduli, Theorem Appendix A.1 implies that $\frac{1}{-t - \Psi_0} \cap [\overline{\mathcal{M}}_{(\Gamma'_2)_1}^\sim(D)]^{\text{vir}}$ also satisfies the product rule. Note that the statement of the theorem uses Ψ_∞ in order to match [25, Theorem 4.1]. But turning the rubber target upside down turns Ψ_∞ into Ψ_0 with the rest unchanged. To fit the exact statement of the theorem, we also need to change t into $-t$.

Now we can apply (A.2) to the left-hand side of Lemma Appendix A.4. Note that a vertex in $(\Gamma'_2)_1$ can be either stable or unstable. On the right-hand side, we expand the product $\prod_{v \in V(\Gamma'_2)} p_v^* \left(1 + \frac{\Psi}{-t - \Psi} \right)$ and apply (A.2) again to each $\frac{\Psi}{-t - \Psi} \cap [\overline{\mathcal{M}}_{\gamma_v}(X, D)]^{\text{vir}}$. The summand 1 in $p_v^* \left(1 + \frac{\Psi}{-t - \Psi} \right)$ covers the case when γ_v splits into $((\gamma_v)_1, (\gamma_v)_2)$, where all the vertices in $(\gamma_v)_1$ (corresponding to rubber moduli) are unstable. Such cases might appear on the left-hand side of Lemma Appendix A.4 but are missing when applying (A.2) to each $\frac{\Psi}{-t - \Psi} \cap [\overline{\mathcal{M}}_{\gamma_v}(X, D)]^{\text{vir}}$ on the right-hand side. It is straightforward to check that both sides match. □

Acknowledgments. We thank Michel van Garrel, Rahul Pandharipande, Hyenho Lho, Younghwan Bae, and Jie Zhou for helpful discussions. We also thank the anonymous referees for valuable suggestions. P.B. was supported by Dr. Max Rössler, the Walter Haefner Foundation and the ETH Zurich Foundation. H.F. was supported by ERC-2012-AdG-320368-MCSK and SwissMAP. S.G. is supported by NSFC 12061131014, 11890660, 11890661 and the 2019 National Youth Talent Support Program of China. L.W. is supported by grant ERC-2017-AdG-786580-MACI.

This project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation program (grant agreement No. 786580).

Conflict of Interest: None.

References

- [1] D. Abramovich, S. Marcus, and J. Wise, ‘Comparison theorems for Gromov-Witten invariants of smooth pairs and of degenerations’, *Ann. Inst. Fourier (Grenoble)* **64** (2014), no. 4, 1611–1667.
- [2] M. Aganagic, V. Bouchard, and A. Klemm, ‘Topological strings and (almost) modular forms’, *Comm. Math. Phys.* **277** (2008), no. 3, 771–819.
- [3] M. Alim, J. D. Lange, and P. Mayr, ‘Global properties of topological string amplitudes and orbifold invariants’, *J. High Energy Phys.* **3** (2010), 113, 30.
- [4] M. Alim, E. Scheidegger, S.-T. Yau, and J. Zhou, ‘Special polynomial rings, quasi modular forms and duality of topological strings’, *Adv. Theor. Math. Phys.* **18** (2014), no. 2, 401–467.
- [5] M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, ‘Holomorphic anomalies in topological field theories’, *Nuclear Phys. B* **405** (1993), no. 2-3, 279–304.
- [6] M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, ‘Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes’, *Comm. Math. Phys.* **165** (1994), no. 2, 311–427.
- [7] S. Bloch and A. Okounkov, ‘The character of the infinite wedge representation’, *Adv. Math.* **149** (2000), no. 1, 1–60.
- [8] P. Bousseau, *A proof of N. Takahashi’s conjecture for (\mathbb{P}^2, E) and a refined sheaves/Gromov–Witten correspondence*, Preprint, 2019, arXiv:1909.02992.
- [9] P. Bousseau, *Scattering diagrams, stability conditions and coherent sheaves on \mathbb{P}^2* , Preprint, 2019, arXiv:1909.02985.
- [10] P. Bousseau, ‘Tropical refined curve counting from higher genera and lambda classes’, *Invent. Math.* **215** (2019), no. 1, 1–79.
- [11] P. Bousseau, ‘The quantum tropical vertex’, *Geom. Topol.* **24** (2020), no. 3, 1297–1379.
- [12] Y. Cao, D. Maulik, and Y. Toda, ‘Genus zero Gopakumar-Vafa type invariants for Calabi-Yau 4-folds’, *Adv. Math.* **338** (2018), 41–92.
- [13] H.-L. Chang, S. Guo, and J. Li, *BCOV’s Feynman rule of quintic 3-folds*, Preprint, 2018, arXiv:1810.00394.
- [14] T.-M. Chiang, A. Klemm, S.-T. Yau, and E. Zaslow, ‘Local mirror symmetry: calculations and interpretations’, *Adv. Theor. Math. Phys.* **3** (1999), no. 3, 495–565.
- [15] J. Choi, S. Katz, and A. Klemm, ‘The refined BPS index from stable pair invariants’, *Comm. Math. Phys.* **328** (2014), no. 3, 903–954.
- [16] T. Coates and A. Givental, ‘Quantum Riemann–Roch, Lefschetz and Serre’, *Ann. of Math.* **165** (2007), no. 1, 15–53.
- [17] T. Coates and H. Iritani, *Gromov-Witten Invariants of Local \mathbb{P}^2 and Modular Forms*, Preprint, 2018, arXiv:1804.03292
- [18] F. Diamond and J. Shurman, *A first course in modular forms*, *Graduate Texts in Mathematics*, vol. **228**, Springer-Verlag, New York, 2005.
- [19] R. Dijkgraaf, *Mirror symmetry and elliptic curves*, in *The moduli space of curves* (Texel Island, 1994), 1995, pp. 149–163.
- [20] Y. Eliashberg, A. Givental, and H. Hofer, *Introduction to symplectic field theory*, *Geom. Funct. Anal. Special Volume, Part II* (2000), 560–673. GAFA 2000 (Tel Aviv, 1999).
- [21] B. Eynard, M. Marino, and N. Orantin, ‘Holomorphic anomaly and matrix models’, *J. High Energy Phys.* **58** (2007).
- [22] B. Eynard and N. Orantin, ‘Computation of open Gromov-Witten invariants for toric Calabi-Yau 3-folds by topological recursion, a proof of the BKMP conjecture’, *Comm. Math. Phys.* **337** (2015), no. 2, 483–567.
- [23] C. Faber and R. Pandharipande, ‘Hodge integrals and Gromov–Witten theory’, *Invent. Math.* **139** (2000), no. 1, 173–199.
- [24] C. Faber and R. Pandharipande, ‘Logarithmic series and Hodge integrals in the tautological ring’, 2000, pp. 215–252. With an appendix by Don Zagier, Dedicated to William Fulton on the occasion of his 60th birthday.
- [25] H. Fan, L. Wu, and F. You, ‘Higher genus relative Gromov–Witten theory and double ramification cycles’, *Journal of the London Mathematical Society*, available at <https://onlinelibrary.wiley.com/doi/pdf/10.1112/jlms.12417>.
- [26] H. Fan, L. Wu, and F. You, ‘Structures in genus-zero relative Gromov–Witten theory’, *J. Topol.* **13** (2020), no. 1, 269–307.
- [27] B. Fang, M.C.-C. Liu, and Z. Zong, *On the remodelling conjecture for toric Calabi-Yau 3-orbifolds*, Preprint, 2016, arXiv:1604.07123.
- [28] B. Fang, Y. Ruan, Y. Zhang, and J. Zhou, *Open Gromov-Witten theory of $K_{\mathbb{P}^2}$, $K_{\mathbb{P}^1 \times \mathbb{P}^1}$, $K_{W\mathbb{P}^1[1,2]}$, K_{F_1} and Jacobi forms*, *Comm. Math. Phys.* **369** (2019), no. 2, 675–719.
- [29] M. van Garrel, T. Graber, and H. Ruddat, ‘Local Gromov-Witten invariants are log invariants’, *Adv. Math.* **350** (2019), 860–876.
- [30] A. Givental, ‘Equivariant Gromov-Witten invariants’, *Internat. Math. Res. Notices* **13** (1996), 613–663.
- [31] A. Givental, ‘Gromov–Witten invariants and quantization of quadratic hamiltonians’, *Moscow Mathematical Journal* **1** (2001), no. 5, 551–568.
- [32] T. Graber and R. Pandharipande, ‘Localization of virtual classes’, *Invent. Math.* **135** (1999), no. 2, 487–518.

- [33] T. Graber and R. Vakil, 'Relative virtual localization and vanishing of tautological classes on moduli spaces of curves', *Duke Math. J.* **130** (2005), 1–37.
- [34] T. Graefnitz, *Tropical correspondence for smooth del Pezzo log Calabi-Yau pairs*, Preprint, 2020, arXiv:2005.14018.
- [35] S. Guo, F. Janda, and Y. Ruan, *Structure of higher genus Gromov-Witten invariants of quintic 3-folds*, Preprint, 2018, arXiv:1812.11908.
- [36] B. Haghighat, A. Klemm, and M. Rauch, 'Integrability of the holomorphic anomaly equations', *J. High Energy Phys.* **10** (2008), 097, 37.
- [37] X. Hu, 'Localized standard versus reduced formula and genus 1 local Gromov-Witten invariants', *Int. Math. Res. Not. IMRN* **20** (2015), 9921–9990.
- [38] M.-x. Huang, A.-K. Kashani-Poor, and A. Klemm, *The Ω -deformed B-model for rigid $N = 2$ theories*, *Ann. Henri Poincaré* **14** (2013), no. 2, 425–497.
- [39] M.-x. Huang and A. Klemm, 'Holomorphic anomaly in gauge theories and matrix models', *J. High Energy Phys.* **9** (2007), 054, 33.
- [40] M.-x. Huang and A. Klemm, 'Direct integration for general backgrounds', *Adv. Theor. Math. Phys.* **16** (2012), no. 3, 805–849.
- [41] M.-x. Huang, A. Klemm, and S. Quackenbush, *Topological string theory on compact Calabi-Yau: modularity and boundary conditions*, Homological mirror symmetry, 2009, pp. 45–102.
- [42] E.-N. Ionel and T.-H. Parker, 'Relative Gromov-Witten invariants', *Ann. of Math. (2)* **157** (2003), no. 1, 45–96.
- [43] D. Joyce and Y. Song, 'A theory of generalized Donaldson-Thomas invariants', *Mem. Amer. Math. Soc.* **217** (2012), no. 1020, iv+199.
- [44] E. Katz, *Line-bundles on stacks of relative maps*, Preprint, 2005, arXiv:0507322.
- [45] S. Katz, 'Genus zero Gopakumar-Vafa invariants of contractible curves', *J. Differential Geom.* **79** (2008), no. 2, 185–195.
- [46] B. Kim, H. Lho, and H. Ruddat, *The degeneration formula for stable log maps*, Preprint, 2018, arXiv:1803.04210.
- [47] A. Klemm and E. Zaslow, 'Local mirror symmetry at higher genus', in Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds (Cambridge, MA, 1999), 2001, pp. 183–207.
- [48] M. Kontsevich and Y. Soibelman, *Stability structures, motivic Donaldson-Thomas invariants and cluster transformations*, Preprint, 2008, arXiv:0811.2435.
- [49] D. Krefl and J. Walcher, 'Extended holomorphic anomaly in gauge theory', *Lett. Math. Phys.* **95** (2011), no. 1, 67–88.
- [50] H. Lho, *Gromov-Witten invariants of Calabi-Yau manifolds with two Kähler parameters*, Preprint, 2018, arXiv:1804.04399.
- [51] H. Lho and R. Pandharipande, 'Stable quotients and the holomorphic anomaly equation', *Adv. Math.* **332** (2018), 349–402.
- [52] H. Lho and R. Pandharipande, *Crepant resolution and the holomorphic anomaly equation for $[\mathbb{C}^3/\mathbb{Z}_3]$* , *Proc. Lond. Math. Soc. (3)* **119** (2019), no. 3, 781–813.
- [53] H. Lho and R. Pandharipande, 'Holomorphic anomaly equations for the formal quintic', *Peking Mathematical Journal* **2** (2019), no. 1, 1–40.
- [54] A.-M. Li and Y. Ruan, 'Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds', *Invent. Math.* **145** (2001), no. 1, 151–218.
- [55] J. Li, 'Stable morphisms to singular schemes and relative stable morphisms', *J. Differential Geom.* **57** (2001), no. 509–578.
- [56] J. Li, 'A Degeneration formula of GW-invariants', *J. Differential Geom.* **60** (2002), 199–293.
- [57] B.H. Lian, K. Liu, and S.-T. Yau, 'Mirror principle. I', *Asian J. Math.* **1** (1997), no. 4, 729–763.
- [58] R.S. Maier, 'On rationally parametrized modular equations', *J. Ramanujan Math. Soc.* **24** (2009), no. 1, 1–73.
- [59] R.S. Maier, 'Nonlinear differential equations satisfied by certain classical modular forms', *Manuscripta Math.* **134** (2011), no. 1–2, 1–42.
- [60] Maulik, Nekrasov, Okounkov, and Pandharipande, 'Gromov-Witten theory and Donaldson-Thomas theory, I', *Compos. Math.* **142** (2006), no. 5, 1263–1285.
- [61] D. Maulik and J. Shen, *Cohomological χ -independence for moduli of one-dimensional sheaves and moduli of Higgs bundles*, Preprint, 2020, arXiv:2012.06627.
- [62] T. Milanov, Y. Ruan, and Y. Shen, 'Gromov-Witten theory and cycle-valued modular forms', *J. Reine Angew. Math.* **735** (2018), 287–315.
- [63] D. Mumford, 'Towards an enumerative geometry of the moduli space of curves', *Arithmetic and geometry*, Vol. II, 1983, pp. 271–328.
- [64] N. Nekrasov and A. Okounkov, 'Membranes and sheaves', *Algebr. Geom.* **3** (2016), no. 3, 320–369.
- [65] G. Oberdieck and A. Pixton, 'Holomorphic anomaly equations and the Igusa cusp form conjecture', *Invent. Math.* **213** (2018), no. 2, 507–587.
- [66] G. Oberdieck and A. Pixton, 'Gromov-Witten theory of elliptic fibrations: Jacobi forms and holomorphic anomaly equations', *Geom. Topol.* **23** (2019), no. 3, 1415–1489.
- [67] A. Okounkov and R. Pandharipande, 'Gromov-Witten theory, Hurwitz theory, and completed cycles', *Ann. of Math.* **163** (2006), 517–560.
- [68] A. Okounkov and R. Pandharipande, 'Virasoro constraints for target curves', *Invent. Math.* **163** (2006), 47–108.
- [69] R. Pandharipande, A. Pixton, and D. Zvonkine, *Relations on $\overline{M}_{g,n}$ via 3-spin structures*, *J. Amer. Math. Soc.* **28** (2015), no. 1, 279–309.

- [70] R. Pandharipande and R.P. Thomas, 'Curve counting via stable pairs in the derived category', *Invent. Math.* **178** (2009), no. 2, 407–447.
- [71] X. Wang, *Finite generation and holomorphic anomaly equation for equivariant Gromov-Witten invariants of $K_{\mathbb{P}^1 \times \mathbb{P}^1}$* , Preprint, 2019, arXiv:1908.03691.
- [72] X. Wang, *Quasi-modularity and Holomorphic Anomaly Equation for the Twisted Gromov-Witten Theory: $\mathcal{O}(3)$ over \mathbb{P}^2* , *Acta Math. Sin. (Engl. Ser.)* **35** (2019), no. 12, 1945–1962.
- [73] L. Wu, 'A remark on Gromov-Witten invariants of quintic threefold', *Adv. Math.* **326** (2018), 241–313.
- [74] D. Zagier, *Elliptic modular forms and their applications*, in *The 1-2-3 of modular forms*, 2008, pp. 1–103.
- [75] D. Zagier and A. Zinger, *Some properties of hypergeometric series associated with mirror symmetry*, in *Modular forms and string duality*, 2008, pp. 163–177.
- [76] J. Zhou, *Arithmetic Properties of Moduli Spaces and Topological String Partition Functions of Some Calabi-Yau Threefolds* (2014). Ph.D. thesis, Harvard University, available at <https://dash.harvard.edu/handle/1/12274581>.