

---

## INSTRUCTIONS TO CONTRIBUTORS

The *Journal of Helminthology* publishes papers on all aspects of animal parasitic helminths, particularly those of medical or veterinary importance.

Manuscripts, which must be in English or French (with an English summary), should be addressed to:

The Editor, Journal of Helminthology  
London School of Hygiene and Tropical Medicine  
Keppel Street, London WC1E 7HT  
England

Two copies of a typescript, on size A4 paper with double spacing, should be submitted. Papers should be preceded by a short abstract and will normally have the following sections: brief introduction (unheaded); Materials and Methods; Results; Discussion; Acknowledgements; References. However, the form of the paper may vary, depending on its subject matter; recent past issues should be consulted for a suitable form. Illustrations should be drawn in Indian ink, preferably not more than double the final size. Care should be taken that all illustrations fit into the format of the Journal. The maximum size an illustration may be printed is 13.5 × 18 cm. Where many separate drawings are made, some indication of how they may be grouped to make a corporate plate without undue wastage of space should be indicated. Some indication of scale should normally be given on the figure. Photocopies of illustrations should also be enclosed for refereeing purposes. Lettering and numbering, which must be of a high standard, should be added by the author, with due regard for subsequent reduction.

Photographs should be glossy prints, preferably of the same size as they are to appear in the Journal (maximum size 13.5 × 18 cm). Composite prints must be mounted and can have the separate photographs abutting; they will then have a separating line inserted by the printers. All figures and letters on photographs must be inserted by the author.

Information should not be repeated in the text and in tables or figures. The legends to tables and to figures should be sufficiently detailed for the information to be understood without reference to the text.

References should be given in alphabetical order with the full title of the journal. The following are examples:

DUKE, B. O. L. (1971) The ecology of onchocerciasis in man and animals. In *Ecology and physiology of parasites* (ed A. M. Fallis) pp. 213-222. Adam Hilger Ltd.: London.

JAMES, C. and WEBBE, G. (1973) A comparison of Egyptian and East African strains of *Schistosoma haematobium*. *Journal of Helminthology*, **47**, 49-59.

50 offprints are provided free of charge; additional copies may be ordered at the proof stage.

---

# Contents

|   | Pages   |
|---|---------|
| Effect of gamma-irradiation on the survival and development of the infective larvae of the hookworm, <i>Gaigeria pachyscelis</i> . MD. Z. ANSARI and K. S. SINGH  | 283-286 |
| The use of gamma-radiation-attenuated larvae of <i>Gaigeria pachyscelis</i> Railliet and Henry, 1910, for vaccination in lambs. MD. Z. ANSARI and K. S. SINGH   | 287-290 |
| The morphology and distribution of <i>Eupolystoma</i> species (Monogenoidea) in Africa, with a description of <i>E. anterorchis</i> sp. n. from <i>Bufo pardalis</i> at the Cape. R. C. TINSLEY   | 291-302 |
| Homologous immunity in schistosomiasis mansoni using ultra-violet irradiated cercariae. Research Note. A. M. GHANDOUR and M. A. MAGID   | 303-304 |
| Efficacy of Avermectins against <i>Ancylostoma caninum</i> in dogs. Research Note. L. S. BLAIR and W. C. CAMPBELL   | 305-307 |
| Efficacy of Avermectins against <i>Dirofilaria immitis</i> in dogs. Research Note. W. C. CAMPBELL and L. S. BLAIR   | 308-310 |
| Revision of the genus <i>Cooperioides</i> Daubney, 1933 (Nematoda: Trichostrongylidae). L. M. GIBBONS   | 311-322 |
| Endoparasites of the spotted ground squirrel, <i>Spermophilus spilosoma</i> Bennett, 1833, from Colorado. R. J. BRODA and G. D. SCHMIDT   | 323-326 |
| Morphological and histochemical observations on the digestive system of <i>Cotylophoron cotylophorum</i> . V. R. PARSHAD and S. S. GURAYA   | 327-333 |
| The occurrence of <i>Oesophagostomum multifoliatum</i> Daubney and Hudson, 1932, in Nigerian cattle. J. P. FABIYI   | 335-338 |
| A study of the relationship between the energy contents of <i>Schistosoma mansoni</i> cercariae and their death during penetration of mammalian host skin. A. M. GHANDOUR and A. M. IBRAHIM   | 339-342 |
| SUPPLEMENTARY REVIEW ARTICLES   |         |
| Morphological variations within the species <i>Onchocerca armillata</i> and <i>O. gutturosa</i> in buffaloes and cattle in India, with special reference to the male tail and cuticular ornamentation. P. P. S. CHAUHAN and B. P. PANDE | 343-353 |
| Ascaridoid nematodes of amphibians and reptiles: <i>Polydelphis Travassosascaris</i> n. g. and <i>Hexametra</i> . J. F. A. SPRENT   | 355-384 |