

M. Bauer. *University Hospital Carl Gustav Carus, Dresden, Germany*

Objective: To describe the pattern of antidepressant (AD) therapy in routine care over a 6-month period and to explore associations with health-related quality of life (HRQoL).

Methods: FINDER was a 6-month prospective, observational study to investigate the HRQoL of 3,468 depressed outpatients receiving AD treatment. Type and dose of AD(s) prescribed at baseline and throughout the follow-up period was recorded and grouped into SSRIs, SNRIs, TCAs, others and combinations (ADs from >1 group). 'Switching' groups were defined when medication taken changed between period 1 (baseline-3 months) and period 2 (3-6 months). HRQoL measures included the EQ-5D Visual Analogue Scale (VAS), from 'best imaginable health' (100) to 'worst imaginable health' (0).

Results: Complete information to assess switching patterns was available for 2,672 (77%) patients. Of those, 8.0% discontinued their AD, 5.6% decreased dose, 60.5% remained on stable dose, 9.6% increased dose; 5.1% and 8.6% switched within and between AD groups, respectively. In addition, 2.7% re-started treatment or remained untreated. The mean(sd) EQ-5D VAS changes from baseline to 3-months were: 20(22), 20(22), 18(21), 17(21), 12(21), 12(24), and 13(22), respectively and from baseline to 6-months were: 24(24), 28(25), 26(24), 24(24), 16(23), 21(26) and 15(24), respectively. Those patients switching within classes and those without treatment in period 1 had worst HRQoL outcomes.

Conclusions: The majority of patients treated for depression remained on the same medication at a stable dose. HRQoL may have contributed to the decision to change AD therapy.

FINDER was supported by Eli Lilly and Company Limited & Boehringer Ingelheim GmbH

Symposium: Effective treatment in borderline personality disorder – Approaches from general and disorder-specific psychotherapies

S41.01

The currency makes all the difference! Why the therapeutic relationship should be tailored around the patients' motives

F. Caspar. *Clinical Psychology and Psychotherapy, University of Berne, Bern, Switzerland*

The therapeutic relationship is the factor which has most consistently shown an impact on psychotherapy outcome. While for other disorders, such as circumscribed phobias, the relationship seems to determine a smaller percentage of variance, for the treatment of patients with BPD it is crucial. This is accounted for by disorder specific elements also in manualized treatment (DBT) with the principle of validation. Validation can be conceptualized as criticizing behaviors while accepting motives.

In this paper it will be argued that this principle can be used even more systematically based on the concept of "Plan Analysis" by Grawe and Caspar. It will be shown how the functioning of patients can be analyzed and described in a hierarchical structure of Plans. Such a description serves as a basis for reflections about how to react

in a complementary way to problematic patient behavior while avoiding reinforcement of maladaptive behavior.

It will then be elaborated and demonstrated that to the extent to which the individual motives of a patient are met with precision, it becomes realistic to satisfy them and to reduce the motivational basis of problem behavior. In contrast, if attention is given reluctantly and reactive to patient pressure, we have to expect the well known bottomless pit.

S41.02

Effectiveness of dialectical behavioral therapy for borderline personality disorder under inpatient conditions: A controlled trial and follow-up data

M. Bohus. *Department of Psychosomatics and Psychotherapy, Central Institute of Mental Health, Mannheim, Germany*

Dialectical Behavioral Therapy (DBT) was initially developed and evaluated as an outpatient treatment program for borderline personality disorder (BPD). Within the last few years, several adaptations have been developed. This study aims to evaluate a three-month DBT inpatient treatment program. Clinical outcomes, including changes on measures of psychopathology and frequency of self-mutilating acts, were assessed for 50 female patients meeting criteria for BPD. Thirty-one patients had participated in a DBT inpatient program, and 19 patients had been placed on a waiting list and received treatment as usual in the community. Post-testing was conducted four months after the initial assessment. The DBT group improved significantly more than participants on the waiting list on seven of the nine variables analyzed, including depression, anxiety, interpersonal functioning, social adjustment, global psychopathology and self-mutilation. Analyses based on Jacobson's criteria for clinically relevant change indicated that 42% of those receiving DBT had clinically recovered on a general measure of psychopathology. The effect sizes ranged between moderate and strong (see fig. 1). The data suggest that three months of inpatient DBT treatment is significantly superior to non-specific outpatient treatment. Within a relatively short time frame, improvement was found across a broad range of psychopathological features. Nine and 21 month follow-up data suggest stability of the recovery.

S41.03

Schema-focused therapy for borderline personality disorder: Effectiveness and cost-effectiveness, evidence from a multicenter trial

A. Arntz. *Clinical Psychological Science, University of Maastricht, Maastricht, The Netherlands*

Background and Aims: Although there is general consensus that only prolonged and intensive psychotherapy can provide real recovery from Borderline Personality Disorder (BPD), almost nothing is known about the relative effectiveness of different approaches. The present study compared the (cost-)effectiveness of two psychotherapies for BPD aiming at a fundamental change: a modern psychodynamic approach (Transference-Focused Psychotherapy, TFP) and schema-focused cognitive therapy (SFT).

Methods: In a multicenter trial 86 patients were randomised to either TFP or SFT and treated for max. 3 years. In Maastricht, patients also participated in fundamental studies on emotion regulation (attentional bias, fMRI, peripheral nervous system responses).

Results: TFP had more (early) drop-outs than SFT. SFT was about twice as effective as TFP in terms of recovery from BPD. This effect could not be explained by differences in drop-outs. SFT was superior