

INSTRUCTIONS FOR CONTRIBUTORS

SUBMISSION OF MANUSCRIPTS

Manuscripts should be submitted online via our manuscript submission and tracking site, <http://www.editorialmanager.com/psm/>. Full instructions for electronic submission are available directly from this site. To facilitate rapid reviewing, communications for peer review will be electronic and authors will need to supply a current e-mail address when registering to use the system.

Papers for publication from Europe (except those on genetic topics, irrespective of country), and all papers on imaging topics, should be submitted to the UK Office.

Papers from the Americas, Asia, Africa, Australasia and the Middle East (except those dealing with imaging topics), and all papers dealing with genetic topics, irrespective of country, should be sent to US Office.

Generally papers should not have text more than 4500 words in length (excluding these sections) and should not have more than a combined total of 5 tables and/or figures. Papers shorter than these limits are encouraged. For papers of unusual importance the editors may waive these requirements. A structured abstract of no more than 250 words should be given at the beginning of the article using the headings: Background; Methods; Results; Conclusions. The name of an author to whom correspondence should be sent must be indicated and a full postal address given in the footnote. Any acknowledgements should be placed at the end of the text (before the References section).

Declaration of Interest: A statement must be provided in the acknowledgements listing all financial support received for the work and, for all authors, any financial involvement (including employment, fees, share ownership) or affiliation with any organization whose financial interests may be affected by material in the manuscript, or which might potentially bias it. This applies to all papers including editorials and letters to the editor.

Contributors should also note the following:

1. S.I. units should be used throughout in text, figures and tables.
2. Authors should spell out in full any abbreviations used in their manuscripts.
3. Foreign quotations and phrases should be followed by a translation.
4. If necessary, guidelines for statistical presentation may be found in: **Altman DG, Gore SM, Gardner MJ & Pocock SJ** (1983). Statistical guidelines for contributors to medical journals. *British Medical Journal* **286**, 1489–1493.

REFERENCES (1) The Harvard (author-date) system should be used in the text and a complete list of References cited given at the end of the article. In a text citation of a work by more than two authors cite the first author's name followed by et al. (but the names of all of the authors should be given in the References section). Where several references are cited together they should be listed in rising date order. (2) The References section should be supplied in alphabetical order (authors' names in **bold**, journal titles in full), following the text. Some examples follow:

Miller PM, Byrne M, Hodges A, Lawrie SM, Johnstone EC (2002). Childhood behaviour, psychotic symptoms and psychosis onset in young people at high risk of schizophrenia: early findings from the Edinburgh high risk study. *Psychological Medicine* **32**, 173–179.

Cleckley HJ (1941). *The Mask of Sanity*, 2nd edn. Mosby: St. Louis, MO.

Brewer WJ, Wood SJ, DeLuca C, Pantelis C (2006). Models of olfaction for exploring neurodevelopment. In *Olfaction and the Brain* (ed. W. J. Brewer, D. Castle and C. Pantelis), pp. 97–121. Cambridge University Press: Cambridge.

(3) Online citations

doi (when published online prior to printed issue)

Lauritsen MB, Pedersen CB, Mortensen CB (2004). The incidence and prevalence of pervasive developmental disorders: a Danish population-based study. *Psychological Medicine*. Published online: 21 October 2004. doi:10.1017/S0033291704002387.

URL

World Bank (2003). Quantitative techniques for health equity analysis – Technical Notes (http://siteresources.worldbank.org/INTPAH/Resources/Publications/Quantitative-Techniques/health.eq_tn07.pdf). Accessed 15 February 2006.

[Authors are requested to print-out and keep a copy of any online-only material, in case the URL changes or is no longer maintained.]

FIGURES AND TABLES Only essential figures and tables should be included. Further tables, figures, photographs and appendices, may be included with the online version on the journal website. To ensure that your figures are reproduced to the highest possible standards, Cambridge Journals recommends the following formats and resolutions for supplying electronic figures. Please ensure that your figures are saved at final publication size and are in our recommended file formats. Following these guidelines will result in high quality images being reproduced in both the print and the online versions of the journal. **Line artwork:** Format: tif or eps, Colour mode: black and white (also known as 1-bit), Resolution: 1200 dpi; **Combination artwork (line/tone):** Format: tif or eps, Colour mode: grayscale (also known as 8-bit), Resolution: 800 dpi; **Black and white halftone artwork:** Format: tif, Colour mode: grayscale (also known as 8-bit), Resolution: 300 dpi; **Colour halftone artwork:** Format: tif, Colour mode: CMYK colour, Resolution: 300 dpi. All photographs, graphs, and diagrams should be referred to as figures and should be numbered consecutively in Arabic numerals. Captions for figures should be typed double-spaced on separate sheets. **Tables** Tables should be numbered consecutively in the text in Arabic numerals and each typed on a separate sheet after the References section. Titles should be typed above the table.

PROOFS AND OFFPRINTS Page proofs will be sent to the author designated to receive correspondence. corrections other than to printer's errors may be charged to the author. The corresponding author of each paper will receive a PDF file of their article and hard copy offprints may be purchased if they are ordered on the form supplied when the proof is returned.

PSYCHOLOGICAL MEDICINE

CONTENTS

REVIEW ARTICLE

- A new look at the supposed risks of early institutional rearing**
Woodhouse S, Miah A & Rutter M

1 Koops S, Gruber O, Richter A, Krämer B, Kelly S, Donohoe G, McDonald C, Cannon DM, Corvin A, Gill M, Di Giorgio A, Bertolino A, Lawrie S, Nickson T, Whalley HC, Neilson E, Calhoun VD, Thompson PM, Turner JA & Ehrlrich S 82

ORIGINAL ARTICLES

- Cognitive profiles in childhood and adolescence differ between adult psychotic and affective symptoms: a prospective birth cohort study**

Koike S, Barnett J, Jones PB & Richards M

11 Phelps AJ, Steel Z, Metcalf O, Alkemade N, Kerr K, O'Donnell M, Nursey J, Cooper J, Howard A, Armstrong R & Forbes D 95

- The natural history of risky drinking and associated harms from adolescence to young adulthood: findings from the Australian Temperament Project**

Betts KS, Alati R, Baker P, Letcher P, Hutchinson D, Youssef G & Olsson CA

23 Schweizer S, Kievit RA, Emery T, Cam-CAN & Henson RN 104

- Transmission of alcohol use disorder across three generations: a Swedish National Study**

Kendler KS, Ohlsson H, Sundquist J & Sundquist K

33 Symptoms of depression in a large healthy population cohort are related to subjective memory complaints and memory performance in negative contexts 115

- Psychotic patients who used cannabis frequently before illness onset have higher genetic predisposition to schizophrenia than those who did not**

Aas M, Melle I, Bettella F, Djurovic S, Le Hellard S, Bjella T, Ringen PA, Lagerberg TV, Smeland OB, Agartz I, Andreassen OA & Tesli M

43 Schweizer S, Kievit RA, Emery T, Cam-CAN & Henson RN 104

- Impact of dysfunctional maternal personality traits on risk of offspring depression, anxiety and self-harm at age 18 years: a population-based longitudinal study**

Pearson RM, Campbell A, Howard LM, Bornstein MH, O'Mahen H, Mars B & Moran P

50 Symptoms of depression in a large healthy population cohort are related to subjective memory complaints and memory performance in negative contexts 115

- Cognitive reserve as a moderator of the negative association between mood and cognition: evidence from a population-representative cohort**

Opdebeeck C, Matthews FE, Wu Y-T, Woods RT, Brayne C, Clare L & On behalf of the Cognitive Function AND Ageing Study Collaboration

50 Symptoms of depression in a large healthy population cohort are related to subjective memory complaints and memory performance in negative contexts 115

- Adjunctive memantine for schizophrenia: a meta-analysis of randomized, double-blind, placebo-controlled trials**

Zheng W, Li X-H, Yang X-H, Cai D-B, Ungvari GS, Ng CH, Wang S-B, Wang Y-Y, Ning Y-P & Xiang Y-T

50 Symptoms of depression in a large healthy population cohort are related to subjective memory complaints and memory performance in negative contexts 115

- Prefrontal cortical thinning links to negative symptoms in schizophrenia via the ENIGMA consortium**

Walton E, Hibar DP, van Erp TGM, Potkin SG, Roiz-Santiañez R, Crespo-Facorro B, Suarez-Pinilla P, van Haren NEM, de Zwarte SMC, Kahn RS, Cahn W, Doan NT, Jørgensen KN, Gurholt TP, Agartz I, Andreassen OA, Westlye LT, Melle I, Berg AO, Mørch-Johnsen L, Færden A, Flyckt L, Fatouros-Bergman H, Karolinska Schizophrenia Project Consortium (KaSP), Jönsson EG, Hashimoto R, Yamamori H, Fukunaga M, Jahanshad N, De Rossi P, Piras F, Banaj N, Spalletta G, Gur RE, Gur RC, Wolf DH, Satterthwaite TD, Beard LM, Sommer IE,

50 Symptoms of depression in a large healthy population cohort are related to subjective memory complaints and memory performance in negative contexts 115

61 **Altered interoceptive activation before, during, and after aversive breathing load in women remitted from anorexia nervosa**

Berner LA, Simmons AN, Wierenga CE, Bischoff-Grethe A, Paulus MP, Bailer UF, Ely AV & Kaye WH 142

61 **Post-traumatic stress disorder associated with sexual assault among women in the WHO World Mental Health Surveys**

Scott KM, Koenen KC, King A, Petukhova MV, Alonso J, Bromet EJ, Bruffaerts R, Bunting B, de Jonge P, Haro JM, Karam EG, Lee S, Medina-Mora ME, Navarro-Mateu F, Sampson NA, Shahly V, Stein DJ, Torres Y, Zaslavsky AM & Kessler RC and on behalf of the WHO World Mental Health Survey Collaborators 142

61 **Subclinical levels of anxiety but not depression are associated with planning performance in a large population-based sample**

Unterrainer JM, Domschke K, Rahm B, Wiltink J, Schulz A, Pfeiffer N, Lackner KJ, Müntzel T, Wild PS & Beutel M 142

72 **Correspondence** 155

72 **Correspondence** 168

72 **Correspondence** 175

Cambridge Core

For further information about this journal
please go to the journal web site at:
cambridge.org/psm



MIX
Paper from
responsible sources
FSC® C007785

CAMBRIDGE
UNIVERSITY PRESS