



The complete mitochondrial genome of *Ogmocotyle ailuri*: gene content, composition and rearrangement and phylogenetic implications

Research Article

Cite this article: Gao J-F *et al* (2023). The complete mitochondrial genome of *Ogmocotyle ailuri*: gene content, composition and rearrangement and phylogenetic implications. *Parasitology* **150**, 661–671. <https://doi.org/10.1017/S0031182023000379>

Received: 7 March 2023

Revised: 2 April 2023

Accepted: 4 April 2023

First published online: 13 April 2023

Keywords:

Gene arrangement; mitochondrial genome; molecular phylogeny; *Ogmocotyle ailuri*

Corresponding author:

Chun-Ren Wang;

E-mail: chunrenwang@sohu.com

Jun-Feng Gao¹, Ai-Hui Zhang¹, Wei Wei¹, Bin Jia², Jun Zhang², Ben Li¹, Ying-Yu Chen¹, Yun-Yi Sun¹, Mei-Ru Hou¹, Xue-Wei Liu¹, Jia-Wen Wang¹, Xin-Hui Zhang¹ and Chun-Ren Wang¹ 

¹Key Laboratory of Bovine Disease Control in Northeast China, Ministry of Agriculture and Rural Affairs; Heilongjiang Provincial Key Laboratory of Prevention and Control of Bovine Diseases; College of Animal Science and Veterinary Medicine, Heilongjiang Bayi Agricultural University, Daqing 163319, China and ²Branch of Animal Husbandry and Veterinary of Heilongjiang Academy of Agricultural Sciences, Qiqihar, China

Abstract

Trematodes of the genus *Ogmocotyle* are intestinal flukes that can infect a variety of definitive hosts, resulting in significant economic losses worldwide. However, there are few studies on molecular data of these trematodes. In this study, the mitochondrial (mt) genome of *Ogmocotyle ailuri* isolated from red panda (*Ailurus fulgens*) was determined and compared with those from Pronocephalata to investigate the mt genome content, genetic distance, gene rearrangements and phylogeny. The complete mt genome of *O. ailuri* is a typical closed circular molecule of 14 642 base pairs, comprising 12 protein-coding genes (PCGs), 22 transfer RNA genes, 2 ribosomal RNA genes and 2 non-coding regions. All genes are transcribed in the same direction. In addition, 23 intergenic spacers and 2 locations with gene overlaps were determined. Sequence identities and sliding window analysis indicated that *cox1* is the most conserved gene among 12 PCGs in *O. ailuri* mt genome. The sequenced mt genomes of the 48 Plagiorchiida trematodes showed 5 types of gene arrangement based on all mt genome genes, with the gene arrangement of *O. ailuri* being type I. Phylogenetic analysis using concatenated amino acid sequences of 12 PCGs revealed that *O. ailuri* was closer to *Ogmocotyle sikae* than to *Notocotylus intestinalis*. These data enhance the *Ogmocotyle* mt genome database and provide molecular resources for further studies of Pronocephalata taxonomy, population genetics and systematics.

Introduction

The red panda (*Ailurus fulgens*) is a globally threatened species with an estimated >40% population decline over the past 50 years in the wild. It occurs in the wild in India, Nepal, Myanmar and China, where it is mainly distributed in Sichuan, Yunnan, Guizhou and Tibet, with numbers estimated to be 3000–7000 (Wei *et al.*, 1999). The International Union for Conservation of Nature upgraded the status of the species from vulnerable to endangered in 2015 (Goździewska-Harłajczuk *et al.*, 2021). The red panda was also included in a national key protected wildlife list in China by the State Forestry and Grassland Administration and the Ministry of Agriculture and Rural Affairs in 2021 (www.forestry.gov.cn/main/5461/20210205/122418860831352.html).

Disease is one of the most important factors to consider when managing endangered species (Sharma and Achhami, 2022). Parasitic diseases are particularly serious given that they can influence the reproductive rate and population abundance in captive wild animals (Albon *et al.*, 2002). *Ogmocotyle* spp. are intestinal trematodes of various mammalian species and belong to the family Notocotylidae, suborder Pronocephalata, order Plagiorchiida. *Ogmocotyle* trematodes have been reported in a variety of definitive hosts, including ruminants (Ma *et al.*, 2016), primates (Coil, 1966; Yoshimura *et al.*, 1969; Iwaki *et al.*, 2012), waterfowl (Xu *et al.*, 2021), panda (Li *et al.*, 2020) and red panda (Pice, 1954). The flukes mainly inhabit the small intestine, with large numbers of flukes resulting in ogmocotylosis, which manifests clinically as diarrhoea, mucosal haemorrhage, haemorrhagic enteritis, malnutrition and even death (Ma *et al.*, 2016). Trematodes of this genus are mainly distributed in Asian countries, including China, Korea, Japan and India (Coil, 1966; Eom *et al.*, 1984; Iwaki *et al.*, 2012; Ma *et al.*, 2016).

Morphological descriptions have had a key role in identifying and differentiating parasites, with some synonym species within the genus *Ogmocotyle* from various hosts being reported. Sey and Graber (1991) conducted a systematic morphological description of *Ogmocotyle* trematodes known at that time, and identified 6 valid species in the genus: *Ogmocotyle africanum*, *Ogmocotyle ratti*, *Ogmocotyle fujianensis*, *Ogmocotyle indica* (synonym *Ogmocotyle capricori*), *Ogmocotyle sikae* (synonym *Ogmocotyle pygargi*) and *Ogmocotyle ailuri* (synonym *Ogmocotyle macacae*) (Sey and Graber, 1991; Yang and Zhang, 2013). Two species, *O. indica* and *O. ailuri*,

© The Author(s), 2023. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

were reported to infest red pandas. The main morphological identification characteristics distinguishing the 2 species were the location of the cirrus pouch and testis shape. In *O. indica*, the cirrus pouch occurs longitudinally in one-third of the body, and the testis is either kidney shaped or oval with a smooth surface; by contrast, in *O. ailuri*, the cirrus pouch lies on the front half of the body in an arc shape, and the testis is lobular (Sey and Graber, 1991).

The emergence of molecular data has resolved many of the limitations associated with morphological approaches for comparing related species and any further in-depth analyses. For example, genus *Orientobilharzia* was originally called *Schistosoma*, and then re-named *Ornithobilharzia* based upon the presence of numerous testes. Subsequently, it transferred to the new genus *Orientobilharzia* to accommodate blood flukes whose definitive hosts were mammals (Wang *et al.*, 2009). Aldhoun and Littlewood (2012) then proposed on molecular grounds that the genus *Orientobilharzia* was in fact a junior synonym of *Schistosoma* (Aldhoun and Littlewood, 2012). In addition, controversy has surrounded whether the hookworms *Bunostomum trigonocephalum* and *Bunostomum phlebotomum* represent different species or strains, given significant similarities in morphology. However, molecular data were used to show that *B. trigonocephalum* and *B. phlebotomum* represent distinct but closely related species (Gao *et al.*, 2014). Thus, more molecular data are needed to reassess traditional morphological classifications.

As a result of maternal inheritance, an apparent lack of recombination, rapid evolutionary rates and comparatively conserved genomic structures, mitochondrial (mt) genomes are highly suitable for trematode taxonomy, population genetics and systematics studies (Ichikawa-Seki *et al.*, 2017; Le *et al.*, 2020; Valentyne *et al.*, 2020). Although 6 valid species have been recognized in the genus *Ogmocotyle*, only the *O. sikae* mt genome is currently available in GenBank. The paucity of molecular data is a limitation for population genetic and phylogenetic studies of these parasites.

Therefore, the objectives of the current study were to determine the complete *O. ailuri* mt genome, compare the mtDNA genome with those previously reported for Pronocephalata trematodes, and reconstruct the phylogenetic relationships with other trematodes based on a dataset comprising the concatenation of 12 protein-coding gene (PCG) nucleotide sequences. The results provide a molecular resource for further studies of Pronocephalata taxonomy, population genetics and systematics.

Materials and methods

Parasites and total genomic DNA isolation

An autopsy was performed to determine the cause of death in red pandas in Longsha Zoological and Botanical Garden, Heilongjiang Province, China. Trematodes were collected from the small intestine of red panda following Wildlife Protection Law protocols of the People's Republic of China (Fang *et al.*, 2022). The trematodes were identified to the species level based on previously described morphological keys (Sey and Graber, 1991) (Fig. S1). They were then fixed in 75% ethanol and stored at -20°C until use. Total genomic DNA was isolated from a single fluke using a sodium dodecyl sulphate/proteinase K treatment, followed by spin-column purification using Genomic DNA Purification System (Promega, Madison, Wisconsin, USA) according to the manufacturer's protocol.

Sequencing and mt genome assembly

Illumina paired-end shotgun libraries were prepared using the standard protocol of the Nextera™ DNA Sample Prep Kit

(Epicentre, Madison, Wisconsin, USA) and sequenced using an Illumina NovaSeq sequencing platform (Personal Biotechnology Co, Ltd, Shanghai, China) using 2×100 cycles. Raw sequence data were deposited into the Short Read Archive database (www.ncbi.nlm.nih.gov/sra/) under accession number PRJNA896356. Clean data without sequencing adapters were assembled using NOVOPlasty software (Dierckxsens *et al.*, 2017) with the parameters of the genome range (13 000–15 000) and k-mer 39. The completeness of the mt genome assembly was further verified by polymerase chain reaction and Sanger sequencing using 4 pairs of primers designed on the basis of conserved regions (Table S1; Fig. S2).

Sequence analysis and gene annotation

Mt genome sequences were used for a BLAST search of the NCBI database (<http://blast.ncbi.nlm.nih.gov/Blast>) (Altschul *et al.*, 1997). Twelve PCGs were initially identified using 'ORF Finder' through NCBI and the MITOS Web Server (Rombel *et al.*, 2002) to specify the mt genetic code of invertebrates. The MITOS Web Server was then used to calculate the potential stem-loop secondary structures within these tRNA gene sequences (Bernt *et al.*, 2013). The codon usage of the 12 PCGs was analysed using the invertebrate genetic code and the Codon Usage web server (www.bioinformatics.org/sms2/codon_usage.html). An analysis of compositional skews was conducted using equations (1) and (2) (Perna and Kocher, 1995):

$$\text{AT-skew} = (A - T)/(A + T) \quad (1)$$

$$\text{GC-skew} = (G - C)/(G + C) \quad (2)$$

The relative synonymous codon usage of the 12 PCGs was calculated using the Sequence Manipulation Suite (www.detaibio.com/sms2/codon_usage.html) (Stothard, 2000). A sliding window analysis [window length = 300 base pairs (bp), step size = 10 bp] was conducted using DnaSP v.5 (Librado and Rozas, 2009) to assess nucleotide diversity π between 12 PCGs from 14 Pronocephalata mt genomes. Nucleotide diversity was plotted against the mid-point positions of each window, and gene boundaries were identified. The differences between nucleotide and amino acid sequences were calculated using MEGA 11.0 and MegAlign (Burland, 2000; Tamura *et al.*, 2021). A gene map of the mt genome was constructed using the online mitochondrial visualization tool OrganellarGenomeDRAW (Lohse *et al.*, 2013). To determine the occurrence of *Plagiorchiida* spp. gene recombination, the gene arrangement of the *O. ailuri* mt genome was compared with those of 47 *Plagiorchiida* species currently available. The circular mt genomes were linearized at the 5' end of their *cox3* genes in the H-strand direction, as previously reported (Gao *et al.*, 2021).

Phylogenetic analysis

Phylogenetic relationships among the 48 representative members of *Plagiorchiida* available in GenBank (Table S2) were determined based on concatenated amino acid sequences of 12 PCGs, using *Postharmostomum commutatum* (NC_044643) as the outgroup. The PCG amino acid sequences were aligned using MAFFT v7.471 (Katoh and Standley, 2013), and the alignment was processed for elimination of poorly aligned sites and divergent regions using the GBlocks Server (http://molevol.cmima.csic.es/castresana/Gblocks_server.html).

Phylogenetic trees were reconstructed using Bayesian inference (BI) methods. The best-fit substitution model for phylogenetic

Table 1. Features of the mt genome of *Ogmocotyle ailuri*

Genes	Location	Length (bp)	Start codons/stop codons	Anti-codons	Intergenic spacers (bp)
<i>cox3</i>	1–645	645	ATG/TAG		7
<i>trnH</i>	653–719	67		GTG	3
<i>cytb</i>	723–1832	1110	ATG/TAA		26
<i>nad4L</i>	1859–2110	252	ATA/TAG		–40
<i>nad4</i>	2071–3357	1287	ATG-TAG		4
<i>trnQ</i>	3362–3424	63		TTG	14
<i>trnF</i>	3439–3506	68		GAA	14
<i>trnM</i>	3521–3586	66		CAT	4
<i>atp6</i>	3591–4106	516	ATG/TAG		14
<i>nad2</i>	4121–4990	870	ATG/TAG		12
<i>trnV</i>	5003–5067	65		TAC	8
<i>trnA</i>	5076–5140	65		TGC	1
<i>trnD</i>	5142–5211	70		GTC	0
<i>nad1</i>	5212–6111	900	ATG/TAA		6
<i>trnN</i>	6118–6187	70		GTT	9
<i>trnP</i>	6197–6265	69		TGG	0
<i>trnI</i>	6266–6330	65		GTA	10
<i>trnK</i>	6341–6407	67		CTT	3
<i>nad3</i>	6411–6770	360	ATG/TAG		4
<i>trnS1</i>	6777–6836	60		GCT	5
<i>trnW</i>	6842–6907	66		TCA	0
<i>cox1</i>	6908–8455	1548	ATA/TAA		9
<i>trnT</i>	8465–8533	69		TGT	0
<i>rrnL</i>	8534–9527	995			0
<i>trnC</i>	9528–9596	69		GCA	0
<i>rrnS</i>	9597–10 344	748			0
<i>cox2</i>	10 345–10 920	576	ATG/TAG		6
<i>nad6</i>	10 927–11 379	453	ATG/TAG		18
<i>trnY</i>	11 398–11 461	64		GTA	0
<i>trnL1</i>	11 462–11 529	68		TAG	–1
<i>trnS2</i>	11 529–11 591	63		TGA	37
<i>trnL2</i>	11 629–11 691	63		TAA	2
<i>trnR</i>	11 694–11 758	65		TCG	0
<i>nad5</i>	11 759–13 339	1581	GTG/TAG		4
<i>trnG</i>	13 344–13 410	67		TCC	0
SNCR	13 411–14 087	677			0
<i>trnE</i>	14 088–14 155	68		TTC	0
LNCR	14 156–14 642	487			0

The rearrangement events mainly occurred in 2 gene blocks (*trnN-trnP-trnI-trnK* and *trnG-trnE*) of the mt genome in Plagiorchiida. Compared with type I rearrangements, *trnG* and *trnE* were interchanged within *nad5* and *cox3* in type II rearrangements. Compared with type II rearrangements, *trnP* and *trnI* were interchanged with *nad1* and *nad3* in type III rearrangements. Compared with type III rearrangements, *trnE* and *trnG* were interchanged with *nad5* and *cox3* in type IV rearrangements. *trnE* between *nad5* and *cox3* in type IV rearrangements was translocated between *nad1* and *nad3* in type V rearrangements.

Additionally, *trnY*, which is between *nad6* and *nad5* in type IV rearrangements, was translocated between *nad5* and *cox3* in type V rearrangements.

Phylogenetic analysis

BI approaches were used to estimate the phylogenetic position of *O. ailuri* within 47 other Plagiorchiida trematodes based on the concatenated amino acid sequences of 12 PCGs (Fig. 5). The result showed that the Pronocephalata and Opisthorchiata




Figure 2. A + T content and nucleotide skew of genes, individual elements and the complete mt genome of 14 Pronocephalata trematodes. Lines colour and point represent a specific Pronocephalata species.

suborders form independent monophyletic groups in the phylogenetic tree. Xiphidiata trematodes have a complicated taxonomic relationship, although each family clustered together in 1 branch; however, Xiphidiata trematodes formed 4 lineages rather than a monophyletic group. *Prosthogonimus pellucidus* and *Prosthogonimus cuneatus* clustered together with *Tamerlania zarudnyi* to form a group on the outermost branch, which was next to other Plagiorchiida trematodes. Except for *T. zarudnyi* (Eucotylidae), 12 Echinostomata trematodes clustered together to form a group.

The branch of Pronocephalata includes the clades Paramphistomidae, Gastrothylacidae, Diplodiscidae and Notocotylidae. Paramphistomidae, Gastrothylacidae and Diplodiscidae clustered together to form a group, with Notocotylidae forming another group. In the Notocotylidae branch, *O. ailuri* isolated from this study was closer to *O. sikae* than to *N. intestinalis*.

Discussion

In this study, the entire mt genome of *O. ailuri* was sequenced for the first time. The mt genome of *O. ailuri* was the classical structure of trematodes, contained 12 PCGs, 22 tRNA genes, 2 rRNA genes and 2 non-coding regions. There were a 40 bp gene overlap between *nad4L* and *nad4* in *O. ailuri* mt genome, the result was consistent with most Plagiorchiida, except *Dicrocoelium dendriticum* (8 bp), *Dicrocoelium chinensis* (8 bp), *Lyperosomum longicauda* (37 bp), *Paragonimus westermani* (14 bp) and *N. intestinalis* (0 bp) (Biswal *et al.*, 2014; Liu *et al.*, 2014; Suleman *et al.*, 2020; Xu *et al.*, 2021).

A common feature of the mt genome in most metazoans is a bias towards a higher representation of A and T nucleotides, which leads to subsequent biases in the corresponding encoded amino acids (Hu *et al.*, 2020). The AT content of the entire mt genome and the 12 PCGs was 65.51 and 64.8%, respectively,

and the result was consistent with other Plagiorchiida trematodes biased towards A and T (Atopkin *et al.*, 2021; Ivashko *et al.*, 2022). The negative AT-skew indicated a higher incidence of T compared with A nucleotides, and the positive GC-skew indicated that G was more abundant than C (Le *et al.*, 2019). The AT-skew values were negative, and the GC-skew values were positive in *O. ailuri* mt genome; these trends in AT- and GC-skew were similar to those of other Pronocephalata (Yan *et al.*, 2013; Xu *et al.*, 2021).

Three types of initiation codon (ATG, ATA and GTG) and 2 types of termination codon (TAG and TAA) occurred in the *O. ailuri* mt genome; these codons were commonly used in Plagiorchiida trematodes. The result was consistent with most Plagiorchiida trematodes, but different from some Plagiorchiida trematodes, such as *Eurytrema pancreaticum*, *L. longicauda* and *Plagiorchis maculosus* that utilized incomplete stop codon T as termination codon (Chang *et al.*, 2016; Suleman *et al.*, 2019, 2020). Two NCRs were identified in the *O. ailuri* mt genome, which is consistent with other related Plagiorchiida, such as *Fiscooederius elongatus*, *G. crumenifer* and *Paragonimus ohirai* (Yang *et al.*, 2015, 2016; Le *et al.*, 2020). However, this result is inconsistent with other related Plagiorchiida, such as *Diplodiscus japonicus* and *Tracheophilus cymbius*, in which only 1 NCR has been identified in the mt genome (Li *et al.*, 2019; An *et al.*, 2022).

The gene *cox1* was considered to be a useful barcode for metazoans, and widely used for trematode studies (Mioduchowska *et al.*, 2018). The mitochondrial genes *cox1* have also been used to study the population genetic structure of trematode on a local and global scale (Djuikwo-Teukeng *et al.*, 2019; Shumenko and Tatonova, 2022; Wang *et al.*, 2022). In the present study, the sequence identities and sliding window analysis indicated *cox1* was the most conserved gene among 12 PCGs in 14 Pronocephalata species; this result was similar to those from a study of *L. longicauda*, *O. sikae*, *N. intestinalis*, *Paramphistomum*

Table 2. Codon usage analysis of 12 PCGs in the mitochondrial genome of *Ogmocotyle ailuri*

Codon	Number	/1000	Fraction	Codon	Number	/1000	Fraction
GCG(Ala)	26	7.74	0.23	CCG(Pro)	25	7.44	0.3
GCA(Ala)	21	6.25	0.18	CCA(Pro)	15	4.47	0.18
GCU(Ala)	60	17.86	0.52	CCU(Pro)	37	11.02	0.45
GCC(Ala)	8	2.38	0.07	CCC(Pro)	6	1.79	0.07
UGU(Cys)	103	30.66	0.89	CAG(Gln)	18	5.36	0.6
UGC(Cys)	13	3.87	0.11	CAA(Gln)	12	3.57	0.4
GAU(Asp)	56	16.67	0.85	CGG(Arg)	6	1.79	0.1
GAC(Asp)	10	2.98	0.15	CGA(Arg)	14	4.17	0.23
GAG(Glu)	57	16.97	0.7	CGU(Arg)	35	10.42	0.58
GAA(Glu)	24	7.14	0.3	CGC(Arg)	5	1.49	0.08
UUU(Phe)	351	104.5	0.96	AGG(Ser)	49	14.59	0.13
UUC(Phe)	14	4.17	0.04	AGA(Ser)	38	11.31	0.1
GGG(Gly)	97	28.88	0.36	AGU(Ser)	93	27.69	0.25
GGA(Gly)	34	10.12	0.13	AGC(Ser)	12	3.57	0.03
GGU(Gly)	122	36.32	0.45	UCG(Ser)	31	9.23	0.08
GGC(Gly)	17	5.06	0.06	UCA(Ser)	34	10.12	0.09
CAU(His)	47	13.99	0.85	UCU(Ser)	95	28.28	0.26
CAC(His)	8	2.38	0.15	UCC(Ser)	14	4.17	0.04
AUU(Ile)	128	38.11	0.91	ACG(Thr)	20	5.95	0.25
AUC(Ile)	12	3.57	0.09	ACA(Thr)	12	3.57	0.15
AAG(Lys)	53	15.78	0.64	ACU(Thr)	43	12.8	0.54
AAA(Lys)	30	8.93	0.36	ACC(Thr)	4	1.19	0.05
UUG(Leu)	244	72.64	0.48	GUG(Val)	122	36.32	0.3
UUA(Leu)	182	54.18	0.36	GUA(Val)	76	22.63	0.19
CUG(Leu)	14	4.17	0.03	GUU(Val)	199	59.24	0.49
CUA(Leu)	26	7.74	0.05	GUC(Val)	12	3.57	0.03
CUU(Leu)	44	13.1	0.09	UGG(Trp)	61	18.16	0.58
CUC(Leu)	2	0.6	0	UGA(Trp)	45	13.4	0.42
AUG(Met)	100	29.77	0.55	UAU(Tyr)	144	42.87	0.85
AUA(Met)	82	24.41	0.45	UAC(Tyr)	26	7.74	0.15
AAU(Asn)	52	15.48	0.88	UAG(*)	9	2.68	0.75
AAC(Asn)	7	2.08	0.12	UAA(*)	3	0.89	0.25

leydeni and *T. zarudnyi* (Ma *et al.*, 2015, 2016; Suleman *et al.*, 2020, 2021; Xu *et al.*, 2021).

Gene arrangement in mt genomes provides a source of information for phylogenetic inference (Le *et al.*, 2001). Wey-Fabrizius *et al.* (2013) reviewed how mt genome data contribute to the phylogeny debates within Platyzoan, which showed that the gene order of PCGs and rRNA genes within Platyhelminthes were highly conserved with exceptions of partial Monogenea and Seriata trematodes. For majority trematodes, the gene order is *cox3 > cyt b > nad4L > nad4 > atp6 > nad2 > nad1 > nad3 > cox1 > rrrnL > rrrnS > cox2 > nad6 > nad5* (Wey-Fabrizius *et al.*, 2013), which was well supported by the present study. Solà *et al.* (2015) reported the mt genomes and gene order of 2 free-living flatworm species (*Crenobia alpina* and *Obama* sp.), and compared with other relevant helminthes, it was found that the gene order among free-living flatworms differs considerably from the parasitic flatworms. However, the gene order of both free-living and parasitic flatworms was highly conservative with

only a few exceptions in each kind of flatworm (Solà *et al.*, 2015). The phenomenon was also found in Plagiorchiida trematodes in the present study; 48 Plagiorchiida trematodes available in GenBank shared an identical gene order based on PCGs and rRNA genes.

The gene rearrangement showed 5 types among 48 Plagiorchiida trematodes, and only occurred in transposed tRNAs. This was different from *Schistosoma* trematodes, the gene arrangement of which was radically altered by PCGs (Le *et al.*, 2001; Zhang *et al.*, 2019). In addition, the rearrangement events mainly occurred in *nad1-N-P-I-K-nad3* and *nad5-G-E-cox3* gene blocks of the mt genome in Plagiorchiida. Zhang *et al.* reported that type I (*trnG-trnE*) and type II (*trnE-trnG*) rearrangements were generated by the gene recombination of ancestral gene rearrangements in Trematoda, and hypothesized that type II evolutionarily pre-dates type I gene rearrangements (Zhang *et al.*, 2019). Thus, these gene rearrangements might be an important molecular marker of Plagiorchiida trematode evolution.

Table 3. Comparison of 12 PCGs among *Ogmocotyle ailuri* and other suborder Pronocephalata trematodes

Genes	Dn	Dm	Dj	Ni	Ogs	Oga	Pc	Cm	Ee	Ors	Gc	Fc	Fe	Pl	Nucleotides similarity (%)	Amino acid similarity (%)
	No. nucleotides/No. amino acid															
<i>cox3</i>	645/214	645/214	645/214	645/214	645/214	645/214	645/214	645/214	645/214	645/214	645/214	645/214	645/214	645/214	53.0–89.5	47.2–95.8
<i>cytb</i>	1116/371	1116/371	1113/370	1089/362	1110/369	1110/369	1113/310	1116/371	1113/370	1113/370	1113/370	1113/370	1113/370	1113/370	71.7–91.2	73.7–97.0
<i>nad4l</i>	264/87	264/87	264/87	264/87	267/88	252/83	264/87	264/87	264/87	264/87	264/87	264/87	264/87	264/87	68.9–91.3	58.6–98.9
<i>nad4</i>	1287/428	1287/428	1287/428	1287/428	1287/428	1287/428	1281/426	1281/426	1281/426	1281/426	1281/426	1281/426	1281/426	1281/426	65.3–90.0	62.2–95.1
<i>atp6</i>	516/171	516/171	516/171	516/171	516/171	516/171	516/171	516/171	516/171	516/171	516/171	516/171	516/171	516/171	67.8–92.6	60.2–98.8
<i>nad2</i>	870/289	870/289	870/289	870/289	870/289	870/289	873/290	873/290	876/291	858/285	858/285	873/290	876/291	873/290	62.9–90.0	54.0–92.4
<i>nad1</i>	903/300	903/300	903/300	891/296	891/296	900/299	897/298	897/298	897/298	897/298	903/300	897/298	897/298	897/298	70.5–91.0	68.2–97.0
<i>nad3</i>	354/117	354/117	354/117	357/118	357/118	360/119	357/118	357/118	357/118	356/118	378/125	357/118	357/118	357/118	65.2–88.5	50.4–95.8
<i>cox1</i>	1548/515	1572/523	1548/515	1548/515	1557/518	1548/515	1545/514	1542/513	1542/513	1542/513	1542/513	1542/513	1542/513	1545/514	72.4–92.5	74.3–97.3
<i>cox2</i>	585/194	585/194	585/194	579/192	579/192	576/191	579/192	585/194	582/193	582/193	582/193	582/193	582/191	582/193	67.0–94.0	63.5–97.9
<i>nad6</i>	456/151	456/151	456/151	453/150	453/150	453/150	453/150	453/150	453/150	453/150	453/150	453/150	501/166	453/150	63.4–89.3	56.7–95.4
<i>nad5</i>	1563/520	1563/520	1563/520	1581/526	1581/526	1560/519	1581/526	1581/526	1581/526	1581/526	1581/526	1578/525	1581/526	1584/527	64.5–89.0	57.3–94.9
Total nt/aa	10 107/3357	10 131/3365	10 104/3356	10 080/3348	10 113/3359	10 077/3347	10 104/3296	10 110/3358	10 110/3357	10 088/3351	10 116/3360	10 047/3355	10 155/3373	10 110/3358	68.2–90.3	51.5–95.3
Total size(bp)	14 697	14 210	14 179	14 317	14 307	14 642	14 014	14 028	13 968	13 800	14 801	14 256	14 120	14 050	64.3–89.8	

Note: Dn, *Diplodiscus nigromaculati*; Dm, *Diplodiscus mehari*; Dj, *Diplodiscus japonicus*; Ni, *Notocotylus intestinalis*; Ogs, *Ogmocotyle sikae*; Oga, *Ogmocotyle ailuri*; Pc, *Paramphistomum cervi*; Cm, *Calicophoron microbothrioides*; Ee, *Explanatum explanatum*; Ors, *Orthocoelium streptocoelium*; Gc, *Gastrothylax crumenifer*; Fc, *Fischoederius cobboldi*; Fe, *Fischoederius elongatus*; Pl, *Paramphistomum leydeni*.




Fig. 3. Sliding window analysis of the complete mt genome sequences of 14 Pronocephalata trematodes. A sliding window of 300 bp (in 10 bp overlapping steps) was used to estimate nucleotide diversity π across the alignments. Nucleotide diversity was plotted against the mid-point positions of each window. Each gene boundary is identified.

Previous research reported the phylogenetic position of *O. sikae* based on the complete mt genome (Ma et al., 2016). However, molecular data were rare at that time, and only morphological-based approaches could not fully reflect the phylogenetic relationship within Plagiorchiida trematodes. Thus, the phylogenetic relationships with other Pronocephalata trematodes previously reported were reconstructed based on the concatenation of 12 PCGs. In the phylogenetic tree, most of trematodes were consistent with the current taxonomic status of Plagiorchiida (Olson et al., 2003). Notably, except for *T. zarudnyi* (Eucotylidae), 12 Echinostomata trematodes clustered together to form a group. However, *T. zarudnyi* clustered together with Prosthonimidae flukes, a result consistent with previous

phylogenetic assessments (Guo et al., 2022). Thus, the current results showed that suborder Echinostomata is polyphyletic. Eucotylidae was considered to belong to Cyclocoeloidea (Strigeida) by Kanev et al., while Olson et al. (2003) showed that Eucotylidae should belong to the suborder Xiphidiata as a long-branched clade based on nuclear ribosomal DNA (Kanev et al., 2002; Olson et al., 2003). This hypothesis was also supported by the present study based on the complete mt genome of *O. ailuri*.

In the branch of Pronocephalata, 3 family trematodes (Paramphistomidae, Gastrothylacidae, Diplodiscidae) clustered together to form a group, and 1 family trematode (Notocotylidae) forming another group. A similar result was



Figure 4. Mt genome arrangement in *Ogmocotyle ailuri* compared with those in Plagiorchiida trematodes. The circular mt genomes were linearized at the 5' end of *cox3* gene for illustration purpose. Non-coding regions were not shown.




Figure 5. Phylogenetic relationships of *Ogmocotyle ailuri* with other 48 Plagiorchiida trematodes based on concatenated amino acid sequences of 12 protein-coding genes analysed by BI using *Postharmostomum commutatum* as the outgroup. Posterior probability values are indicated. Suborders and families are highlighted by individual colours. Circles indicate BI = 1.0, other values are given above the nodes.

also reported in previous studies using the mtDNA genome (Atopkin *et al.*, 2021; An *et al.*, 2022; Ivashko *et al.*, 2022). However, this was not consistent with those of a previous study using ITS2 for phylogenies, which demonstrated a close genetic affinity of Notocotylidae with Bucephalidae (of the order Strigeidida), rather than with Paramphistomidae and Gastrothylacidae (Xu *et al.*, 2021).

In the Notocotylidae branch, *O. ailuri* was closer to *O. sikae* than to *N. intestinalis*, which is consistent with the current taxonomic status of Notocotylidae. These data provide novel and useful genetic markers for studying systematics and population genetics, but only 3 complete mt genomes have been published for Notocotylidae species in GenBank. Therefore, more mt genomes should be sequenced to re-examine the phylogenetic relationships of Pronocephalata and other trematodes.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182023000379>.

Data availability. All data generated or used during the study appear in the submitted article.

Acknowledgements. We thank Longsha Zoological and Botanical Garden for collecting the trematode specimens, and we also thank International Science Editing (<http://www.international-scienceediting.com>) for editing this manuscript.

Author's contribution. The research was designed by J.-F. Gao and C.-R. Wang. Material preparation and data collection were performed by W. Wei, B. Jia, J. Zhang and B. Li. Y.-Y. Chen, Y.-Y. Sun and M.-R. Hou analysed the data. The first draft of the manuscript was written by A.-H. Zhang, X.-W. Liu, J.-W. Wang and X.-H. Zhang critically revised the manuscript.

Financial support. This work was supported by the National Natural Science Foundation of China (32172886), Heilongjiang Provincial Natural Science Foundation of China (LH2021C071), Heilongjiang Province Postdoctoral Science Foundation (LBH-Z19191), Heilongjiang Bayi Agricultural University support Program for San Heng San Zong

(ZRCQC202204) and Personnel Foundation of Heilongjiang Bayi Agricultural University (XYB202108).

Conflict of interest. None.

Ethical standards. The autopsy of red panda and collection of trematode specimens following the standards and procedures by the Animal Ethics Committee of Heilongjiang Bayi Agricultural University (Approval ID: WKJXY-2022020).

References

- Albon SD, Stien A, Irvine RJ, Langvatn R, Ropstad E and Halvorsen O (2002) The role of parasites in the dynamics of a reindeer population. *Proceedings of the Royal Society B* **269**, 1625–1632.
- Aldhoun JA and Littlewood DT (2012) *Orientobilharzia* Dutt & Srivastava, 1955 (Trematoda: Schistosomatidae), a junior synonym of *Schistosoma* Weinland, 1858. *Systematic Parasitology* **82**, 81–88.
- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W and Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research* **25**, 3389–3402.
- An Q, Qiu YY, Lou Y, Jiang Y, Qiu HY, Zhang ZH, Li B, Zhang AH, Wei W, Chen YY, Gao JF and Wang CR (2022) Characterization of the complete mitochondrial genomes of *Diplostiscus japonicus* and *Diplostiscus mehari* (Trematoda: Diplostiscidae): comparison with the members of the superfamily Paramphistomoidea and phylogenetic implication. *International Journal for Parasitology – Parasites and Wildlife* **19**, 9–17.
- Atopkin DM, Semenchenko AA, Solodovnik DA, Ivashko YI and Vinnikov KA (2021) First next-generation sequencing data for Haploporidae (Digenea: Haploporata): characterization of complete mitochondrial genome and ribosomal operon for *Parasaccocoeleium mugili* Zhukov, 1971. *Parasitology Research* **120**, 2037–2046.
- Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsche G, Pütz J, Middendorf M and Stadler PF (2013) MITOS: improved de novo metazoan mitochondrial genome annotation. *Molecular Phylogenetics and Evolution* **69**, 313–319.
- Biswal DK, Chatterjee A, Bhattacharya A and Tandon V (2014) The mitochondrial genome of *Paragonimus westermani* (Kerbert, 1878), the Indian

- isolate of the lung fluke representative of the family Paragonimidae (Trematoda). *PeerJ* 2, e484.
- Burland TG** (2000) DNASTAR's Lasergene sequence analysis software. *Methods in Molecular Biology* 132, 71–91.
- Chang QC, Liu GH, Gao JF, Zheng X, Zhang Y, Duan H, Yue DM, Fu X, Su X, Gao Y and Wang CR** (2016) Sequencing and characterization of the complete mitochondrial genome from the pancreatic fluke *Eurytrema pancreaticum* (Trematoda: Dicrocoeliidae). *Gene* 2016, 160–165.
- Coil WH** (1966) Egg shell formation in the notocotylid trematode, *Ogmocotyle indica* (Bhalerao, 1942) Ruiz, 1946. *Zeitschrift für Parasitenkunde* 27, 205–209.
- Darriba D, Taboada GL, Doallo R and Posada D** (2012) jModelTest 2: more models, new heuristics and high-performance computing. *Nature Methods* 9, 772.
- Dierckxsens N, Mardulyn P and Smits G** (2017) NOVOPlasty: de novo assembly of organelle genomes from whole genome data. *Nucleic Acids Research* 45, e18.
- Djuikwo-Teukeng FF, Kouam Simo A, Allienne JF, Rey O, Njayou Ngapagna A, Tchuem-Tchuente LA and Boissier J** (2019) Population genetic structure of *Schistosoma bovis* in Cameroon. *Parasites & Vectors* 12, 56.
- Eom KS, Rim HJ and Jang DH** (1984) A study on the parasitic helminths of domestic duck (*Anas platyrhynchos* var. *domestica* Linnaeus) in Korea. *Kisaengch'unghak Chapchi. The Korean Journal of Parasitology* 22, 215–221.
- Fang G, Liu H, Wu K, Wei T and Wang Q** (2022) Changing legislative thinking in China to better protect wild animals and human health. *Conservation Biology* 36, e13837.
- Gao JF, Zhao Q, Liu GH, Zhang Y, Zhang Y, Wang WT, Chang QC, Wang CR and Zhu XQ** (2014) Comparative analyses of the complete mitochondrial genomes of the two ruminant hookworms *Bunostomum trigonocephalum* and *Bunostomum phlebotomum*. *Gene* 541, 92–100.
- Gao JF, Mao RF, Li Y, Sun YY, Gao ZY, Zhang XG, Jin ZH, An Q, Zhang ZH, Zhang AH, Wei W, Lan Z and Wang CR** (2021) Characterization of the mitochondrial genome of *Tetrameres grusi* and insights into the phylogeny of Spirurina. *International Journal for Parasitology – Parasites and Wildlife* 17, 35–42.
- Goździewska-Harłajczuk K, Hamouzová P, Klečková-Nawrot J and Čížek P** (2021) The tongue of the red panda (*Ailurus fulgens fulgens* Cuvier, 1825) – a stereoscopy, light microscopy and ultrastructural analysis. *PeerJ* 9, e12559.
- Guo XR, Li Y, Gao Y, Qiu YY, Jin ZH, Gao ZY, Zhang XG, An Q, Chang QC, Gao JF and Wang CR** (2022) The complete mitochondrial genome of *Prosthogonimus cuneatus* and *Prosthogonimus pellucidus* (Trematoda: Prosthogonimidae), their features and phylogenetic relationships in the superfamily Microphalloidea. *Acta Tropica* 232, 106469.
- Hu L, Zhang M, Sun Y and Bu Y** (2020) Characterization and phylogenetic analysis of the first complete mitochondrial genome of *Cylicocycclus radia-tus*. *Veterinary Parasitology* 281, 109097.
- Huelsenbeck JP and Ronquist F** (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17, 754–755.
- Ichikawa-Seki M, Peng M, Hayashi K, Shoriki T, Mohanta UK, Shibahara T and Itagaki T** (2017) Nuclear and mitochondrial DNA analysis reveals that hybridization between *Fasciola hepatica* and *Fasciola gigantica* occurred in China. *Parasitology* 144, 206–213.
- Ivashko YI, Semenchenko AA, Solodovnik DA and Atopkin DM** (2022) Characterization of complete mitochondrial genome and ribosomal operon for *Carassotrema koreanum* Park, 1938 (Digenea: Haploporidae) by means of next-generation sequencing data. *Journal of Helminthology* 96, e54.
- Iwaki T, Okada T, Seki K, Izawa K and Sakurai F** (2012) *Ogmocotyle ailuri* (Price, 1954) (Digenea: Notocotylidae) found in the Japanese monkey, *Macaca fuscata*. *The Journal of Veterinary Medical Science* 74, 1211–1212.
- Kanev I, Radev V and Fried B** (2002) Superfamily Cyclocoeloidea Stossich, 1902. In Gibson DJ, Jones A, Bray RA (eds), *Keys to the Trematoda*, vol. 1. Wallingford: CABI Publishing and The Natural History Museum, pp. 147–152.
- Katoh K and Standley DM** (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30, 772–780.
- Le TH, Humair PF, Blair D, Agatsuma T, Littlewood DT and McManus DP** (2001) Mitochondrial gene content, arrangement and composition compared in African and Asian schistosomes. *Molecular and Biochemical Parasitology* 117, 61–71.
- Le TH, Nguyen KT, Nguyen NTB, Doan HTT, Agatsuma T and Blair D** (2019) The complete mitochondrial genome of *Paragonimus ohirai* (Paragonimidae: Trematoda: Platyhelminthes) and its comparison with *P. westermani* congeners and other trematodes. *PeerJ* 7, e7031.
- Le TH, Pham LTK, Doan HTT, Le XTK, Saijuntha W, Rajapakse RPVJ and Lawton SP** (2020) Comparative mitogenomics of the zoonotic parasite *Echinostoma revolutum* resolves taxonomic relationships within the 'E. revolutum' species group and the *Echinostomata* (Platyhelminthes: Digenea). *Parasitology* 147, 566–576.
- Li Y, Ma XX, Lv QB, Hu Y, Qiu HY, Chang QC and Wang CR** (2019) Characterization of the complete mitochondrial genome sequence of *Tracheophilus cymbius* (Digenea), the first representative from the family Cyclocoelidae. *Journal of Helminthology* 94, e101.
- Li J, Karim MR, Li J, Zhang L and Zhang L** (2020) Review on parasites of wild and captive giant pandas (*Ailuropoda melanoleuca*): diversity, disease and conservation impact. *International Journal for Parasitology – Parasites and Wildlife* 13, 38–45.
- Librado P and Rozas J** (2009) DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25, 1451–1452.
- Liu GH, Yan HB, Otranto D, Wang XY, Zhao GH, Jia WZ and Zhu XQ** (2014) *Dicrocoelium chinensis* and *Dicrocoelium dendriticum* (Trematoda: Digenea) are distinct lancet fluke species based on mitochondrial and nuclear ribosomal DNA sequences. *Molecular Phylogenetics and Evolution* 79, 325–331.
- Lohse M, Drechsel O, Kahlau S and Bock R** (2013) OrganellarGenomeDRAW – a suite of tools for generating physical maps of plastid and mitochondrial genomes and visualizing expression data set. *Nucleic Acids Research* 41, W575–W581.
- Ma J, He JJ, Liu GH, Zhou DH, Liu JZ, Liu Y and Zhu XQ** (2015) Mitochondrial and nuclear ribosomal DNA dataset supports that *Paramphistomum leydeni* (Trematoda: Digenea) is a distinct rumen fluke species. *Parasites & Vectors* 8, 201.
- Ma J, He JJ, Liu GH, Blair D, Liu LZ, Liu Y and Zhu XQ** (2016) Mitochondrial genome of *Ogmocotyle sikae* and implications for phylogenetic studies of the Notocotylidae trematodes. *Infection, Genetics and Evolution* 37, 208–214.
- Mioduchowska M, Czyż MJ, Goldyn B, Kur J and Sell J** (2018) Instances of erroneous DNA barcoding of metazoan invertebrates: are universal cox1 gene primers too 'universal'? *PLoS ONE* 13, e0199609.
- Olson PD, Cribb TH, Tkach VV, Bray RA and Littlewood DT** (2003) Phylogeny and classification of the Digenea (Platyhelminthes: Trematoda). *International Journal for Parasitology* 33, 733–755.
- Perna NT and Koehler TD** (1995) Patterns of nucleotide composition at four-fold degenerate sites of animal mitochondrial genomes. *Journal of Molecular Evolution* 41, 353–358.
- Pice EW** (1954) A new trematode from the lesser panda, *Ailurus fulgens*. *Journal of Parasitology* 40, 38–39.
- Rombel IT, Sykes KF, Rayner S and Johnston SA** (2002) ORF-FINDER: a vector for high throughput gene identification. *Gene* 282, 33–41.
- Sey O and Graber M** (1991) Description of *Ogmocotyle africanum* sp. n. (Trematoda, Notocotylidae) from the hippopotamus and a review of the genus *Ogmocotyle* Skrijabim et Schulz, 1933. *Acta Zoologica Hungarica* 37, 257–261.
- Sharma HP and Achhami B** (2022) Gastro-intestinal parasites of sympatric red panda and livestock in protected areas of Nepal. *Veterinary Medicine and Science* 8, 568–577.
- Shumenko PG and Tatonova YV** (2022) Assessing the population structure of trematode *Metagonimus suisfunensis* using three mitochondrial markers. *Parasitology Research* 121, 915–923.
- Solà E, Álvarez-Presas M, Frías-López C, Littlewood DT, Rozas J and Riutort M** (2015) Evolutionary analysis of mitogenomes from parasitic and free-living flatworms. *PLoS ONE* 10, e0120081.
- Stothard P** (2000) The sequence manipulation suite: JavaScript programs for analyzing and formatting protein and DNA sequences. *Biotechniques* 28, 1104.
- Suleman, Ma J, Khan MS, Tkach VV, Muhammad N, Zhang D and Zhu XQ** (2019) Characterization of the complete mitochondrial genome of *Plagiorchis maculosus* (Digenea, Plagiorchiidae), representative of a taxonomically complex digenean family. *Parasitology International* 71, 99–105.
- Suleman, Khan MS, Tkach VV, Muhammad N, Zhang D, Zhu XQ and Ma J** (2020) Molecular phylogenetics and mitogenomics of three avian

- dicrocoeliids (Digenea: Dicrocoeliidae) and comparison with mammalian dicrocoeliids. *Parasites & Vectors* **13**, 74.
- Suleman, Muhammad N, Khan MS, Tkach VV, Ullah H, Ehsan M, Ma J and Zhu XQ** (2021) Mitochondrial genomes of two eucotyliids as the first representatives from the superfamily Microphalloidea (Trematoda) and phylogenetic implications. *Parasites & Vectors* **14**, 48.
- Tamura K, Stecher G and Kumar S** (2021) MEGA11: molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution* **38**, 3022–3027.
- Valentyne H, Spratt DM, Aghazadeh M, Jones MK and Šlapeta J** (2020) The mitochondrial genome of *Angiostrongylus mackerrasae* is distinct from *A. cantonensis* and *A. malaysiensis*. *Parasitology* **147**, 681–688.
- Wang CR, Chen J, Zhao JP, Chen AH, Zhai YQ, Li L and Zhu XQ** (2009) *Orientobilharzia* species: neglected parasitic zoonotic agents. *Acta Tropica* **109**, 171–175.
- Wang XF, Zhang K, Zhang GW, Li ZY, Shang YX, Ning CC, Ji CH, Qiao J, Meng QL and Cai XP** (2022) Molecular characteristics and genetic diversity of *Fasciola hepatica* from sheep in Xinjiang, China. *Journal of Veterinary Research* **66**, 199–207.
- Wei F, Feng Z, Wang Z and Hu J** (1999) Current distribution, status and conservation of wild red pandas *Ailurus fulgens* in China. *Biological Conservation* **89**, 285–291.
- Wey-Fabrizius AR, Podsiadlowski L, Herlyn H and Hankeln T** (2013) Platyzoan mitochondrial genomes. *Molecular Phylogenetics and Evolution* **69**, 365–375.
- Xu G, Zhu P, Zhu W, Ma B, Li X and Li W** (2021) Characterization of the complete mitochondrial genome of *Notocotylus* sp. (Trematoda, Notocotylidae) and its phylogenetic implications. *Parasitology Research* **120**, 1291–1301.
- Yan HB, Wang XY, Lou ZZ, Li L, Blair D, Yin H, Cai JZ, Dai XL, Lei MT, Zhu XQ, Cai XP and Jia WZ** (2013) The mitochondrial genome of *Paramphistomum cervi* (Digenea), the first representative for the family Paramphistomidae. *PLoS ONE* **8**, e71300.
- Yang GY and Zhang ZH** (2013) *Parasitic Diseases of Wildlife*. Beijing: Science Press, pp. 324–327 (in Chinese).
- Yang X, Zhao Y, Wang L, Feng H, Tan L, Lei W, Zhao P, Hu M and Fang R** (2015) Analysis of the complete *Fischoederius elongatus* (Paramphistomidae, Trematoda) mitochondrial genome. *Parasites & Vectors* **8**, 279.
- Yang X, Wang L, Chen H, Feng H, Shen B, Hu M and Fang R** (2016) The complete mitochondrial genome of *Gastrothylax crumenifer* (Gastrothylacidae, Trematoda) and comparative analyses with selected trematodes. *Parasitology Research* **115**, 2489–2497.
- Yoshimura K, Hishinuma Y and Sato M** (1969) *Ogmocotyle ailuri* (Price, 1954) in the Taiwanese monkey, *Macaca cyclopis* (Swinhoe, 1862). *The Journal of Parasitology* **55**, 460.
- Zhang D, Li WX, Zou H, Wu SG, Li M, Jakovlić I, Zhang J, Chen R and Wang GT** (2019) Homoplasy or plesiomorphy? Reconstruction of the evolutionary history of mitochondrial gene order rearrangements in the subphylum Neodermata. *International Journal for Parasitology* **49**, 819–829.