

harmful use of illegal substances. Two patients had excessive amount of alcohol use to the point of dependence since teenage. Mean age of onset of both substance and alcohol use was 11, with range of 0 to 20. The most commonly misused substances were alcohol and cannabis (11 out of 12 patients). Eight patients had abused drugs other than cannabis. All but one of these then progressed to using other substances. The reasons for using substances and attitudes to substance and alcohol use were not explored in admission assessment in most cases.

**Conclusion.** Admission assessment to a rehabilitation ward is also an opportunity to screen for any barriers to recovery as well to use brief motivational interviewing intervention if appropriate clinically. There is a need to improve the quality of our admission assessment in relation to substance use history.

Most of our patients had a very early onset of alcohol and substance use, as young as age 8. Apart from one outlier, all had started using substances and alcohol by age 15. This raises concerns regarding missed early prevention and safeguarding opportunities.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

### Statistical Machine Learning Methods to Handle Missing PHQ-8 Score – Assuming Missing at Random

Mr Khalid Suliman<sup>1\*</sup>, Dr Mitha Al Balushi<sup>1</sup>, Prof Hannah Holliday<sup>2</sup>, Dr Manal Alblooshi<sup>1</sup> and Dr Amar Ahmad<sup>1</sup>

<sup>1</sup>Public Health Research Center, New York University in Abu Dhabi, Abu Dhabi, UAE and <sup>2</sup>New York University, New York, USA

\*Presenting author.

doi: 10.1192/bjo.2024.440

**Aims.** Missing data is a challenge that most researchers encounter. It is a concern that continues to be analyzed and addressed for solutions. Missing data occurs when there is no data stored for certain variables relating to participants. In health surveys, when participants answer in the form of “I don’t know” or “I’d prefer not to answer”, these responses can, in many cases, be categorized as missing data responses from a participant in a specific category or question.

The eight-item Patient Health Questionnaire (PHQ-8) is an essential tool in healthcare and clinical settings to assess an individual’s mental health, specifically related to symptoms of depression. The items are scored on a scale from 0 to 3 with the total score obtained by summing the scores for each item. Higher PHQ-8 scores indicate the presence of depressive symptoms.

We used empirical data from a previous study on depression symptoms in patients with coronary heart disease to study the effect of considering the answers “I do not know” and “I prefer not to answer” as missing values when estimating the percentage of depression using PHQ-8. Moreover, we studied the effect of the complete case analysis and multiple imputation on parameter estimates and confidence intervals. The outcome of this study aims to shed light on the development of missing data procedural knowledge and provide methodological support for public health decision-making when data with missing values are collected.

Furthermore, this study aims to prevent the exclusion of missing data rather than to generate data.

**Methods.** A simulation study with 1000 replicates was performed. Four common statistical machine learning methods for handling

missing values were included in this study. These are K-Nearest Neighbor (KNN), K-Means, Classification and Regression Trees (CART), and Random Forest (RF) imputations. Five clusters were used for KNN and K-mean. Likewise, five multiple imputations were used for the CART and RF methods. The simulation was based on publicly available data with available PHQ-8 data for 1096 subjects. In the simulation study and for each replication, multivariate missing values were generated using the missing-at-random (MAR) assumption with 10%, 20%, 30%, 40%, and 50% proportions of missingness. The percent of depression was calculated using the PHQ-8 questionnaire and a comparison was made between estimated actual depression, complete-case analysis, KNN, Kmean, RF, and CART, respectively.

**Results.** The Median age of the subjects was 69 (interquartile range: 61–67) and more males (72.9%) than females were included in the data. The estimated actual depression was 16.8, whereas the estimated percentage of depression varies between 6.9–13.5, 16.2–16.7, 16.3–16.7, 16.6–16.7 and 16.7–16.8 for the complete case, KNN, Kmean, RF and CART respectively.

**Conclusion.** The results of this simulation study show that missing PHQ-8 data are best handled by applying multiple imputations based on CART or RF. However, using K-Means or KNN leads to a good estimate of the true percentage of depression. Furthermore, the results of this simulation study show that complete-case analysis leads to biased estimates of the true percentage of depression. Nevertheless, further investigation is needed to address the problem of missing PHQ-8 data under the assumption of missing not at random.

Abstracts were reviewed by the RCPsych Academic Faculty rather than by the standard *BJPsych Open* peer review process and should not be quoted as peer-reviewed by *BJPsych Open* in any subsequent publication.

### Inpatient Psychiatry Placement Quality Improvement Project for Medical Students at University College London

Ms Maja Swirska<sup>1\*</sup>, Dr Mustafa Abbas<sup>2</sup>, Dr Petros Lekkos<sup>2</sup>, Prof Suzanne Reeves<sup>1,2</sup> and Dr Stephen Ginn<sup>1,3</sup>

<sup>1</sup>University College London Medical School, London, United Kingdom; <sup>2</sup>North London Mental Health Partnership, London, United Kingdom and <sup>3</sup>Central North West London NHS Trust, London, United Kingdom

\*Presenting author.

doi: 10.1192/bjo.2024.441

**Aims.** UCL 5th year medical students undertake 3-week North London Mental Health Partnership inpatient psychiatric wards placements.

Before this project the management of these placements was at the discretion of individual ward teams. A varied, and potentially unsatisfactory, medical student experience resulted.

This project sought to implement a structured approach to placements.

Ward teaching best practice was, for the purposes of this project, considered to be (i) sending students a welcome email prior to placement, (ii) issuing placement timetable, (iii) using tutorial materials for onward tuition.

Project aims: 75% of wards sending welcome email, 75% issuing timetables, 75% using tutorial materials, 75% of students stating placement exceeded expectations.

**Methods.** This project consists of 1 PDSA cycle.