

This study and previous reports suggest that the combined analysis of the MIE and a-sequence amplified products can be used for typing CMV strains. Although these regions represent relatively small samples of the whole viral genome, this PCR-based system was shown to be reproducible, stable, and discriminatory. The combined analysis of these two regions appears to be a reasonable alternative to traditional molecular analysis of CMV strains. This approach will facilitate epidemiological studies because it allows analysis of CMV strains directly from stored clinical samples and offers a rapid means for analyzing large numbers of specimens. RFLP analysis of PCR products also can be used as an important laboratory quality control procedure to differentiate wild CMV strains from laboratory control strains that might contaminate PCR reactions.

REFERENCES

1. Winston DJ, Huang ES, Miller MJ, et al. Molecular epidemiology of cytomegalovirus infections associated with bone marrow transplant. *Ann Intern Med* 1985;102:16-20.
2. Murph JR, Bale JF, Murray JC, Stinski ME, Perlman S. Cytomegalovirus transmission in a Midwest day care center: possible relationship to child care practices. *J Pediatr* 1986;109:35-39.
3. Willfert CM, Huang ES, Stagno S. Restriction endonuclease analysis of cytomegalovirus deoxyribonucleic acid as an epidemiologic tool. *Pediatrics* 1982;70:711-721.
4. Adler SP. The molecular epidemiology of cytomegalovirus transmission among children attending a day care center. *J Infect Dis* 1985;152:760-768.
5. Spector SA, Neuman TR, Hirata KK. Rapid determination of molecular relatedness of isolates of human cytomegalovirus. *J Infect Dis* 1985;152:755-759.
6. Huang ES, Alford CA, Reynolds DW, Stagno S, Pass RF. Molecular epidemiology of cytomegalovirus infections in women and their infants. *N Engl J Med* 1980;303:958-962.
7. Chou S. Acquisition of donor strains of cytomegalovirus by renal transplant recipients. *N Engl J Med* 1986;22:1418-1423.
8. Chou S. Differentiation of cytomegalovirus strains by restriction analysis of DNA sequences amplified from clinical specimens. *J Infect Dis* 1990;162:738-742.
9. Zaia JA, Galleg-Hawkins G, Churchill MA. Comparative analysis of human cytomegalovirus a sequence in multiple clinical isolates by using polymerase chain reaction and restriction fragment polymorphism assays. *J Clin Microbiol* 1990;28:2602-2607.
10. Bale JF, O'Neil ME, Fowler SS, Murph JR. Analysis of acquired human cytomegalovirus infections by polymerase chain reaction. *J Clin Microbiol* 1993;31:2043-2048.
11. Souza IE, Nicholson D, Matthey S, et al. Detection of human cytomegalovirus in peripheral blood leukocytes by the polymerase chain reaction and a nonradioactive probe. *Diagn Microbiol Infect Dis* 1994;20:13-19.
12. Haugen TH, Alden B, Matthey S, Nicholson D. Restriction enzyme fragment length polymorphism of amplified herpes simplex Type-1 DNA provide epidemiologic information. *Diagn Microbiol Infect Dis* 1993;17:129-133.
13. Winston DJ, Ho WG, Champlin RE. Cytomegalovirus infections after allogeneic bone marrow transplantation. *Infect Dis* 1990;12(suppl):776S-792S.
14. Williams SD, Kwok S. *Laboratory Diagnosis of Viral Infections*. 2nd ed. New York, NY: Marcel Dekker, Inc; 1992:147.
15. Souza IE, Greg A, Pfab D, et al. Intrafamilial CMV transmission: relationship to congenital infection. Presented at the 6th International Congress for Infectious Diseases; April 26-30, 1994; Prague, Czech Republic. Abstract 579.
16. Souza IE, Benson P, Murph J, et al. Polymerase chain reaction analysis of cytomegalovirus strains from congenitally-infected infants. Presented at the 34th Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC); October 4-7, 1994; Orlando, Florida. Abstract J21.
17. Sokol DM, Demmler GJ, Buffone GJ. Rapid epidemiologic analysis of cytomegalovirus by using polymerase chain reaction amplification of the L-S junction. *J Clin Microbiol* 1991;30:839-844.

Risk of HIV Transmission from Healthcare Worker to Patient Is Very Small

by Gina Pugliese, RN, MS
Medical News Editor

The CDC recently reported the results of several investigations from health departments, hospitals, and other agencies that had elected to notify patients who had received care from HIV-infected workers.

As of January 1, 1995, investigations of 64 healthcare workers infected with HIV had been reported to CDC; HIV test results were available for 22,171 patients of 51 of the 64 healthcare workers. For 37 of the 51 workers, no seropositive patients were reported among 13,063 patients tested for HIV. For the remaining 14

healthcare workers, 113 seropositive patients were reported among 9,108 patients. Epidemiologic and laboratory follow up did not show any healthcare worker to have been the source of HIV for any of the patients tested.

This report summarizes the largest existing database on retrospective evaluations of patients treated by healthcare workers infected with HIV. Excluding the previously described Florida dental practice, no transmission of HIV from healthcare workers has been documented among the 22,171 patients tested.

The CDC concluded that, despite limitations, these data are consistent with previous assessments that the risk for transmission of HIV from a

healthcare worker to a patient during an invasive procedure is very small.

Many of the retrospective investigations involved intensive effort and the expenditure of substantial resources by hospitals, state and local health departments, and the CDC. Failure to find additional instances of transmission to patients supports current recommendations that retrospective patient notification need not be done routinely, but should be considered on a case-by-case basis, taking into account an assessment of risks, confidentiality issues, and available resources.

FROM: Robert LM, Chamberland ME, Cleveland JL, et al. *Ann Intern Med* 1995;122:653-657.