

Mineralogy and space weathering found in the fine-grained samples returned from the C-type asteroid Ryugu

Takaaki Noguchi¹, the Hayabusa2-initial-analysis Sand team, some members from the Hayabusa2-initial-analysis Chemistry team, and the Hayabusa2-initial-analysis core.

¹ Kyoto University, Division of Earth and Planetary Science, Kyoto, Kyoto, Japan.

* Corresponding author: noguchi.takaaki.2i@kyoto-u.ac.jp

Introduction: The JAXA Hayabusa2 spacecraft returned samples from near-Earth C-type asteroid Ryugu. Our Hayabusa2-initial-analysis Min-Pet Fine (“Sand”) team has investigated the mineralogy and petrology and space weathering of grains in the fine-grained fraction of the samples. Space weathering is alteration induced mainly by solar wind irradiation and micrometeoroid impacts. Here, we report the mineralogy, microstructural, and chemical features related to space weathering of the Ryugu grains.

Samples and Methods: Surface morphologies of ~700 grains were observed by field emission scanning electron microscopy (FE-SEM) at Kyoto University and focused ion beam (FIB)-SEM at Kyushu University. Thin foil samples were prepared by FIB-SEM at both universities. A polished sample of a fragment originating from a large grain A0058 (~3 mm wide), which was allocated to the Hayabusa2-initial-analysis Chemistry team, was also investigated. We performed transmission electron microscopy (TEM), synchrotron radiation X-ray absorption fine structure (XANES and EXAFS) spectroscopy, nanotomography, and atom probe analysis at 16 universities and laboratories spread across the world. In addition, more than 200 grains are being investigated until the end of May, 2022.

Results and discussion: Mineralogy of fine-grained fraction of the Ryugu samples. Major minerals of the small Ryugu grains are saponite, serpentine, Fe-Ni sulfides, magnetite, dolomite, and breunnerite. The mineralogy and petrology of the Ryugu FIB sections investigated is similar to that of CI chondrites, apart for the lack of ferrihydrite and sulfates, which are commonly found in those meteorites [1]. Considering the effects of terrestrial weathering of CI chondrites, we infer that the mineralogy of investigated Ryugu grains represents that of CI chondrites prior to their arrival on Earth.

Space weathering of Ryugu samples. Recognizable surface modifications of the phyllosilicate-rich matrix were found on 6 to 7% of the observed grains. A variety of surface modifications are observed: melt splashes, amorphous layers, and melt layers. The amorphous layers form a continuous sheet ~0.1 μm thick composed of amorphous silicate material at the top surface. Their bulk chemical compositions are indistinguishable from those of the underlying phyllosilicate-rich matrix apart for a higher ratio of Fe²⁺ relative to Fe³⁺. The melt layers have bubbles and numerous submicroscopic (<100 nm) rounded Fe-Ni sulfide beads. These data suggest that both silicate and Fe-Ni sulfides were melted and then immiscibly separated into silicate and sulfide melts. Such melt layers have higher Fe and lower Si+Al and Mg ratios, as well as a higher ratio of Fe²⁺ relative to Fe³⁺ than the interior phyllosilicate-rich matrix.

The surface morphology of the amorphous layers resembles the surface of an experimental product of Orgueil CI chondrite that was irradiated by 4 keV He⁺. To a first approximation, it appears that solar-wind irradiation likely played an important role in modifying the space-exposed surface of the phyllosilicate-rich matrix. Structures of the melt layers resemble those of the products from the laser

irradiation experiments that simulate micrometeoroid impacts [2], suggesting that they had an important role in forming the melt layers. Only a small amount of nanophase (np) Fe⁰ was formed on the surface of the melt layer, which may be related to the high relative abundances of Fe³⁺ to Fe²⁺ in Ryugu grains compared to space weathered Itokawa and lunar grains. The H₂O in Ryugu phyllosilicates may also be responsible for this difference.

Given the assumptions that solar wind irradiation is rapid and that micrometeoroid impact processing is slow [3], Ryugu grains preserve a range of stages of space weathering. Some have a melt layer that covers an amorphous layer, others show the effects of multiple micrometeoroid impacts. Furthermore, some have blisters (bubbles formed near the surface) on the surface of a melt layer. These multiple space weathering textures suggest that effects of solar wind irradiation accumulate rapidly and the effects of micrometeoroid impact overprint later. In some cases, solar wind gaseous species implanted on the surface of a melt layer and later micrometeoroid impact heating may have induced formation of blisters on the surface. Of course, the natural overturn or gardening of regolith grains on the asteroid parent body interrupts these stages so that the weathering process on any single grain does not necessarily progress linearly.

The Hayabusa2-initial-analysis Sand team: Takaaki Noguchi, Toru Matsumoto, Akira Miyake, Yohei Igami, Mitsutaka Haruta, Hikaru Saito, Satoshi Hata, Yusuke Seto, Masaaki Miyahara, Naotaka Tomioka, Hope A. Ishii, John P. Bradley, Kenta Ohtaki, Elena Dobrică, Hugues Leroux, Corentin Le Guillou, Damien Jacob, Maya Marinova, Francisco de la Peña, Falko Langenhorst, Dennis Harries, Pierre Beck, Thi H. V. Phan, Rolando Rebois, Neyda M. Abreu, Jennifer Gray, Thomas Zega, Pierre-M. Zanetta, Michelle S. Thompson, Rhonda Stroud, Kate Burgess, Brittany A. Cymes, John C. Bridges, Leon Hicks, Martin R. Lee, Luke Daly, Phil A. Bland, Michael E. Zolensky, David R. Frank, James Martinez, Akira Tsuchiyama, Masahiro Yasutake, Junya Matsuno, Shota Okumura, Itaru Mitsukawa, Kentaro Uesugi, Masayuki Uesugi, Akimasa Takeuchi, Mingqi Sun, Satomi Enju, Aki Takigawa, Tatsuhiro Michikami, Tomoki Nakamura, Megumi Matsumoto, Yusuke Nakauchi.

Some members from the Hayabusa2-initial-analysis Chemistry team: Kazuhide Nagashima, Noriyuki Kawasaki, Naoya Sakamoto, Hisayoshi Yurimoto

The Hayabusa2-initial-analysis core: Shogo Tachibana, Tomoki Nakamura, Takaaki Noguchi, Hiroshi Naraoka, Ryuji Okazaki, Kanako Sakamoto, Hikaru Yabuta, Hisayoshi Yurimoto, Yuichi Tsuda, Sei-ichiro Watanabe.

References:

- [1] Brearley, A. J., Jones, R. H., in “Planetary Materials (Rev. Min. **36**)”, ed. J.J. Papike, (Mineralogical Society of America, City) p.C1.
- [2] Thompson, M. S. et al. *Icarus* **319** (2019), p. 499. doi: 10.1016/j.icarus.2018.09.022.
- [3] Vernazza, P. et al. *Nature* **458** (2009), p. 993. doi: 10.1038/nature07956.

[4] The authors acknowledge funding from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) (grants JPMXP09A21KU0393, JP19H00725, JP19KK0094, JP21H05424, JP20H01965, JP20H00198, JP20H00205, JP15H05695, JP16H06348, JP20H00188, Core-to-core program “International Network of Planetary Sciences”), the UK Science and Technology Facilities Council (STFC) (grant ST/T002328/1, ST/S000429/1), Purdue University (Start-up funds), University of Glasgow (COVID-5 19 Researcher Support Scheme), ISITE ULNE, Métropole Européenne de Lille, the region “Hauts-de-France”, and the European Research Council (ERC) (grant 771691).