




RESEARCH ARTICLE

A quasi-cohort trend analysis of adult obesity in Colombia

Paula Andrea Castro-Prieto^{1,2} , Jeroen Spijker²  and Joaquín Recaño^{1,2} 

¹Universitat Autònoma de Barcelona, Barcelona, Spain and ²Centre d'Estudis Demogràfics (CED-CERCA), Bellaterra, Spain
Corresponding author: Paula Andrea Castro-Prieto; Email: pcastro@ced.uab.es

(Received 29 July 2022; revised 8 May 2023; accepted 18 May 2023; first published online 26 June 2023)

Abstract

In Colombia, the prevalence of obesity has been increasing in recent years due to changes in dietary and nutritional patterns. While previous studies have focussed on describing obesity and its associated factors, they have mainly used a cross-sectional methodology. Accordingly, this study aims to conduct a descriptive quasi-cohort analysis to capture age-specific cohort trends in body mass index (BMI) according to sex and ethnicity (indigenous, Afro-Colombian, and the remaining population). The study utilised data from the National Survey of the Nutritional Situation in Colombia (ENSIN) conducted in 2005, 2010, and 2015 that included 214,136 individuals aged 20–64 years after screening. Data on ethnicity were only available from the 2010 and 2015 surveys. Overall, the prevalence of obesity increased by 6.1 percentage points (from 15.2% to 21.3%) between 2005 and 2015 (men from 10.4% to 15.7%; women from 18.2% to 25.7%). Among Afro-Colombians, obesity rose 6.6 percentage points (from 19.4% to 26.0%), again more so in women than in men (2015: 35.2% versus 17.8%). Among indigenous people, the proportion increased by 5.3 percentage points (from 13.5% to 18.8%), with women reporting highest rates (2015: 23.7% against 12.6% in men). Age- and cohort-specific results also indicate that recent adult cohorts are experiencing sharp increases in BMI, for example, while 25–29-year-old males born in 1975–1979 had a BMI of 24.2 kg/m², among 40–44-year-olds of the same cohort, this equalled 26.8 kg/m². In the case of women, these age differences in BMI among the same cohort are even greater (24.4 and 28.0 kg/m²). In summary, the results of this study indicate that Colombia is still in the early stages of the obesity transition, urging the need to monitor obesity trends in Colombia from both an age and cohort perspective. To achieve this, longitudinal surveys or repeated cross-sectional surveys like the ENSIN could be utilised.

Keywords: Obesity; Demography; Population Health; Colombia

Introduction

Excessive body weight is a risk marker for cardiovascular disease and is associated with an increased prevalence of diabetes, hypertension, metabolic syndrome, and premature mortality (Bozkurt et al., 2016) and is also a potential mediator of female infertility and cancer (Iyengar et al., 2016; Broughton & Moley, 2017). In addition, the rising incidence and prevalence of obesity constitute an important issue in global public health in terms of economic burden (Tremmel et al., 2017). Although obesity trends differ greatly between and within countries, according to Jaacks et al. (2019) there are discernible patterns in broad changes in obesity prevalence over time. These patterns can be grouped into predictable stages of, what the authors call, the ‘obesity transition’. The first stage is characterised by an increase in obesity from very low initial levels, especially in

women. The second stage corresponds to a more generalised increase in obesity in adults, and smaller increases in children, leading to a narrowing of the gender gap. According to Jaacks *et al.* (2019), many Latin American and Middle Eastern countries are presently at this stage, although, as our results show, in Colombia gender differences are not yet declining. The third stage is identified by the closing of the gender gap and an acceleration of obesity prevalence in the low socioeconomic status (SES) subpopulation that still had a low body mass index (BMI) during the previous stage to the point that they surpass the high SES population. The obesity prevalence in children also increases. The USA and all European countries that were analysed pertained to this stage in 2016 (Jaacks *et al.*, 2019).

Current evidence from Europe suggests a stabilisation of obesity prevalence. In the case of Austria, this has been shown for women since 2007 (Großschädl & Stronegger, 2021). In Scotland, while there was an increase in BMI between 1995 and 2008 among adults, it stayed level between 2008 and 2014 (Tod *et al.*, 2017). In Spain, the evidence shows a stabilisation in the excess body weight in men since 2017 and in women since 1997 (García & Martín, 2022).

However, recent studies in Latin America have observed an increase in the prevalence of overweight, particularly obesity, which has been unevenly distributed according to SES and gender. A study of 14 Latin American countries found that on average the highest prevalence of obesity was found in the fourth wealth quintile (26.1%), the third education quintile (27.1%), and urban areas (26.0%) (Jiwani *et al.*, 2019). Furthermore, the ethnic dimension implies an additional issue in Latin America, due to its diverse socio-cultural and economic conditions and characteristics. Afro-descendants and indigenous people are an examples of this, as they have experienced situations of extreme poverty, lower levels of education, poorer nutrition and health, in addition to being victims of crime, violence, and forced displacement (CEPAL, 2014; Freire *et al.*, 2018)

Due to its social, ethnic, and geographical diversity, Colombia is therefore an interesting case for the study of obesity trends making it also a potential model for countries with similar characteristics. According to the Nutritional Situation Survey (ENSIN, by its acronym in Spanish), obesity in adults has been increasing in recent years. In 2005, 13.7% of the adult population (men: 8.8% and women: 16.6%) suffered from obesity (ICBF, 2006). In 2010, the figure rose to 16.5% (men: 11.5% and women: 20.1%) (Profamilia, 2011) and by 2015 it stood at 18.7% (men: 14.4% and women: 22.4%) (Ministerio de Salud y Protección Social de Colombia, 2019). The information available at the ethnic dimension indicates that the highest levels of obesity are found in the Afro-Colombian population and lowest in the indigenous population. In 2010, 18.2% of Afro-Colombians were found to suffer from obesity, while the figure for indigenous people was 15.1%, compared to 16.5% for Colombia as a whole (Profamilia, 2011). In 2015, obesity had increased to 22.9% in the case of Afro-Colombians and remained stable among indigenous people (14.9%) as well as for the entire population (18.7%) (Ministerio de Salud y Protección Social de Colombia, 2019). The ENSIN has been used in different studies for food and nutrition analysis in Colombia (Flórez Pregonero *et al.*, 2012; Kasper *et al.*, 2013; Herrán *et al.*, 2016; Vecino-Ortiz & Arroyo-Ariza, 2018). However, descriptive quasi-cohort analyses have not been conducted, even though they have been used in demographic and other fields elsewhere when longitudinal data were not available (Preston & Wang, 2006; Cámara & Spijker, 2010; Suissa, 2015; Waite, 2015; Qian *et al.*, 2017; Pensiero & Green, 2018; García & Martín, 2022). One important advantage of the quasi-cohort approach is that allows cohort trends and differences between age groups to be analysed simultaneously, while standard cross-sectional analyses may mask generation-specific life course experiences related to nutrition (Cámara & Spijker, 2010). The aim of our research is therefore to perform a descriptive quasi-cohort analysis to capture age and cohort trends in BMI for different social and ethnic segments of the population in Colombia.

Table 1. Sample size of the ENSIN according to subgroup and age group used in the analysis

Survey year	Individuals		Households	Segments	Primary sampling units
	0–64	20–64	0–64	0–64	0–64
2005	156076	89740	17740	1920	209
2010	202253	119677	50670	4987	258
2015	136138	81501	44000	4379	295

Source: ENSIN 2005–2015.

Methods

Study design and data source

National Survey of the Nutritional Situation (ENSIN)

The ENSIN survey has been conducted every five years since 2005 and currently the microdata of three editions are publicly available (2005, 2010, and 2015). The ENSIN survey contains a representative sample of the Colombian population aged 0–64 years. It collects data on food security, nutritional status by anthropometry and biochemical indicators, and information on the 24-hour dietary recall and food frequency questionnaires (Ministerio de Salud y Protección Social de Colombia, 2019); however, the last has not been widely disseminated or published.

The sample constructed for this survey is probabilistic by multi-stage sampling. In the ENSIN 2005 and 2010, the rural population of Orinoquía and Amazonia, which corresponds to less than 1% of the country's population, was not included. In turn, the ENSIN only started to include the variable of ethnicity since 2010. The sample sizes according to each sampling unit level are described in Table 1.

For our analysis, we used the demographic information (age, sex) of the respondents aged 20–64 years born between 1940–1944 and 1995–1999 as well as their anthropometric components height and weight to calculate BMI. This age threshold was previously applied by other national and international studies and surveys (Cámara & Spijker, 2010; Briceño et al., 2012; García & Martín, 2022). Specifically, our study covers the period 2005–2015 and includes 214,136 individuals after screening (see Table 2). The respondents were aggregated into one database and subsequently designated into quasi birth-cohorts by subtracting their age from the survey year.

Cross-sectional and quasi-cohort approaches:

In Colombia, the main data source for data on nutrition is the ENSIN, which is a cross-sectional rather than a panel survey, as it does not follow the same subjects over time. When longitudinal data are absent, one alternative is merge existing cross-sectional data as this enables cohort differences in age-related changes to be analysed (Thomas, 2018). Given the large sample size in each ENSIN survey, we were able to apply a quasi-cohort to study trends in the nutritional indicators weight, height, and BMI. A quasi-cohort approach provides a more nuanced understanding of the complex relationship between sex, age, and nutritional indicators not only over time but also across different birth-cohorts as they age, thereby aiding the construction of more precise evidence-based policy decisions aimed at reducing the prevalence of overweight and obesity and its associated health risks.

Variables

The anthropometric nutritional status indicator BMI was obtained by taking the ratio of weight in kilograms over height in metres squared. Before calculating the BMI of the respondents, the data

Table 2. Number of after-screening sample cases used in the analysis by sex, age, and quasi birth-cohort

Quasi birth-cohort	Age group								
	20–24	25–29	30–34	35–39	40–44	45–49	50–54	55–59	60–64
Males									
1940–1944									1040
1945–1949								1388	1891
1950–1954							1788	2290	2021
1955–1959						2065	2760	2340	743
1960–1964					2409	3358	2939	887	
1965–1969				2541	3717	3164	1050		
1970–1974			2604	3774	3282	1078			
1975–1979		2944	3977	3173	1105				
1980–1984	3560	4494	3700	1111					
1985–1989	5055	3891	1273						
1990–1994	4421	1383							
1995–1999	1489								
Females									
1940–1944									1517
1945–1949								2139	2395
1950–1954							2791	3051	2517
1955–1959						3735	4194	3074	947
1960–1964					3886	4728	3734	1228	
1965–1969				4299	5146	4007	1393		
1970–1974			4302	5181	4141	1335			
1975–1979		4584	5293	4395	1366				
1980–1984	5096	5773	4638	1557					
1985–1989	6011	4743	1610						
1990–1994	5189	1721							
1995–1999	1715								

Source: ENSIN 2005–2015. Own calculations.

were screened on omitted values in height and/or weight (52,275 cases), errors, and unusual values (weight less than 30 kg (35 cases) and height less than 130 cm (116 cases)). This led to the removal of 52,426 cases (see also Appendix Table A1). Finally, additional 1054 cases were also removed because they were in an age-cohort category that contained too few cases to be analysed. In other words, a total of 53,480 cases were not included in the study (20.0% of all cases in the selected age and birth-cohort range).

It is also important to note that 68.3% of the cases eliminated were men. This could be due to two reasons. Firstly, in Colombia men are more likely to be employed than women (DANE, 2023) and therefore not at home at the time of the survey. Secondly, taking into account that women are the main caregivers of children under 12 and the primary source of information about their

children's nutrition, it was mandatory during the data collection phase to make initial contact with them rather than with men (ICBF, 2015).

While BMI is perhaps the most commonly used indicator to assess the prevalence of obesity among the adult population, in part due to its precision, accuracy, and validity, one important criticism is the lack of information on body fat mass and fat location (Nuttall, 2015), characteristics that are known to differ between population groups due to its relation to body shape (Norgan, 1994). However, BMI is widely used by the World Health Organization (2021) and health ministries in different countries (Ministerio de Salud y Protección Social, 2016; Estado Plurinacional de Bolivia, 2017; INEGI, 2018), especially in lower-middle-income countries, because of its low cost, simple acquisition, and comparability.

Hence, we analysed 20–64-year-olds according to 5-year age groups, 5-year birth-cohorts, sex (male and female), and for the 2010 and 2015 data, ethnicity, distinguishing between the indigenous, Afro-Colombian, and remaining population (labelled as *others*, the majority of whom are of mixed ethnicity).

We analysed both the continuous values of BMI and the standard BMI categories underweight (levels below 18.5 kg/m²), normal weight (values between 18.5 kg/m² and 24.9 kg/m²), overweight (levels between 25 and 29.9 kg/m²), and obesity (values above 30 kg/m²) (World Health Organization, 2021).

Analysis

This was a descriptive study, starting with an analysis of the weight by survey year, sex, and age group to understand its evolution in Colombia between 2005 and 2015. We then analysed weight gain by quasi birth-cohort and sex to discern the age-specific trend within each quasi birth-cohort. We also analysed the trend in height of the 1940–1944 to 1995–1995 quasi birth-cohorts by ethnicity as weight increases may partly be explained by changes in height across the different generations. Finally, to control for changing height we conducted an analysis of BMI by age group, sex, and survey year, as well as by age group, quasi birth-cohort, and ethnicity. We also performed significance tests on mean weight by quasi birth-cohort and age group.

The statistical program R was used to process and perform the descriptive analyses and for the construction of the graphs. The code is available upon request.

Results

Cross-sectional analysis of average weight by sex and age

Between 2005 and 2015, the average weight of Colombian adult men aged 20–64 years increased from 69.7 kg to 73.2 kg (+3.5 kg, equal to an increase of 4.7%). In women, the increase was from 62.3 kg to 65.9 kg (i.e. +3.6 kg and +5.4%, respectively). As Figure 1 shows, for both men and women and all age groups analysed, similar increases were observed between 2005 and 2010 as between 2010 and 2015, whereby increases over the 10-year period were statistically significant at the 95% level, except for the 20–24-year age group in the case of men and the 45–49, 50–54, and 55–59-year age groups in the case of women (see also Appendix Table A2). The highest increases between the first and last survey were observed for the 35–39-year age group among men (+4.6 kg) and 45–49-year age group among women (+2.7 kg).

Quasi birth-cohort analysis of average weight by age and sex

Figure 2a and 2b shows the evolution in average body weight experienced by people belonging to quasi birth-cohorts at different ages. In men, in almost all age groups each quasi birth-cohort reported an average weight increase over the previous quasi-cohort. For example, in the 50–54-

year age group between the 1950–1969 quasi-cohorts, weight increased from 70.2 kg to 74.4 kg (+4.2 kg). In women, the results indicate a similar trend to that of men. For example, in the same age and quasi-cohort group, the weight increased from 65.2 to 68.0 kg, indicating an increase of 2.8 kg, that is, 1.4 kg less. The results also clearly show that the slope of average weight changes between two five-year age groups in the same quasi-cohort is steeper in the younger quasi-cohorts.

Quasi-cohort analysis of average height by quasi birth-cohort and sex

Figure 3 presents the evolution of the average height experienced by people belonging to each quasi birth-cohorts by sex. Successive male and female birth-cohorts saw their average height increase. For instance, between the five-year cohorts 1940–1944 and 1995–1999, the increase in men was 5.6 cm and in women 5.8 cm.

It should be noted that the above reflects the national average height of the Colombian population. However, when calculating the average height in Afro-Colombian and indigenous communities, the figure changes considerably. While only aggregated data on ethnicity were available for the 2010 and 2015 surveys, Afro-Colombian men born in 1995–1999 were, on average, 3.5 cm taller than those born in 1945–1949 (171.6 cm vs. 168.1 cm). In the case of women, height increased by 5.2 cm (154.7–159.5 cm). On average, indigenous men born in 1995–1999 were 3.7 cm taller than the 1945–1949 cohort (162.6 cm vs. 158.9 cm). In the case of women, it increased by 4.7 cm (147.3–152.0 cm). Lastly, for the remainder of the Colombian population, men born in 1995–1999 were 5.4 cm taller on average than those born in 1945–1949 (169.8 cm vs. 164.4 cm). In the case of women, height increased by 5.8 cm (157.1–151.3 cm).

Table 3 shows the evolution of height by ethnicity by age group for the aggregated samples (2010 and 2015). The results indicate the younger the taller, with Afro-Colombians being the tallest on average whereas indigenous people are the shortest. The remaining group has an average height that is in-between the two ethnic groups.

Cross-sectional analysis of BMI by sex and age

The average BMI in both males and females also increased between the studied periods (Figure 4a and 4b). In men, this was most evident in the 35–39-year age group (from 25.3 kg/m² in 2005 to 26.7 kg/m² in 2015), but all age groups observed notable rises in BMI. Among women, the highest increase was observed in the 55–59-year age group (from 27.8 kg/m² in 2005 to 28.4 kg/m² in 2015).

When analysing the BMI categories, the proportion of the population with normal weight decreased by 9.9 percentage points between 2005 (47.7%) and 2015 (37.8%), with similar decreases occurring among both sexes. In men, the decrease was 10.7 percentage points (from 53.0% to 42.3%) and in women 10.1 (from 44.3% to 34.2%). In men, the decline was concentrated among ages 35–59 years and in women among 20–44-year-olds (Figure 5) (see also Appendix Table A3).

Regarding the proportion of the overweight population, between 2005 and 2015 it increased from 33.8% to 38.9% (i.e. 5.1 percentage points). In men, this was 6.3 percentage points (from 33.6% to 39.9%) and in women 4.2 (from 33.9% to 38.1%). In terms of obesity, between 2005 and 2015 it increased by 6.1 percentage points (from 15.2% to 21.3%). In contrast to overweight, the proportion of obese men was lower compared to women. By 2005, 10.4% of men suffered from obesity, in 2010 12.9%, and in 2015 15.7%. The respective figures for women were 18.2% in 2005, 22% in 2010, and 25.7% in 2015, which translates to an increase of 7.5 percentage points. Similar to men, obesity was highest in the 55–59-year age group, which in this case exceeded 30%.

At the ethnic level, information was only available from the ENSIN 2010 and 2015. In Afro-Colombians, normal weight decreased from 44.3% to 36.5%, while overweight increased from 33.1% to 35.8%, and obesity from 19.4% to 26.0%. Overweight and obesity were more prevalent in




Figure 1. Weight by age group and survey year. Source: ENSIN 2005–2015. Own calculations.




Figure 2. Weight by sex, age group, and quasi-cohort. Source: ENSIN 2005–2015. Own calculations.




Figure 3. Evolution of height according to quasi-cohort. Source: ENSIN 2005–2015. Own calculations.

Table 3. Mean height by ethnicity and age group

Age group	Sex	Afro-Colombian	Indigenous	Others	Total
20–24	Men	171.5	162.8	169.3	167.9
25–29		170.8	162.9	169.1	167.6
30–34		170.9	162.9	168.7	167.5
35–39		170.4	162.2	168.0	166.9
40–44		170.0	162.8	167.6	166.8
45–49		169.6	161.2	167.0	165.9
50–54		168.8	161.0	166.3	165.4
55–59		168.3	161.5	165.5	165.1
60–64		167.9	160.2	164.2	164.1
Total		169.8	161.9	167.3	166.3
20–24	Women	159.1	151.7	156.8	155.9
25–29		158.6	151.2	156.4	155.4
30–34		158.7	151.2	155.9	155.3
35–39		158.2	150.7	155.5	154.8
40–44		157.9	150.4	155.0	154.4
45–49		157.4	150.4	154.5	154.1
50–54		156.7	149.6	153.7	153.3
55–59		156.6	148.8	152.8	152.7
60–64		155.2	147.8	151.9	151.6
Total		157.6	150.2	154.7	154.2

Source: ENSIN 2010–2015. Own calculations

women than in men in 2010: 34.3% of women vs. 31.6% of men were overweight and for obesity it was, respectively, 24.9% and 12.5%. In 2015, however, overweight increased only slightly among women (to 35.3%) but more so among men (36.2%), although sex differences in obesity increased (35.2% vs. 17.8%). Proportions were especially high in the 45–49-year age group (see also Appendix Figure A1).

Results for indigenous people indicated a decrease in normal weight by 9.6% from 49.3% to 39.7%, an increase in overweight from 35.7% to 40.2%, and in obesity from 13.5% to 18.8%. Similar to Afro-Colombians, women reported the highest figures for overweight. In 2010, the proportion overweight was 36.9% vs. 34.2% for men, while 18.1% of women and 8.3% of men were obese. In 2015, the proportion overweight increased to 39.6%, while the increase was even greater among men, as they surpassed women in overweight (41.1%). Obesity was still close to twice as prevalent among women (23.7% vs. 12.6%). Regarding age, the highest proportion of obese men in 2015 was found among 60–64-year-olds, while among women it is the group aged 40–44 years (see also Appendix Figure A2).

Concerning subjects with another ethnicity (mainly of mixed ancestry), the decrease in normal weight was five percentage points (from 42.8% to 37.8%), while overweight and obesity both increased close to three percentage points (respectively, from 36.4% to 39.1% and from 18.5% to 21.0%). Being overweight was more common in males than in females, but both sexes observed




Figure 4. BMI by sex, age, and survey year. Source: ENSIN 2005–2015. Own calculations.

small increases between the two surveys. In 2010, the proportions were 36.9% vs. 35.9%, respectively, while in 2015 it was 40.2% for men against 38.2% for women. In the case of obesity, it was higher in women than in men (22.1% vs. 13.6% in 2010) and increased slightly during the 5-year period (25.1% vs. 15.8% in 2015). The ages most affected by obesity in 2015 were 55–59-year-olds (both sexes) (see also Appendix Table A4 and Figure 3).




Figure 5. Nutritional status by sex, age group, and survey year. Source: ENSIN 2005–2015. Own calculations.




Figure 6. BMI trends by sex, quasi birth-cohort, and age. Source: ENSIN 2005–2015. Own calculations.

BMI trends by quasi birth-cohort, age, and sex

The increasing height of successive birth-cohorts can only explain part of the increasing weight of the Colombian population between 2005 and 2015 as within the same quasi-cohort the average BMI increased as individuals became older, except for the 1950–1954 and 1945–1949 cohorts. As Figure 6 and Table 4 show, especially the most recent adult cohorts are experiencing sharp increases in BMI. For instance, while 25–29-year-old males born in 1975–1979 had a BMI of 24.2 kg/m², among 40–44-year-olds of same cohort this equalled 26.8 kg/m². In the case of women, these age differences in BMI among the same cohort are even greater (24.4 kg/m² and 28.0 kg/m²). But even the 1955–1959 cohort has seen its BMI increase over age by about 1 kg/m².

Interestingly, if we consider the oldest age of each cohort analysed (which would translate to the most recent period studied), we observe very few age differences in BMI between ages 35 and 64 in the case of men and ages 45 and 59 in the case of women. This is in part because the younger cohorts experienced very large increases in BMI during the 10-year period studied. As this occurred among all but the oldest cohorts, Figure 6 also clearly shows the cohort differences in BMI at each age group. For instance, the average BMI of a male aged 35–39 born in 1965–1969 was 25.3 kg/m², compared to 26.9 kg/m² for those of the same age but born in 1980–1984. In the case of women of the same age, cohort differences are less but still substantial (respectively 26.2 kg/m² and 27.3 kg/m²).

BMI trends at the ethnicity level (based on the 2010 and 2015 surveys) indicate that Afro-Colombian men reported an increase of up 1.2 kg/m² in the 40–44-year age group between the quasi birth-cohorts 1965–1969 and 1975–1979. In Afro-Colombian women, the highest values were found in the 45–49-year age group between the 1960–1964 and 1970–1974 quasi birth-cohorts (+1.3 kg/m²).

In indigenous men, the average BMI of 55–59-year-olds born in 1960–1964 was 1.3 kg/m² higher than in the 1960–1964 quasi birth-cohort. In indigenous women, the largest increase was observed among the 40–44-year age group between the 1965–1969 and 1970–1974 cohorts (+1.2 kg/m²). In other ethnicities (mainly of mixed ethnicity), in men the highest increase in average BMI was found in the 35–39-year age group between the 1970–1974 and 1980–1984 quasi birth-cohorts (+0.8 kg/m²). In women, this was among 60–64-year-olds, where the average BMI of the 1955–1959 cohort was 0.5 kg/m² higher than that of the 1945–1949 cohort (see Figure 7).

Table 4. 95% confidence intervals of BMI by sex, age group, and quasi birth-cohort

Quasi birth-cohort	Age group								
	20–24	25–29	30–34	35–39	40–44	45–49	50–54	55–59	60–64
Males									
1940–1944									25.29–25.74
1945–1949								25.44–25.89	25.49–25.85
1950–1954							25.44–25.83	25.82–26.17	25.85–26.22
1955–1959						25.6–25.95	25.85–26.16	26.12–26.46	26.37–27.01
1960–1964					25.38–25.70	25.98–26.27	26.35–26.66	26.51–27.05	
1965–1969				25.19–25.50	26.03–26.29	26.42–26.72	26.46–26.99		
1970–1974			24.87–25.18	25.72–25.97	26.34–26.62	26.59–27.10			
1975–1979		24.06–24.34	25.48–25.74	26.38–26.68	26.52–27.04				
1980–1984	22.92–23.16	24.75–24.99	25.86–26.13	26.67–27.18					
1985–1989	23.28–23.48	24.99–25.25	25.52–25.99						
1990–1994	23.58–23.81	24.65–25.10							
1995–1999	22.83–23.17								
Females									
1940–1944									27.12–27.65
1945–1949								27.62–28.06	27.79–28.22
1950–1954							27.61–27.98	27.97–28.33	27.84–28.24
1955–1959						27.39–27.70	28.01–28.32	28.3–28.66	28.41–29.06
1960–1964					26.59–26.89	27.63–27.91	28.03–28.35	28.12–28.70	
1965–1969				26.07–26.35	27.18–27.45	27.96–28.27	28.14–28.70		
1970–1974			25.36–25.63	26.61–26.87	27.68–27.99	28.08–28.62			
1975–1979		24.25–24.51	26.06–26.32	27.33–27.63	27.72–28.28				
1980–1984	23.08–23.30	24.95–25.19	26.62–26.90	27.05–27.58					
1985–1989	23.65–23.86	25.65–25.93	26.53–27.03						
1990–1994	24.33–24.59	25.38–25.83							
1995–1999	23.86–24.30								

Source: ENSIN 2005–2015. Own calculations.

Discussion

Based on data from the National Survey of the Nutritional Situation (ENSIN) from 2005, 2010, and 2015, we were able to observe an increase in height, weight, and BMI across successive generations of working age and born between 1945 and 1999. Specifically, regarding BMI, 38.9% of Colombians aged 20–64 years were overweight in 2015, up from 33.8% in 2005. Regarding obesity, 21.3% of Colombians were obese in 2015 compared to 15.2% in 2005, that is, an increase of more than 40%.

Increases were found among adults of successive cohorts at each 5-year age category that was analysed. In the case of men, the largest increase across successive generations occurred at age 35–39 (+1.6 kg/m² between the 1965–1969 and 1980–1984 quasi birth-cohorts). In the case of women, the largest increase across successive generations occurred at age 60–64 (1.4 kg/m² between the 1940–44 and 1955–1959 quasi birth-cohorts). In general, the 20–24-year age group born between 1980–1984 and 1995–1999 observed the lowest levels of BMI, whereby the youngest cohort of men even observed a lower average BMI (−0.4 kg/m²), while for women the average level was still increasing among younger quasi birth-cohorts (+0.8 kg/m²).

The evidence provided here goes in line with increases observed in other Colombian studies. For instance, Herrán *et al.* (2016) noted that overweight changed from 36.4% in 2005 to 37.6% in 2010, and Kasper *et al.* (2013) indicated that the prevalence of obesity increased from 13.9% in 2005 to 16.4% in 2010. Our study also found that obesity is more prevalent in women than in men and increases with age, but whereby sex differences were still not converging, which is consistent with the literature and places Colombia towards the end of stage one of the obesity transition (Jaacks *et al.*, 2019).

In addition to analyzing a more recent period (2015), our study incorporated an ethnic component, which distinguishes it from previous research on overweight and obesity in Colombia. Increases in BMI occurred among all three ethnic groups, although results also indicated that the prevalence of obesity was higher in people of Afro-Colombian communities. This insinuates parallels to black races in other countries such as the USA (Hales *et al.*, 2017; McTigue *et al.*, 2002).

Potential factors involved

In terms of age-specific increases in (over)weight, the weight of Colombian men stabilises after age 35–39. For instance, 35–39-year-olds have a similar average BMI as 40–44-year-olds and 60–64-year-olds when we consider the most recent cohort. On the other hand, in the case of women, our results showed that among all but the oldest cohorts, average BMI kept increasing with age. This sex difference can, at least partly, be explained by genetic and physiology factors. Within genetics, a meta-analysis of waist-to-hip ratio developed by genome-wide associations studies found that, when adjusted for total fat in more than 200,000 individuals, 20 of 49 loci identified showed sex-specific effects, and 19 of these had stronger effects in women (Zore *et al.*, 2018). Physiologically, women have a higher percentage of body fat and a lower percentage of fat-free mass than men (Fu, 2019) as well as a low prevalence of exclusive breastfeeding (Victora *et al.*, 2016) which is associated with postpartum weight loss (da Silva *et al.*, 2015). In addition, after menopause, women accumulate fat in subcutaneous area facilitated due to a decrease oestrogen, which makes them more prone to central obesity than men (Fu, 2019).

Moreover, there are also other potential factors involved in the increase of overweight and obesity in Colombian adults, in particular dietary, nutritional, historical, social, and political factors. Among the dietary factors is calorie intake, which has increased from an average of 1950 calories in 1960 (Bourges *et al.*, 2000) to 2117 calories in 2015 (Herrán *et al.*, 2021). Similarly, ultra-processed foods, especially junk food (*chatarra*), increasingly form part of the diet of Colombians (Bejarano-Roncancio *et al.*, 2015; García & Contreras, 2022).




Figure 7 BMI trends by sex, quasi birth-cohort, age, and ethnicity. Source: ENSIN 2005–2015. Own calculations.

Research indicates that in some indigenous communities the diet is based on carbohydrates due to food scarcity (Hernández et al., 2014). For example, the Guambiana community located in the southwest of Colombia bases its diet on preparations based on corn, ullucos (type of tubercle), and beans (Molano & Molano, 2018). A study that determined traditional foods in indigenous and Afro-Colombian communities in ten Colombian departments found that of the 92 foods reported, 39 came from plants and only 18 were classified as meat (Rivas et al., 2010). Some approaches have quantified the level of some macronutrients and micronutrients in the diet, reporting low levels of protein and vitamin A adequacy, such as the indigenous Tules in Antioquia (Carmona et al., 2005). In other cases, qualitative research has described how indigenous people have been immersed in the promotion of industrialised foods through the media, which affects their food consumption (Farfán et al., 2019).

Regarding nutritional factors, the so-called double nutritional burden, defined as the manifestation of undernutrition and overweight simultaneously (Popkin et al., 2020), can occur at different levels: population, household, and even on an individual level (Shrimpton & Rokx, 2012). In a Colombian study conducted in 2010, it was found that this double burden occurred in 4.6% of households, that is, the coexistence of chronic malnutrition in a child under five years of age and an overweight mother. The prevalence increased to 4.9% in indigenous communities (Rueda, 2019). In 2015, in rural Colombia, 7.8% of households were double burdened (Sansón-Rosas et al., 2021).

Previous studies have also linked physical inactivity and sedentary lifestyles to obesity in Colombia, primarily related to watching television (Lear *et al.*, 2014) and the use of private motor vehicles, especially motorcycles, where time spent riding a motorcycle per week could exceed 150 minutes (Parra *et al.*, 2009; Flórez Pregonero *et al.*, 2012). Regarding the latter, one study suggested that the increased use of motorcycles is due to vehicle restrictions known as ‘pico y placa’ and congested traffic (Niño-Muñoz & Morera-Ubaque, 2018). In addition, one of the best explanations for motorbikes and cars was per capita income, car prices (Mercado Díaz, 2015), and the lack of safety in public transport systems (Kash, 2019).

In the case of Colombia, there are also historical, social, and geographical factors to consider. This includes forced displacement, which caused obstacles for families to obtain food, a longing for some foods and preparations usually consumed (Ruiz Pascua, 2015), as well as an increase in the consumption of calorie-dense foods such as flour, panela water (type of sugar), tubers, soups, and sausages (Puentes & Bejarano, 2020). In Girón, Santander, a study reported that the forcibly displaced families ate twice as much corn flour and a third additional of rice and panela (Prada Gómez *et al.*, 2008). These difficulties are exacerbated in ethnic minority populations because the geographical location of many indigenous communities coincides with areas where the armed conflict takes place (Centro Nacional de Memoria Histórica, 2018). As dietary patterns vary greatly across regions, reflecting local food traditions and availability and the influence of globalisation, future research should look at regional BMI patterns.

Economic and political factors also play a role in the nutritional status of the Colombian population. In economic terms, the number of informally employed people, that is, those who do not have social security and no employment relationship with the employer, was 57.9% between November 2022 and January 2023, being higher in men than in women (60.1% vs 54.6%) (DANE, 2023), that could have an impact on the food and nutrition sphere. On political factors, while most packaged foods available in Bogotá are eligible for front-of-package warning labels, this has not yet been done (Mora-Plazas *et al.*, 2019). Regarding the type of labelling, one study showed that respondents liked and trusted those of an octagonal shape the most (Taillie *et al.*, 2020). Evidence also suggests that there has only been a minimal reduction in caloric density and sugar in food and beverages in Colombia (Lowery *et al.*, 2020). The implementation of regulation defined in 2022 therefore needs to be accelerated (Ministerio de Salud y Protección Social, 2022).

It is also worth mentioning that the Colombia government had made continuous efforts to prevent obesity, for instance through their 10-year public health plan (2012–2021). This plan contains a food and nutrition security dimension that includes a series of actions to reduce overweight and obesity (Ministerio de Salud y Protección Social, 2012). This policy is currently being evaluated and the plan for the next period is under construction. In 2009, a law on obesity was also passed, defining this disease as a public health priority and indicating that the state through the Ministries of health, culture, education, transport, environment, agriculture, and rural development will promote policies related to food and nutritional security and physical activity that promote safe environments for the development of the same (Ministerio de Educación Nacional, 2009).

Similarly, the Colombian Institute for Family Welfare (Instituto Colombiano de Bienestar Familiar (ICBF) in Spanish) has developed food guides for pregnant and breastfeeding mothers, infants, children under 2 years of age, and the general population. These guides contain a rigorous study of energy and nutrient needs and are based on regional tastes, knowledge, and food culture. They also contain technical and community outreach material that facilitates the process (ICBF, 2018b, 2018a). Despite these advances, in Colombia there are still problems concerning the unification of criteria on nutrition among professionals and effective mechanisms for the dissemination of information. Moreover, free trade agreements still promote the entry of foodstuffs into Colombia that are high in sugars, saturated fats, and sodium (Bejarano-Roncancio *et al.*, 2015) what restricts compliance with the country’s food sovereignty policy that promotes *‘the right of peoples to nutritious and culturally appropriate, accessible, sustainable and ecologically*

produced food, and their right to decide their food and production system' (Clacso, 2006). At the same time, the problem of seed hoarding and commodification, through genetic engineering and intellectual property rights by biotechnology transnationals (Gutiérrez, 2015), has affected the food and nutritional sphere in Colombia.

Strengths and limitations of the study and ideas for future research

One important strength of this study is that the ENSIN survey uses an interdisciplinary team of nutritionists, nurses, and bacteriologists during data collection. This minimises biases and errors, particularly in the taking of anthropometric measurements of weight and height required for the calculation of BMI. Due to the technical rigor applied by the survey, there is no issue of social desirability bias (i.e., underestimation of weight and overestimation of height) that is common in BMI studies that rely on self-reported measures (Gorber et al., 2007).

While our research provides insights into the basic demographic characteristics of changes in average BMI levels in the Colombian population between 2005 and 2015, there are some limitations in the use of anthropometrics for the study of trends based on cross-sectional survey data. One such limitation is the fact that BMI is only an approximate indicator of obesity, as (age, sex, and race specific) fat level, skinfold thickness, or body shape are not controlled for, although BMI is known to be highly associated with fatness (Norgan, 1994; Tsai et al., 2014). Another limitation is selection as the proportion of a cohort who already died from either underweight or obesity is unknown (Cámara & Spijker, 2010).

Finally, the results suggest the need to continue carrying out synthetic cohort studies based on cross-sectional surveys such as the ENSIN in Colombia until there are longitudinal studies with repeated measurements that allow the weight of the same individuals to be analysed over time. Especially in the context of the COVID-19 pandemic, since during confinement an increased intake of cereals, eggs, fats, sugars, and sugar cane was observed (Pertuz-Cruz et al., 2021), suggesting a possible increase in the rates of overweight and obesity in Colombia.

Acknowledgements. The results presented in this paper are part of the ongoing Doctoral Dissertation of Paula Andrea Castro-Prieto to obtain the degree of Doctor in Demography from the Universitat Autònoma de Barcelona (UAB), Centre d'Estudis Demogràfics (CED-CERCA).

The study supported with funding from the DEMOS_2021 contract through the R&D project "Salud de las personas de edad avanzada el análisis de la comorbilidad las múltiples causas de muerte y las desigualdades de género y socioeconómicas en la salud" (COMORHEALTHSES PID2020-113934RB-I00) financed by the Spanish Ministry of Science and Innovation (PI Jeroen Spijker) and also by the European Research Council (ERC-2019-COG agreement No 864616, HEALIN).

Competing interests. The authors declare no conflicts of interest. Funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Ethical approval. The study complies with the definition in the Declaration of Helsinki regarding the development of research that involves human beings. For this research, only a secondary analysis of information from data systems was carried out; therefore, the risk of the research was minimal.

References

- Bejarano JJ, Gamboa EM, Aya DH and Parra DC (2015) Los alimentos y bebidas ultra-procesados que ingresan a Colombia por el tratado de libre comercio: ¿influirán en el peso de los colombianos? *Revista Chilena de Nutrición* 42(4), 409–413. doi: [10.4067/S0717-75182015000400014](https://doi.org/10.4067/S0717-75182015000400014)
- Bourges H, Bengoa JM and O'Donnell AM (2000) *Historia de la Nutrición en América Latina*. [óéhttps://www.slan.org.ve/libros/HistoriasdeLaNutriciónenAméricaLatina.pdf](https://www.slan.org.ve/libros/HistoriasdeLaNutriciónenAméricaLatina.pdf) (accessed 16 February 2023).
- Bozkurt B, Aguilar D, Deswal A, Dunbar SB, Francis GS, et al. (2016) Contributory risk and management of comorbidities of hypertension, obesity, diabetes mellitus, hyperlipidemia, and metabolic syndrome in chronic heart failure: a scientific statement from the American Heart Association. *Circulation* 134(23), e535–e578. doi: [10.1161/CIR.0000000000000450](https://doi.org/10.1161/CIR.0000000000000450)
- Briceño G, Durán P, Colón E, Line D, Merker A, et al. (2012) Protocolo del estudio para establecer estándares normativos de crecimiento de niños colombianos sanos. *Pediatría* 45(4), 235–242. doi: [10.1016/s0120-4912\(15\)30021-5](https://doi.org/10.1016/s0120-4912(15)30021-5)

- Broughton DE and Moley KH** (2017) Obesity and female infertility: potential mediators of obesity's impact. *Fertility and Sterility* **107**(4), 840–847. doi: [10.1016/J.FERTNSTERT.2017.01.017](https://doi.org/10.1016/J.FERTNSTERT.2017.01.017)
- Cámara AD and Spijker JJ** (2010) Super size Spain? A cross-sectional and quasi-cohort trend analysis of adult overweight and obesity in an accelerated transition country. *Journal of Biosocial Science* **42**(3), 377–393. doi: [10.1017/S0021932009990629](https://doi.org/10.1017/S0021932009990629)
- Carmona J, Correa A and Alcaraz G** (2005) Población, alimentación y estado nutricional entre los tules (kunas) del resguardo Caimán Nuevo (Turbo y Necoclí; Antioquia, Colombia), 2003–2004*. *Iatreia* **18**(3), 259–278.
- Centro Nacional de Memoria Histórica** (2018) *Regiones y conflicto armado (CNMH)*. <https://centrodememoriahistorica.gov.co/regiones-y-conflicto-armado-balance-de-la-contribucion-del-cnmh-al-esclarecimiento-historico/> (accessed 16 February 2023).
- CEPAL** (2014) Los pueblos indígenas en América Latina. Avances en el último decenio y retos pendientes para la garantía de sus derechos. In *Naciones Unidas* (Vol. 1). <https://bit.ly/2QbS1DN> (accessed 16 February 2023).
- Clacso** (2006) *Declaración de Nyéléni Foro Mundial por la Soberanía Alimentaria Nyéléni, Selingue, Malí, 23 al 27 de febrero de 2007*. <http://biblioteca.clacso.edu.ar/ar/libros/osal/osal21/Nyeleni.pdf> (accessed 16 February 2023).
- da Silva MC, Oliveira AM, Pinheiro SM, de Oliveira LP and da Cruz TR** (2015) Breastfeeding and maternal weight changes during 24 months post-partum: a cohort study. *Maternal and Child Nutrition* **11**(4). doi: [10.1111/mcn.12071](https://doi.org/10.1111/mcn.12071)
- DANE** (2023) *Gran Encuesta Integrada de Hogares. Mercado laboral según sexo*. <https://www.dane.gov.co/index.php/estadisticas-por-tema/mercado-laboral/segun-sexo> (accessed 16 February 2023).
- DANE** (2023) *Ocupación informal*. https://www.dane.gov.co/files/investigaciones/boletines/ech/ech/Nueva_medicion_informalidad.pdf (accessed 21 March 2023).
- Estado Plurinacional de Bolivia** (2017) *Manual Integral de Antropometría Serie: Documentos Técnico Normativos*. www.minsalud.gob.bo (accessed 16 February 2023).
- Farfán JC, Torres DA, Gómez MN and Tamayo MP** (2019) Condiciones de seguridad alimentaria en una comunidad indígena de Colombia. *Physis: Revista de Saúde Coletiva* **28**(4), 1–16. doi: [10.1590/S0103-73312018280405](https://doi.org/10.1590/S0103-73312018280405)
- Florez PA, Gómez LF, Parra DC, Cohen DD, Arango PC, et al.** (2012) Time spent traveling in motor vehicles and its association with overweight and abdominal obesity in Colombian adults who do not own a car. *Preventive Medicine* **54**(6), 402–404. doi: [10.1016/j.ypmed.2012.04.002](https://doi.org/10.1016/j.ypmed.2012.04.002)
- Freire G, Bonilla DC, Schwartz OS, Soler LJ and Carbonari F** (2018) Afrodescendientes en Latinoamérica. <http://hdl.handle.net/10986/30201> (accessed 16 February 2023).
- Fu Q** (2019) Sex differences in sympathetic activity in obesity and its related hypertension. *Annals of the New York Academy of Sciences* **1454**(1), 1–11. doi: [10.1111/nyas.14095](https://doi.org/10.1111/nyas.14095)
- García GJ and Martín CE** (2022) A reversal in the obesity epidemic? A quasi-cohort and gender-oriented analysis in Spain. *Demographic Research* **46**, 273–290. doi: [10.4054/DEMRES.2022.46.10](https://doi.org/10.4054/DEMRES.2022.46.10)
- García LM and Contreras A** (2022) A call for implementation of the law against processed foods during the pandemic times in Colombia. *Biomedica* **42**(2), 14–18. doi: [10.7705/biomedica.6303](https://doi.org/10.7705/biomedica.6303)
- Gorber SC, Tremblay M, Moher D and Gorber B** (2007) A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obesity Reviews* **8**(4), 307–326. doi: [10.1111/J.1467-789X.2007.00347.X](https://doi.org/10.1111/J.1467-789X.2007.00347.X)
- Großschädl F and Strongegger WJ** (2021) Regional and social disparities for obesity among Austrian adults: representative long-term trends from 1973–2014 Regionale und soziale Ungleichheiten zu Adipositas bei Erwachsenen in Österreich: repräsentative Langzeittrends data source and sample. *Gesundheitswesen* **83**(1), 59–65. doi: [10.1055/a-0965-6840](https://doi.org/10.1055/a-0965-6840)
- Gutiérrez L** (2015) Soberanía alimentaria. *La Red de Semillas Libres de Colombia. Contextos* **4**(13). [https://scholar.google.es/scholar?q=Guti%C3%A9rrez+L+\(2015\)+Soberan%C3%ADa+alimentaria.+La+Red+de+Semillas+Libres+de+Colombia.+Contextos,+4\(13\).&hl=es&as_sdt=0&as_vis=1&oi=scholar](https://scholar.google.es/scholar?q=Guti%C3%A9rrez+L+(2015)+Soberan%C3%ADa+alimentaria.+La+Red+de+Semillas+Libres+de+Colombia.+Contextos,+4(13).&hl=es&as_sdt=0&as_vis=1&oi=scholar)
- Hales CM, Carroll MD, Fryar CD and Ogden CL** (2017) *Prevalence of Obesity and Severe Obesity among Adults: United States, 2017–2018 Key findings Data from the National Health and Nutrition Examination Survey*. <https://www.cdc.gov/nchs/products/index.htm> (accessed 16 February 2023).
- Hernández A, Velasco C, Oviedo M.P., Mantilla B and Flórez N** (2014) Prácticas de alimentación de la población indígena del departamento de Chocó. *Revista Ustasalud* **13**(2), 106–111.
- Herrán OF, Gamboa DE and Zea MDP** (2021) Energy and protein intake in the Colombian population: results of the 2015 ENSIN population survey. *Journal of Nutritional Science* **10**(e11), 1–10. doi: [10.1017/JNS.2021.2](https://doi.org/10.1017/JNS.2021.2)
- Herrán OF, Patiño GA and Del Castillo SE** (2016) La transición alimentaria y el exceso de peso en adultos evaluados con base en la encuesta de la situación nutricional en Colombia, 2010. *Biomedica* **36**(1), 109–120. doi: [10.7705/biomedica.v36i1.2579](https://doi.org/10.7705/biomedica.v36i1.2579)
- INEGI, Instituto Nacional de Salud Pública, & Secretaría de Salud** (2018) *Encuesta Nacional de Salud y Nutrición 2018 Presentación de resultados*. https://ensanut.insp.mx/encuestas/ensanut2018/doctos/informes/ensanut_2018_presentacion_resultados.pdf (accessed 16 February 2023).
- Instituto Colombiano de Bienestar Familiar** (2006) *Encuesta Nacional de la Situación Nutricional*. https://www.icbf.gov.co/sites/default/files/libro_2005.pdf (accessed 16 February 2023).
- Instituto Colombiano de Bienestar Familiar** (2018a) *Guías Alimentarias Basadas en Alimentos para la Población Colombiana Mayor de 2 Años | Portal ICBF – Instituto Colombiano de Bienestar Familiar ICBF*. <https://www.icbf.gov.co/guias-alimentarias-basadas-en-alimentos-para-la-poblacion-colombiana-mayor-de-2-anos-0> (accessed 16 February 2023).

- Instituto Colombiano de Bienestar Familiar** (2018b) *Guías Alimentarias basadas en Alimentos para mujeres gestantes, madres en periodo de lactancia y niños y niñas menores de 2 años para Colombia* | Portal ICBF – Instituto Colombiano de Bienestar Familiar ICBF. <https://www.icbf.gov.co/guias-alimentarias-basadas-en-alimentos-para-mujeres-gestantes-madres-en-periodo-de-lactancia-y-2> (accessed 16 February 2023).
- Instituto Colombiano de Bienestar Familiar, Ministerio de Salud y Protección Social, Instituto Nacional de Salud and Departamento de Prosperidad Social** (2015) *Encuesta Nacional de la Situación Nutricional – ENSIN 2015. Documento metodológico*. <https://www.minsalud.gov.co/sites/rid/Lists/BibliotecaDigital/RIDE/VS/ED/GCFI/documento-metodologico-ensin-2015.pdf> (accessed 16 February 2023).
- Instituto Nacional de Estadística y Geografía, Instituto Nacional de Salud Pública, Secretaría de Salud** (2023) *Encuesta Nacional de Salud y Nutrición 2018 Presentación de resultados*. https://ensanut.insp.mx/encuestas/ensanut2018/doctos/informes/ensanut_2018_presentacion_resultados.pdf (accessed 16 February 2023).
- Iyengar NM, Gucalp A, Dannenberg AJ and Hudis CA** (2016) Obesity and cancer mechanisms: tumor microenvironment and inflammation. *Journal of Clinical Oncology* **34**(35), 4270–4276. doi: [10.1200/JCO.2016.67.4283](https://doi.org/10.1200/JCO.2016.67.4283)
- Jaacks LM, Vandevijvere S, Pan A, McGowan CJ, Wallace C, et al.** (2019) The obesity transition: stages of the global epidemic. *The Lancet Diabetes & Endocrinology* **7**(3), 231–240. doi: [10.1016/S2213-8587\(19\)30026-9](https://doi.org/10.1016/S2213-8587(19)30026-9)
- Jiwani SS, Carrillo LR, Hernández VA, Barrientos GT, Basto AA, et al.** (2019) The shift of obesity burden by socioeconomic status between 1998 and 2017 in Latin America and the Caribbean: a cross-sectional series study. *The Lancet Global Health* **7**(12), e1644–e1654. doi: [10.1016/S2214-109X\(19\)30421-8](https://doi.org/10.1016/S2214-109X(19)30421-8)
- Kash G** (2019) Always on the defensive: the effects of transit sexual assault on travel behavior and experience in Colombia and Bolivia. *Journal of Transport and Health* **13**, 234–246. doi: [10.1016/j.jth.2019.04.004](https://doi.org/10.1016/j.jth.2019.04.004)
- Kasper NM, Herrán OF and Villamor E** (2013) Obesity prevalence in Colombian adults is increasing fastest in lower socioeconomic status groups and urban residents: results from two nationally representative surveys. *Public Health Nutrition* **17**(11), 2398–2406. doi: [10.1017/S1368980013003418](https://doi.org/10.1017/S1368980013003418)
- Lear SA, Teo K, Gasevic D, Zhang X, Poirier PP, et al.** (2014). The association between ownership of common household devices and obesity and diabetes in high, middle and low income countries. *Canadian Medical Association Journal* **186**(4), 258–266. doi: [10.1503/cmaj.131090](https://doi.org/10.1503/cmaj.131090)
- Lowery CM, Mora-Plazas M, Gómez LF, Popkin B and Taillie LS** (2020) Reformulation of packaged foods and beverages in the Colombian food supply. *Nutrients* **12**(11), 3260. doi: [10.3390/nu12113260](https://doi.org/10.3390/nu12113260)
- McTigue KM, Garrett JM and Popkin B** (2002) The natural history of the development of obesity in a cohort of young U.S. adults between 1981 and 1998. *Annals of Internal Medicine* **136**(12), 857–864. doi: [10.7326/0003-4819-136-12-200206180-00006](https://doi.org/10.7326/0003-4819-136-12-200206180-00006)
- Mercado DM** (2015) Modelos alternativos para la estimación de la tasa de motorización en Colombia. <https://manglar.uninorte.edu.co/handle/10584/9213> (accessed 21th March 2023).
- Ministerio de Educación Nacional** (2009) *Ley 1355 de octubre 14 de 2009*. <https://www.mineduacion.gov.co/portal/normativa/Leyes/381525:Ley-1355-de-octubre-14-de-2009> (accessed 16 February 2023).
- Ministerio de Salud y Protección Social** (2012) *Plan Decenal de Salud Pública*. <https://www.minsalud.gov.co/PlanDecenal/Paginas/home2013.aspx> (accessed 16 February 2023).
- Ministerio de Salud y Protección Social** (2016) *Resolución 2465 de 2016*. https://www.icbf.gov.co/sites/default/files/resolucion_no_2465_del_14_de_junio_de_2016.pdf (accessed 16 February 2023).
- Ministerio de Salud y Protección Social** (2022) *MinSalud expide Resolución 2492_2022 sobre etiquetado nutricional y frontal*. <https://www.minsalud.gov.co/Paginas/MinSalud-expide-Resolucion-2492-2022-sobre-etiquetado-nutricional-y-frontal.aspx> (accessed 16 February 2023).
- Ministerio de Salud y Protección Social de Colombia, Instituto Colombiano de Bienestar Familiar, Instituto Nacional de Salud** (2019) *Encuesta Nacional de la Situación Nutricional-ENSIN, 2015. Documento metodológico*. <https://www.minsalud.gov.co/sites/rid/Lists/BibliotecaDigital/RIDE/VS/ED/GCFI/documento-metodologico-ensin-2015.pdf> (accessed 16 February 2023).
- Molano TN and Molano TD** (2018) Cosmovisión de salud y alimentación en la cultura Guambiana. *Worldview of Health and Food in the Guambiana Culture* **20**(1), 16–25. doi: [10.22267/rus.182001.105](https://doi.org/10.22267/rus.182001.105)
- Mora-Plazas M, Gómez LF, Miles DR, Parra DC and Taillie LS** (2019) Nutrition quality of packaged foods in Bogotá, Colombia: a comparison of two nutrient profile models. *Nutrients* **11**(5), 1011. doi: [10.3390/nu11051011](https://doi.org/10.3390/nu11051011)
- Niño MD and Morera UN** (2018) Percepción de la pobreza en Colombia en los años 2003 y 2016. *Clío América* **12**(23), 25–38. doi: [10.21676/23897848.2614](https://doi.org/10.21676/23897848.2614)
- Norgan NG** (1994) Population differences in body composition in relation to the body mass index. *European Journal of Clinical Nutrition* **48**(3), S10–S25.
- Nuttall FQ** (2015) Body mass index: obesity, BMI, and health: a critical review. *Nutrition Today* **50**(3), 117–128. doi: [10.1097/NT.0000000000000092](https://doi.org/10.1097/NT.0000000000000092)
- Parra DC, Lobelo F, Gómez LF, Rutt C, Schmid T, et al.** (2009) Household motor vehicle use and weight status among Colombian adults: are we driving our way towards obesity? *Preventive Medicine* **49**(2–3), 179–183. doi: [10.1016/j.ypmed.2009.07.010](https://doi.org/10.1016/j.ypmed.2009.07.010)

- Pensiero N and Green A** (2018) The effects of post-compulsory education and training systems on literacy and numeracy skills: a comparative analysis using PISA 2000 and the 2011 survey of adult skills. *European Journal of Education* 53(2), 238–253. doi: [10.1111/EJED.12268](https://doi.org/10.1111/EJED.12268)
- Pertuz CS, Molina ME, Rodríguez PC, Guerra HE, Cobos de Rangel OP, et al.** (2021) Exploring dietary behavior changes due to the COVID-19 confinement in Colombia: a national and regional survey study. *Frontiers in Nutrition* 8(644800), 1–16. doi: [10.3389/fnut.2021.644800](https://doi.org/10.3389/fnut.2021.644800)
- Popkin B, Corvalan C and Grummer SL** (2020) Dynamics of the double burden of malnutrition and the changing nutrition reality. *The Lancet* 395(10217), 65–74. doi: [10.1016/S0140-6736\(19\)32497-3](https://doi.org/10.1016/S0140-6736(19)32497-3)
- Prada GG, Herrán FO and Cárdenas RO** (2008) Patrón alimentario y acceso a los alimentos en familias desplazadas en el municipio de Girón, Santander, Colombia. *Revista Panamericana de Salud Publica/Pan American Journal of Public Health* 23(4), 257–263. doi: [10.1590/S1020-49892008000400005](https://doi.org/10.1590/S1020-49892008000400005)
- Preston SH and Wang H** (2006) Sex mortality differences in the United States: the role of cohort smoking patterns. *Demography* 43(4), 631–646. doi: [10.1353/dem.2006.0037](https://doi.org/10.1353/dem.2006.0037)
- Profamilia, Instituto Nacional de Salud, Instituto Colombiano de Bienestar Familiar and Ministerio de la Protección Social** (2011) *Encuesta Nacional de la Situación Nutricional en Colombia 2010*. <https://www.icbf.gov.co/sites/default/files/resumenfi.pdf> (accessed 16 February 2023).
- Puentes M and Bejarano A** (2020) Prácticas de consumo alimentario de familias desplazadas por el conflicto armado, asentadas en Bosa, Bogotá – Dialnet. *Diversitas: Perspectivas En Psicología* 16(1), 143–155.
- Qian CJ, Coulombe J, Suissa S and Ernst P** (2017) Pneumonia risk in asthma patients using inhaled corticosteroids: a quasi-cohort study. *British Journal of Clinical Pharmacology* 83(9), 2077–2086. doi: [10.1111/bcp.13295](https://doi.org/10.1111/bcp.13295)
- Rivas X, Pazos S, Castillo S and Pachón H** (2010) Alimentos autóctonos de las comunidades indígenas y afrodescendientes de Colombia. *Archivos Latinoamericanos de Nutrición* 60(3). <https://www.alanrevista.org/ediciones/2010/3/art-1/> (accessed 5 June 2023).
- Rueda YP** (2019) Aproximación a los determinantes de la doble carga nutricional en hogares. <https://repositorio.uniandes.edu.co/handle/1992/34121> (accessed 16 February 2023).
- Ruiz PM** (2015) Alimentando la vida frente al desplazamiento forzado: memoria y cocina como propuestas de paz* Nurturing life to face forced displacement: memories and kitchen as peace proposals. *Eleuthera* 12, 112–130. doi: [10.17151/eleu.2015.12.6](https://doi.org/10.17151/eleu.2015.12.6)
- Sansón RA, Bernal RJ, Kubow S, Suarez MA and Melgar QH** (2021) Food insecurity and the double burden of malnutrition in Colombian rural households. *Public Health Nutrition* 24(14), 4417–4429. doi: [10.1017/S1368980021002895](https://doi.org/10.1017/S1368980021002895)
- Shrimpton R and Rokx C** (2012) *The Double Burden of Malnutrition: A Review of Global Evidence*. <https://openknowledge.worldbank.org/handle/10986/27417> (accessed 16 February 2023).
- Suissa S** (2015) The quasi-cohort approach in pharmacoepidemiology: upgrading the nested case-control. *Epidemiology* 26(2), 242–246. doi: [10.1097/EDE.0000000000000221](https://doi.org/10.1097/EDE.0000000000000221)
- Taillie LS, Hall MG, Gómez LF, Higgins I, Bercholz M, et al.** (2020) Designing an effective front-of-package warning label for food and drinks high in added sugar, sodium, or saturated fat in Colombia: an online experiment. *Nutrients* 12(10), 3124. doi: [10.3390/nu12103124](https://doi.org/10.3390/nu12103124)
- Thomas RK** (2018). *Concepts, Methods and Practical Applications in Applied Demography*. <https://link.springer.com/book/10.1007/978-3-319-65439-3> (accessed 16 February 2023).
- Tod E, Bromley C, Millard AD, Boyd A, Mackie P and McCartney G** (2017) Obesity in Scotland: a persistent inequality. *International Journal for Equity in Health* 16(1), 1–13. doi: [10.1186/s12939-017-0599-6](https://doi.org/10.1186/s12939-017-0599-6)
- Tremmel M, Gerdtham UG, Nilsson PM and Saha S** (2017) Economic burden of obesity: a systematic literature review. *International Journal of Environmental Research and Public Health* 14(4), 1–18. doi: [10.3390/ijerph14040435](https://doi.org/10.3390/ijerph14040435)
- Tsai EW, Perng W, Mora-Plazas M, Marín C, Baylin A, et al.** (2014) Accuracy of self-reported weight and height in women from Bogotá, Colombia. *Annals of Human Biology* 41(5), 473–476. doi: [10.3109/03014460.2013.856939](https://doi.org/10.3109/03014460.2013.856939)
- Vecino O and Arroyo A** (2018) A tax on sugar sweetened beverages in Colombia: estimating the impact on overweight and obesity prevalence across socio economic levels. *Social Science and Medicine* 209(40), 111–116. doi: [10.1016/j.socscimed.2018.05.043](https://doi.org/10.1016/j.socscimed.2018.05.043)
- Victora CG, Bahl R, Barros AJ, França GV, Horton S, et al.** (2016) Breastfeeding in the 21st century: epidemiology, mechanisms, and lifelong effect. *The Lancet* 387 (10017), 475–490. doi: [10.1016/S0140-6736\(15\)01024-7](https://doi.org/10.1016/S0140-6736(15)01024-7)
- Waite S** (2015) Does it get better? A quasi-cohort analysis of sexual minority wage gaps. *Social Science Research* 54, 113–130. doi: [10.1016/j.ssresearch.2015.06.024](https://doi.org/10.1016/j.ssresearch.2015.06.024)
- World Health Organization** (2021) *Obesity and Overweight*. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight> (accessed 16 February 2023).
- Zore T, Palafox M and Reue K** (2018) Sex differences in obesity, lipid metabolism, and inflammation—a role for the sex chromosomes? *Molecular Metabolism* 15, 35–44. doi: [10.1016/j.molmet.2018.04.003](https://doi.org/10.1016/j.molmet.2018.04.003)

Appendix A




Figure A1. Nutritional status by sex, age group, and survey year of Afro-Colombians. Source: ENSIN 2010–2015. Own calculations.




Figure A2. Nutritional status by sex, age group, and survey year of indigenous people. Source: ENSIN 2010–2015. Own calculations.




Figure A3. Nutritional status by sex, age group, and survey year of others ethnicities. Source: ENSIN 2010–2015. Own calculations.

Table A1. Missing values

Survey year	Total cases	Age group	Missing values in weight		Missing values in height		Weight less than 30 kg		Height less than 130 cm		Total missing	
			Men	Women	Men	Women	Men	Women	Men	Women		
2005	80006		Total population									
		20–24	2645	1040	1	0	1	3	5	2	3697	
		25–29	2302	866	4	5	0	4	0	6	3187	
		30–34	2031	694	6	3	0	4	2	5	2745	
		35–39	1873	682	4	9	0	3	2	3	2576	
		40–44	1670	666	6	6	0	0	1	0	2349	
		45–49	1368	649	7	7	0	1	1	2	2035	
		50–54	1168	515	4	4	0	1	0	4	1696	
		55–59	788	403	0	7	0	1	0	2	1201	
		60–64	783	556	1	6	0	0	1	2	1349	
	Total	14628	6071	33	47	1	17	12	26	20835		
2010	106125		Afro-Colombians									
		20–24	178	122	0	0	0	0	0	1	301	
		25–29	192	92	0	0	0	0	0	0	284	
		30–34	140	64	0	0	0	0	1	0	205	
		35–39	136	61	0	0	0	0	0	1	198	
		40–44	155	59	1	0	0	0	0	1	216	
		45–49	122	45	0	1	1	0	0	1	170	
		50–54	80	47	0	0	0	0	0	0	127	
		55–59	74	45	0	0	0	0	0	1	120	
		60–64	51	29	0	0	0	0	0	0	80	
			Total	1128	564	1	1	1	0	1	5	1701
				Indigenous								
		20–24	176	96	0	0	0	0	0	0	0	272
		25–29	187	78	0	0	0	0	1	0	0	266
		30–34	159	47	0	0	0	1	1	1	0	209
		35–39	95	50	0	0	0	0	1	0	0	146
		40–44	108	40	0	1	0	0	0	0	0	149
		45–49	106	46	0	0	0	0	0	0	0	152
		50–54	81	44	0	0	0	0	0	1	0	126
55–59	38	33	0	0	0	0	0	0	0	71		
60–64	40	31	0	0	0	0	0	0	0	71		
	Total	990	465	0	1	0	1	3	2	1462		

(Continued)

Table A1. (Continued)

Survey year	Total cases	Age group	Missing values in weight		Missing values in height		Weight less than 30 kg		Height less than 130 cm		Total missing
			Men	Women	Men	Women	Men	Women	Men	Women	
Others											
		20–24	1842	1054	3	0	1	0	1	0	2901
		25–29	1862	879	1	1	1	1	2	0	2747
		30–34	1645	755	0	1	0	2	0	3	2406
		35–39	1456	636	2	4	0	0	3	1	2102
		40–44	1450	608	2	2	0	1	0	2	2065
		45–49	1326	582	0	1	0	1	2	1	1913
		50–54	1067	572	2	2	0	0	0	1	1644
		55–59	826	462	1	2	0	0	1	2	1294
		60–64	572	389	1	2	0	0	0	3	967
		Total	12046	5937	12	15	2	5	9	13	18039
2015	81485	Afro-Colombians									
		20–24	100	33	1	0	0	0	0	1	135
		25–29	77	37	1	0	0	0	0	1	116
		30–34	73	25	1	0	0	0	0	1	100
		35–39	63	34	1	0	0	0	0	0	98
		40–44	54	31	0	0	0	0	0	0	85
		45–49	47	24	1	2	0	0	0	0	74
		50–54	41	25	0	0	0	0	0	0	66
		55–59	44	21	0	0	0	0	1	0	66
		60–64	21	21	0	0	0	0	0	0	42
		Total	520	251	5	2	0	0	1	3	782
		Indigenous									
		20–24	79	26	1	0	0	0	0	0	106
		25–29	61	22	0	1	0	0	0	0	84
		30–34	50	24	0	0	0	0	0	0	74
		35–39	57	17	0	0	0	0	1	0	75
		40–44	47	16	0	0	0	0	0	0	63
		45–49	37	21	0	0	0	0	0	1	59
		50–54	22	20	0	0	0	0	0	0	42
		55–59	27	13	0	0	0	0	0	0	40
		60–64	15	8	0	1	0	0	0	0	24
		Total	395	167	1	2	0	0	1	1	567

(Continued)

Table A1. (Continued)

Survey year	Total cases	Age group	Missing values in weight		Missing values in height		Weight less than 30 kg		Height less than 130 cm		Total missing
			Men	Women	Men	Women	Men	Women	Men	Women	
Others											
		20–24	916	434	12	1	0	2	3	4	1372
		25–29	943	404	6	3	0	0	1	2	1359
		30–34	792	319	2	1	1	3	2	5	1125
		35–39	735	285	0	2	1	0	2	2	1027
		40–44	687	276	1	2	0	0	1	0	967
		45–49	591	311	13	3	0	0	6	2	926
		50–54	604	289	2	5	0	0	1	2	903
		55–59	499	282	3	2	0	0	0	2	788
		60–64	337	227	2	2	0	1	1	3	573
		Total	6104	2827	41	21	2	6	17	22	9040
Total	267616										52426

Table A2. Cross-sectional analysis of average weight by sex and age

Sex and survey year	Age group								
	20–24	25–29	30–34	35–39	40–44	45–49	50–54	55–59	60–64
Men 2005	65.18–65.90	68.06–68.90	70.13–71.06	70.74–71.70	70.93–71.90	70.79–71.87	69.86–70.99	69.39–70.68	67.80–69.26
Men 2010	66.34–66.95	70.20–70.93	72.22–73.00	72.45–73.24	73.04–73.81	72.39–73.21	71.32–72.24	70.51–71.53	68.94–70.00
Men 2015	67.53–68.24	71.99–72.82	73.98–74.87	75.40–76.30	74.61–75.50	74.32–75.27	73.53–74.45	72.68–73.69	71.06–72.15
Women 2005	56.04–56.58	58.75–59.37	61.12–61.81	62.67–63.37	63.43–64.19	64.78–65.55	64.79–65.68	64.11–65.16	61.91–63.12
Women 2010	57.86–58.39	60.92–61.50	63.09–63.72	64.15–64.80	65.23–65.88	65.83–66.51	65.96–66.70	65.38–66.24	63.85–64.84
Women 2015	60.38–61.03	63.33–64.04	65.58–66.32	66.64–67.40	67.38–68.19	67.55–68.33	67.02–67.82	66.52–67.39	65.50–66.42

Source: ENSIN 2005–2015. Own calculations.

Table A3. Nutritional status (%) by sex and, survey year

Survey year	Nutritional status	Men		Women		Total	
		n	%	n	%	n	%
2005	Underweight	642	2.85	1219	3.43	1861	3.20
	Normal weight	11945	53.07	15771	44.33	27716	47.72
	Overweight	7579	33.68	12082	33.96	19661	33.85
	Obese	2340	10.40	6502	18.28	8842	15.22
	Total	22506	100.00	35574	100.00	58080	100.00
2010	Underweight	688	1.87	1242	2.58	1930	2.27
	Normal weight	18009	49.05	19035	39.49	37044	43.62
	Overweight	13271	36.15	17309	35.91	30580	36.01
	Obese	4747	12.93	10621	22.03	15368	18.10
	Total	36715	100.00	48207	100.00	82922	100.00
2015	Underweight	612	1.94	773	1.95	1385	1.95
	Normal weight	13334	42.35	13561	34.20	26895	37.81
	Overweight	12572	39.93	15108	38.10	27680	38.91
	Obese	4966	15.77	10208	25.75	15174	21.33
	Total	31484	100.00	39650	100.00	71134	100.00
Total		90705	42.36	123431	57.64	214136	100.00

Source: ENSIN 2005–2015. Own calculations.

Table A4. Nutritional status (%) by sex, ethnicity, and survey year

Survey year	Nutritional status	Afro-Colombians						Indigenous						Others					
		Men		Women		Total		Men		Women		Total		Men		Women		Total	
		n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
2010	Underweight	96	2.53	169	3.50	265	3.07	38	1.00	77	1.80	115	1.42	554	1.90	996	2.55	1550	2.27
	Normal weight	2027	53.36	1803	37.29	3830	44.36	2153	56.45	1840	43.07	3993	49.38	13829	47.52	15392	39.37	29221	42.84
	Overweight	1201	31.61	1660	34.33	2861	33.14	1308	34.29	1579	36.96	2887	35.70	10762	36.98	14070	35.98	24832	36.41
	Obese	475	12.50	1203	24.88	1678	19.43	315	8.26	776	18.16	1091	13.49	3957	13.60	8642	22.10	12599	18.47
	Total	3799	100	4835	100	8634	100	3814	100	4272	100	8086	100	29102	100	39100	100	68202	100
2015	Underweight	49	1.68	69	1.89	118	1.79	28	1.16	34	1.11	62	1.13	535	2.05	670	2.03	1205	2.04
	Normal weight	1294	44.28	1105	30.20	2399	36.45	1093	45.09	1091	35.60	2184	39.79	10947	41.88	11365	34.52	22312	37.78
	Overweight	1058	36.21	1295	35.39	2353	35.75	997	41.13	1213	39.58	2210	40.26	10517	40.24	12600	38.27	23117	39.14
	Obese	521	17.83	1190	35.52	1711	26.00	306	12.62	727	23.72	1033	18.82	4139	15.84	8291	25.18	12430	21.04
	Total	2922	100	3659	100	6581	100	2424	100	3065	100	5489	100	26138	100	32926	100	59064	100
Total		6721	44.17	8494	55.83	15215	100	6238	45.95	7337	54.05	13575	100	55240	43.41	72026	56.59	127266	100

Source: ENSIN 2010–2015. Own calculations.

Cite this article: Castro-Prieto PA, Spijker J, and Recaño J (2024). A quasi-cohort trend analysis of adult obesity in Colombia. *Journal of Biosocial Science* 56, 63–89. <https://doi.org/10.1017/S002193202300010X>