January 1967

Vol. LXXXI No. 1

The Journal of MML Laryngology and Otology

EDITED BY G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

Contents

LESIONS OF THE APICAL REGION OF THE COCHLEA JOEL M. BERNSTEIN and HAROLD F. SCHUKNECHT TOMOGRAPHY OF THE INNER EAR IN DEAF CHILDREN. RADIOLOGICAL DEMONSTRATION OF TWO JØRGEN JENSEN CASES WITH THE MONDINI MALFORMATION . T. R. BULL, NASO-ALVEOLAR CYSTS K. A. MCNEILL, G. MILNER and S. M. MURRAY DACRYOCYSTORHINOSTOMY IN CHILDREN AND INFANTS M. SPENCER HARRISON and AIIT KUMAR MUKHERJEE HEARING CHILDREN OF DEAF PARENTS EDMUND CRITCHLEY THE ACUITY OF HEARING IN THE KALAHARI BUSHMEN. J. F. JARVIS A PILOT SURVEY 5610 and H. G. VAN HEERDEN POSTERIOR TYMPANOTOMY. ITS IMPORTANCE IN COM-G. D. L. SMYTH, BINED APPROACH TYMPANOPLASTY R. M. ENGLAND, R. GIBSON and A. G. KERR G. D. L. SMYTH, THE PROBLEMS OF TYPE C TYMPANOPLASTY R. M. ENGLAND, R. GIBSON and A. G. KERR A PUNCH CARD SYSTEM FOR PROCESSING DATA FROM HAROLD LUDMAN STAPEDECTOMY OPERATIONS CLINICAL RECORDS-K. G. MALCOMSON OSSIFYING FIBROMA OF THE SPHENOID K. KISHORE CHONDROMA OF NASO-PHARYNX LATERAL EXTENSIONS OF NASOPHARYNGEAL FIB-ROMA . M. L. BHATIA, S. C. MISHRA and I. PRAKASH FIBROSARCOMA OF LARYNX - ITS CLINICAL FEATURES AND TREATMENT S. D. SINGH

SOCIETIES' PROCEEDINGS

GENERAL NOTES

London

Headley Brothers

109 Kingsway WC2

Annual Subscription £5/5/0 net, U.S.A. \$15

Montbly, 12/6 net post free

Published online by Cambridge University Press

ADVERTISEMENTS

The Journal of Laryngology and Otology

(FOUNDED in 1887 by MORELL MACKENZIE and NORRIS WOLFENDEN)

EDITED BY

G. H. BATEMAN

ASSISTANT EDITOR LIONEL TAYLOR

1. Original articles which have not been published elsewhere are invited and should be sent to the Editor. They are considered for publication on the under-standing that they are contributed to this *Journal* solely. Reproduction elsewhere, in whole or in part, is not permitted without the previous written consent of the Author and Editor and the customary acknowledgment must be made.

2. Manuscripts should be typewritten, on one side only of the paper, and well spaced. Captions to illustrations should be typed on a separate sheet and sent at the same time as original photographs, etc. The Harvard system of recording references should be used, e.g. GREEN, C., and BROWN, D. (1951) J. Laryng., 65, 33. Abbreviations of Journals should follow the style recommended in World Medical Periodicals, published by World Health Organization, 1952.

It is most important that authors should verify personally the accuracy of every reference before submitting a paper for publication.

3. Galley proofs and engraver's proofs of illustrations are sent to the author, Corrections, which should be kept to a minimum, must be clearly marked, and no extra matter added. Proofs should be returned within 5 days.

4. Illustration blocks will normally be held by the Printers for three years, after which they will be destroyed. Any author is entitled to have these returned to him, but a request for this must be sent within three years of the appearance of the article, to HEADLEY BROTHERS, 109 Kingsway, London, WC2. Coloured illustra-tions will be charged to authors, unless a special grant is authorized by the Editor.

5. Orders for reprints must be sent when returning galley proofs, and for this purpose special forms are supplied.

6. Authors of original communications on Oto-Laryngology in other journals are invited to send a copy, or two reprints, to the *Journal of Laryngology*. If they are willing, at the same time, to submit their own abstract (in English, French, Italian, or German) it will be welcomed.

7. Editorial communications may be addressed to THE EDITOR, Journal of Laryngology, c/o HEADLEY BROTHERS, 109 Kingsway, London, WC2.

8. The annual subscription is five guineas sterling (U.S.A. \$15) post free, and payable in advance.

9. Single copies will be on sale at 12s. 6d. each; copies of parts up to Vol. LXIII which are available may be purchased at 7s. 6d. each.

10. All subscriptions, advertising and business communications should be sent to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2.

United States of America

Orders for this Journal may be sent through local booksellers, or to STECHERT-HAFNER, INC., 31-33 East 10th Street, New York, or direct to the publishers, HEADLEY BROTHERS, 109 KINGSWAY, LONDON, WC2, England.


C Journal of Laryngology and Otology, 1966

CONTENTS

	PAGE
LESIONS OF THE APICAL REGION OF THE COCHLEA. Joel M. Bernstein and Harold F. Schuknecht (Boston, U.S.A.)	I
TOMOGRAPHY OF THE INNER EAR IN DEAF CHILDREN. RADIOLOGICAL DEMONSTRATION OF TWO CASES WITH THE MONDINI MALFOR- MATION. Jørgen Jensen (Copenhagen)	27
NASO-ALVEOLAR CYSTS. T. R. Bull, K. A. McNeill, G. Milner and S. M. Murray (Kingston, Jamaica)	37
DACRYOCYSTORHINOSTOMY IN CHILDREN AND INFANTS. M. Spencer Harrison (London) and Ajit Kumar Mukherjee (Lincoln) .	45
HEARING CHILDREN OF DEAF PARENTS. Edmund Critchley (London) .	51
THE ACUITY OF HEARING IN THE KALAHARI BUSHMEN. A PILOT SURVEY. J. F. Jarvis and H. G. van Heerden (Cape Town)	63
POSTERIOR TYMPANOTOMY. ITS IMPORTANCE IN COMBINED APPROACH TYMPANOPLASTY. G. D. L. Smyth, R. M. England, R. Gibson and A. G. Kerr (Belfast)	69
THE PROBLEMS OF TYPE C TYMPANOPLASTY. G. D. L. Smyth, R. M. England, R. Gibson and A. G. Kerr (Belfast)	75
A PUNCH CARD SYSTEM FOR PROCESSING DATA FROM STAPEDECTOMY OPERATIONS. Harold Ludman (London)	79
CLINICAL RECORDS—	
Ossifying Fibroma of the Sphenoid. K. G. Malcomson (Bristol) ,	87
CHONDROMA OF NASO-PHARYNX. K. Kishore (Liverpool)	93
LATERAL EXTENSIONS OF NASOPHARYNGEAL FIBROMA. M. L. Bhatia, S. C. Mishra and J. Prakash (Lucknow)	99
FIBROSARCOMA OF LARYNX—ITS CLINICAL FEATURES AND TREATMENT.	
S. D. Singh (Wolverhampton)	107
Societies' Proceedings	111
General Notes	115

14

ii



-that is the triumph of the **MACKIE** hearing aid

The new Mackie post-aural hearing aid is made in several different circuit versions so that the gain and frequency response can be selected to match the patient's needs; and it is available where required with a unique form of automatic volume control which has proved highly successful in cases of recruitment. The forward pick-up microphone gives maximum intelligibility of speech without background interference.

It is exported to many countries including the United States of America–a tribute to its high quality and advanced design.

In Great Britain it is available for trial by your patients from *independent* hearing aid dealers in all parts of the country.

MANUFACTURED BY A. G. R. MACKIE LTD. A member of the Bonochord Group of Companies FOR FURTHER DETAILS WRITE TO 48 WELBECK STREET, LONDON WI



'Otosporin' rapidly destroys virtually all bacteria commonly found in ear infections, reduces irritation, inflammation and pain. 'Otosporin' brand Drops contain polymyxin B sulphate, neomycin sulphate and hydrocortisone in a bland suspension designed to facilitate penetration. Issued in bottles of 5 ml.





Burroughs Wellcome & Co. London (The Wellcome Foundation Ltd.)

BÉKÉSY Accessory

For the very first time a fully automatic unit has been made available for a Clinic Audiometer, making the combined units into a true BÉKÉSY Audiometer.

This automatic accessory is being made for the PETERS SPD/5 CLINIC AUDIOMETER, which has a continuous frequency range and continuously variable attenuation. The result is therefore a true Békésy audiometer.

It can be attached to or detached from the audiometer in under one minute so that the two units together offer the facilities of an advanced clinic audiometer or Békésy audiometry at will.

It can be fitted to any existing SPD/5 Audiometer.

Please write for full details and demonstration to :

ALFRED PETERS & SONS LIMITED 51 GELL STREET, SHEFFIELD 3

ONLY **AMPLIVOX** provides a <u>complete</u> audiometric service!



Transistor Audiometers

with air conduction or air and bone conduction with masking, for hospitals, schools, consulting 'rooms, industrial medical departments, screening programmes, etc.



v

Clinical Audiometers

with narrow band masking for the highest accuracy and extended resolution.

Audiometer Booths

for ideal testing conditions in hospitals, clinics, factories. Standard and special sizes available.



PLUS An efficient and speedy calibration service please write for technical literature to meet your needs 80 New Bond Street, London, W.1. Tel: HYDe Park 9888

ORBENIN and its success in Gram-Positive Infections

Otitis media Otitis externa

Boils Carbuncles Infected dermatoses

Pneumonia Lung abscess Acute bronchitis

Osteomyelitis Osteitis

Tonsillitis Pharyngitis

Post-operative wound infections Burns Skin graft protection

Septicaemia Acute endocarditis

Staphylococcal enterocolitis Staphylococcal urinary tract infections Staphylococcal meningitis

The second seco

tis encouraging that the widespread the of Orbenin in hospitals has not expited in the emergence of resistant

Hore than 300 references in the World Interature testify to the excellent clinical esuits obtained with Orbenin in a wide reviety of Gram-positive infections.

Detailed literature is available on request.



Two capsules (500mg.) or one vial (250mg.) q.i.d.

Availability:



Vials (250mg.)





Orbenin (cloxacillin sodium B.P.)(regd.) is a product of British research at **Beecham Research Laboratories** Brentford, England. Telephone: ISLeworth 4111

WHEN A HEARING AID IS RECOMMENDED ONLY AMPLIVOX OFFERS THIS HEARING AID SERVICE

The design and manufacture of a complete range with individual fitting facilities. Best of other makes in addition to Amplivox aids—Amplivox hearing aids are ONLY available from Amplivox. Highest degree of training of any hearing aid organisation. Master Hearing Aid evaluations. Expert follow-up and rehabilitation advice. 15 branch offices with while-you-wait service. FOR YOUR PATIENTS' GREATEST SATISFACTION SPECIFY AMPLIVOX **CANCELED SCIEVE AMPLIVOX**

AMPLIVOX HEARING ADVISORY SERVICE 80 NEW BOND STREET LONDON, W.1

THE LARYNGOSCOPE

A Monthly Journal devoted to the disease of EAR, NOSE AND THROAT

Official organ for the American Laryngological Rhinological and Otological Society

Price \$18.00 per year Canada \$19.00 per year Foreign \$19.00 per year

Established 1896

Theodore E. Walsh, M.D. Editor

517 SOUTH EUCLID AVENUE SAINT LOUIS MO. 63110.



Please mention The Journal of Laryngology and Otology when replying to advertisements

all clear in sinusitis, otitis media and coryza


'Triogesic' provides decongestion with analgesia.

	TABLETS	ELIXIR
	scored (per tablet)	cherry flavoured (per 5 ml.)
Revised	Phenylpropanolamine-HCl 12.5 mg.	Phenylpropanolamine-HCI 3 mg.
Formula	Paracetamol B.P. 500 mg.	Paracetamol B.P. 125 mg.
		Alcohol (95%) 0.5 ml.
Packs	tubes of 30	bottles of 8 and 20 fl. oz.
Basic N.H.S. Price	3/6d. for 30 tablets	5/10d. for 8 fl. oz.

Literature and sample on request. A. Wander Ltd., 42 Upper Grosvenor Street, London W.1









MULTITONE

Many firms sell hearing aids, only MULTITONE thought of making an electronic ear trumpet; quite apart from having the most comprehensive range of standard and special hearing instruments.

Several important new instruments will be added to the range during 1967. We hope they will help with some of your special problems.

MULTITONE ELECTRIC COMPANY LIMITED UNDERWOOD STREET, LONDON, N.I

Telephone (01) CLE 8022



the antistaphylococcal antibiotic that penetrates tissues at these levels ...

bears the FUCIDIN label



The clinical significance of these levels has been confirmed in many publications during the past five years. Over 100 authors have endorsed the value of Fucidin in the treatment of staphylococcal Hayes Middlesex

infection. Fucidin (Sodium Fusidate B.P.) 250 mgm capsules.

Leo Laboratories Limited






MIDDLE EAR PROSTHESES



Sultana's Stainless Steel Spring Middle Ear Prosthesis, 4 mm. long

Vide Journal of Laryngology Vol. LXXIX, No. 9 Smyth's "Clothes-Pin" Incus replacement Prosthesis



Shepard's Teflon Grommet Drain Tubes

Further details of these and other Middle Ear Prostheses on request

DOWN BROS. AND MAYER & PHELPS LTD CHURCH PATH, MITCHAM, SURREY, ENGLAND

> In Canada 410 Dundas Street West, Toronto, 2B, Canada

KAMPLEX AUDIOMETRIC EQUIPMENT

AUDIOMETERS

SCREENING PORTABLE TA16 DIAGNOSTIC PORTABLE TA15

CLINIC AUDIOMETER

AUTOMATIC AUDIOMETER

DOUBLE CHANNEL



CLINIC AUDIOMETER DA1

ACOUSTIC IMPEDANCE APPARATUS ZA2

PNEUMOMETER ZP1

SKIN RESISTANCE APPARATUS RA4

SOUND LEVEL METER GA6

VISIBLE SPEECH APPARATUS

AUDITORY TRAINERS

Write for full catalogue and detailed descriptions to:


ALFRED PETERS & SONS LTD 51 GELL STREET, SHEFFIELD 3

Please mention The Journal of Laryngology and Otology when replying to advertisements

DA1

DA2

AB1



What makes this antibiotic so valuable in the ENT unit ?

Spectrum?

Exceptionally broad. It covers most gram-positive organisms—including the penicillin-resistant staphylococci—and many gram-negative organisms.

Action?

Intensely bactericidal. It gives a rapid kill of bacteria at or near the minimum inhibitory concentration, especially in the case of gram-negative organisms.

Safety?

Wide clinical usage has confirmed the laboratory evidence that Ceporin has a remarkably low toxicity. It is usually well tolerated by patients allergic to penicillin.

Clinical record?

The latest analysis of clinical results obtained with Ceporin shows a total of 131 ENT infections treated : 109 (83%) responded satisfactorily. There were 28 cases of tonsillitis, of which 24 responded well. The majority of these infections was caused by beta-haemolytic streptococci, but there were several mixed infections involving streptococci, staphylococci and *Corynebacterium diphtheriae*.

Otitis also responded well. Of the 28 cases treated, 24 responded satisfactorily. Most of these infections were caused by staphylococci but infections involving coliforms, pneumococci, Streptococcus pyogenes, proteus and Haemophilus influenzae were also represented. Often rapid clinical response was obtained with Ceporin after other antibiotics had failed. Conditions which respond well to Ceporin include : mastoiditis, otitis media, sinusitis, quinsy, pharyngeal abscess. and intracranial infections secondary to infected sinuses and mastoiditis. broad spectrum, bactericidal action and low toxicity—Ceporin has proved valuable for use in place of more toxic drugs or combinations of antibiotics. Presentation Vials containing 250 mg, 500 mg and 1 gram of cephaloridine, packed singly and in boxes of five.

Ceporin is a Glaxo trade mark

Detailed literature is available from Glaxo Laboratories Limited Greenford, Middlesex