

Shocks and UV radiation around low-mass protostars: the Herschel-PACS legacy

Agata Karska^{1,2,3}, Michael J. Kaufman⁴, Lars E. Kristensen⁵,
Ewine F. van Dishoeck^{3,2} and WISH, DIGIT, and WILL teams

¹Centre for Astronomy, Faculty of Physics, Astronomy and Informatics, Nicolaus Copernicus University, Grudziadzka 5, 87-100 Torun, Poland
email: agata.karska@umk.pl

²Max-Planck Institut für Extraterrestrische Physik (MPE),
Giessenbachstr. 1, D-85748 Garching, Germany

³Leiden Observatory, Leiden University,
P.O. Box 9513, 2300 RA Leiden, The Netherlands

⁴Department of Physics and Astronomy, San Jose State University,
One Washington Square, San Jose, CA 95192-0106, USA

⁵Centre for Star and Planet Formation, Niels Bohr Institute, University of Copenhagen,
Øster Voldgade 5-7, DK-1350 Copenhagen K, Denmark

Abstract. Far-infrared spectroscopy reveals gas cooling and its underlying heating due to physical processes taking place in the surroundings of protostars. These processes are reflected in both the chemistry and excitation of abundant molecular species. Here, we present the *Herschel*-PACS far-IR spectroscopy of 90 embedded low-mass protostars from the WISH (van Dishoeck *et al.* 2011), DIGIT (Green *et al.* 2013), and WILL surveys (Mottram *et al.* 2017). The 5×5 spectra covering the $\sim 50'' \times 50''$ field-of-view include rotational transitions of CO, H₂O, and OH lines, as well as fine-structure [O I] and [C II] in the ~ 50 -200 μm range. The CO rotational temperatures are typically ~ 300 K, with some sources showing additional components with temperatures as high as ~ 1000 K. The H₂O / CO and H₂O / OH flux ratios are low compared to stationary shock models, suggesting that UV photons may dissociate some H₂O and decrease its abundance. Comparison to C shock models illuminated by UV photons show a good agreement between the line emission and the models for pre-shock densities of 10^5 cm^{-3} and UV fields 0.1-10 times the interstellar value. The far-infrared molecular and atomic lines are the unique diagnostic of shocks and UV fields in deeply-embedded sources.

Keywords. Young stellar objects, outflows, jets, shocks

1. Introduction

During the early formation of low-mass protostars, the mass accretion rates are high and thus, the feedback from the protostar on its surrounding is most spectacular. The launching of jets and winds creates outflow cavities and generates shock waves propagating in the envelope. Shocks compress and heat the gas but may also produce UV photons, which penetrate to large distances due to low densities and scattering in the outflow cavities. Characterising these processes requires spectroscopy in the far-infrared (IR), where cooling of the warm, dense gas is very efficient (Kaufman & Neufeld 1996). The question remains, how do these feedback processes influence the initial conditions (both physics and chemistry) of star and planet formation.

In this paper, we briefly summarize the survey of 90 deeply-embedded low-mass protostars observed as part of the three large *Herschel* programs: ‘Water in Star forming regions with Herschel’ (WISH, van Dishoeck *et al.* 2011), ‘Dust, Ice, and Gas in Time’




Figure 1. *Left:* CO rotational diagram of a Class 0 protostar NGC 1333 IRAS 4A. *Right:* Line ratio of H₂O 4₀₄ – 3₁₃ at 125 μm and OH at 84 μm as a function of logarithm of density of the pre-shock gas. The range observed in ~ 90 protostars is shown as filled, light-blue box, and the models of C shocks are shown as solid lines (models of shielded shocks, Kaufman & Neufeld 1996) and violet stars (models of UV illuminated shocks, Melnick & Kaufman 2015).

(DIGIT, Green *et al.* 2013), and ‘William Herschel Line Legacy’ (WILL, Mottram *et al.* 2017). All observations were obtained using the Photodetector Array Camera and Spectrometer (PACS, Poglitsch *et al.* 2010) that provided spectral maps consisting of 25 spatial pixels of $\sim 9.4'' \times 9.4''$, corresponding to pixel sizes of $\sim 2000 \times 2000$ AU for typical distances of 200 pc.

2. Results

Rich molecular line emission is commonly detected in Class 0/I protostars including highly-excited H₂O ($E_{\text{up}} \sim 1000$ K) and CO lines ($E_{\text{up}} \sim 5000$ K). The line fluxes of molecular species correlate strongly with each other and show similar extent of emission, typically following the outflow direction. Multiple transitions of CO lines are used to determine rotational temperatures of the gas (see left panel of Figure 1). A ~ 300 K component is universally seen in almost all protostars, whereas the hotter gas component show a broad distribution with a median at 720 K (detected in 27% of sources). These two components are identified in the spectrally-resolved CO 16-15 lines obtained with HIFI (Kristensen *et al.* 2017). The ~ 300 K component in the PACS CO ladder corresponds to the broad, Gaussian component in the line profiles originating in outflow cavity shocks or disk winds. The hotter component is associated with the offset component likely produced in the irradiated shocks (see Kristensen *et al.* 2017, for discussion).

The right panel of Figure 1 shows a comparison of the measured H₂O / OH line ratios and the predictions from stationary (1D plane-parallel) C shock models from Kaufman & Neufeld (1996). Clearly, these models overestimate the observed line ratios by 1-2 orders of magnitude. Low H₂O / OH line ratios require partial photodissociation of H₂O into OH and O. Predictions from C shock models illuminated by UV photons (Melnick & Kaufman 2015) reproduce the observed ranges of H₂O / OH for pre-shock densities of 10^5 cm^{-3} and UV fields 0.1-10 times the interstellar value. These preliminary comparisons show that UV radiation influences the chemistry in deeply-embedded protostars, as also evidenced by observations of light ionised hydrides (Kristensen *et al.* 2013, Benz *et al.* 2016).




Figure 2. *Left:* Fractions of gas cooling contributed by CO (orange), H₂O (blue), OH (green), and [O I] (red), to the total far-infrared line cooling in Class 0 protostar NGC 1333 IRAS 4B. *Right:* Median line cooling in CO, H₂O, [O I], and OH over bolometric luminosities for Class 0 and Class I sources.

Low abundances of H₂O are also confirmed by relatively low total line cooling in H₂O versus other molecules in far-IR. Kaufman & Neufeld (1996) predict that H₂O should account for 70-90% of total molecular line cooling in the far-IR (H₂O+CO+OH, Karska *et al.* 2014). At the same time, the observed H₂O emission is typically below 30% and at most about 50% in the most H₂O-rich source NGC1333 IRAS4B (left panel of Figure 2). Clearly, some H₂O is missing and the UV photodissociation is a possible mechanism to decrease its abundance.

Far-IR molecular line cooling decreases from Class 0 to Class I (right panel of Figure 2), but the differences are significant only in CO and not in H₂O (Karska *et al.* 2018). Proper interpretation of these trends requires the analysis of velocity-resolved profiles, which reveal multiple kinematic components with varying contributions depending on the transition (e.g. Kristensen *et al.* 2017).

The far-IR data presented here demonstrate the necessity for excellent space-based facilities to quantify the chemistry and excitation of warm molecular and atomic gas. The next major step will occur when the *James Webb Space Telescope* (JWST) is launched in 2018; this telescope will provide crucial data on not only the fmid-IR cooling species, but most importantly on the dominant gas coolant: molecular hydrogen, H₂. Direct measurements of the atomic and molecular abundances will be performed and tested against sophisticated shock models.

Acknowledgement. The authors are grateful to G. Herczeg, J. Mottram, L. Tychoniec, J. Lindberg, N. Evans II, J. Green, Y.-L. Yang, A. Gusdorf, and N. Siodmiak for collaborations on this project. This work is supported by the Polish National Science Center grant 2013/11/N/ST9/00400 and 2016/21/D/ST9/01098.

References

- Benz, A. O., Bruderer, S., van Dishoeck, E. F., *et al.* 2016, *A&A*, 590, A105
 Green, J. D., Evans, II, N. J., Jrgensen, J. K., *et al.* 2013, *ApJ*, 770, 123
 Karska, A., Kristensen, L. E., van Dishoeck *et al.* 2014, *A&A*, 572, A9
 Karska, A., Kaufman, M., Kristensen, L.E., *et al.* 2018, *ApJS*, 235, 30
 Kaufman, M. J. & Neufeld, D. A. 1996, *ApJ*, 456, 611
 Kristensen, L. E., van Dishoeck, E. F., Benz, A. O. *et al.* 2013, *A&A*, 557, A23
 Kristensen, L. E., van Dishoeck, E. F., Mottram, J. C. *et al.* 2017, arXiv No. 1705.10269
 Melnick, G. J. & Kaufman, M. J. 2015, *ApJ*, 806, 227
 Mottram, J. C., van Dishoeck, E. F., Kristensen, L. E., *et al.* 2017, *A&A*, 600, A99
 Poglitsch, A., Waelkens, C., Geis, N., *et al.* 2017, *A&A*, 518, L2
 van Dishoeck, E. F., Kristensen, L. E., Benz, A. O., *et al.* 2011, *PASP*, 123, 138