

# Plumboperloffite, $\text{PbMn}^{2+}_2\text{Fe}^{3+}_2(\text{PO}_4)_3(\text{OH})_3$ , a new member of the bjarebyite group from Wiperaminga Hill, South Australia, Australia

Peter Elliott<sup>1,2\*</sup> and Anthony R. Kampf<sup>3</sup>

<sup>1</sup>Department of Earth Sciences, School of Physical Sciences, The University of Adelaide, Adelaide, South Australia 5005, Australia.

<sup>2</sup>South Australian Museum, North Terrace, Adelaide, South Australia 5000, Australia

<sup>3</sup>Mineral Sciences Department, Natural History Museum of Los Angeles County, 900 Exposition Boulevard, Los Angeles, CA 90007, U.S.A.


\*E-mail: [peter.elliott@adelaide.edu.au](mailto:peter.elliott@adelaide.edu.au)

## ABSTRACT

Plumboperloffite,  $\text{PbMn}^{2+}_2\text{Fe}^{3+}_2(\text{PO}_4)_3(\text{OH})_3$ , is a new mineral and member of the bjarebyite group from Wiperaminga Hill West Quarry, Boolcoomatta Reserve, Olary Province, South Australia, Australia. The mineral was found in a single cavity in triplite-barbosalite matrix associated with fluorapatite, phosphosiderite, natrodufrénite and fluorite. The mineral forms intergrowths of subparallel, thin tabular to bladed crystals. Individual crystals are up to 40  $\mu\text{m}$  in length. Plumboperloffite is brownish orange in colour with a vitreous luster. The mineral has brittle tenacity, an excellent cleavage on  $\{100\}$  and uneven fracture. The calculated density is 4.416  $\text{g}/\text{cm}^3$ . Plumboperloffite is biaxial (+),  $\alpha = 1.87(1)$ ,  $\beta = 1.88(1)$ ,  $\gamma = 1.89(1)$  as measured in white light. The measured 2V is  $88(1)^\circ$ . Dispersion is apparently strong, based on extinction colours and the orientation is  $Y = \mathbf{b}$ . The pleochroism in shades of yellow brown is  $X < Z < Y$ . Electron microprobe analysis gave the empirical formula (based on 15 O *apfu*)

$(\text{Pb}_{0.92}\text{Ca}_{0.04}\text{Ba}_{0.01}\text{K}_{0.01})_{0.98}(\text{Mn}^{2+}_{1.84}\text{Fe}^{2+}_{0.13})_{1.97}(\text{Fe}^{3+}_{1.97}\text{Al}_{0.03})_{2.00}(\text{P}_{3.01}\text{O}_{11.94})(\text{OH})_{3.06}$ .

Plumboperloffite is monoclinic, space group  $P2_1/m$  with  $a = 9.1765(18)$ ,  $b = 12.340(3)$ ,  $c =$



This is a 'preproof' accepted article for Mineralogical Magazine. This version may be subject to change during the production process.  
DOI: 10.1180/mgm.2023.97

5.0092(10) Å,  $\beta = 101.01(3)^\circ$ ,  $V = 556.8(2) \text{ \AA}^3$  and  $Z = 2$ . The crystal structure has been refined using X-ray single-crystal data to a final  $R1 = 0.0207$  on the basis of 1417 reflections with  $F_o > 4\sigma(F_o)$ . The mineral is isostructural with members of the bjarebyite group minerals.

**Keywords:** plumboperloffite, new mineral species, bjarebyite group, crystal structure, pegmatitic phosphate, Wiperaminga Hill West Quarry, Australia

## INTRODUCTION

Bjarebyite-group minerals are secondary phosphates that occur primarily in granitic pegmatites but have also been found in the oxidized zones of ore deposits. Bjarebyite was first described from the Palermo No.1 pegmatite, New Hampshire, USA by Moore *et al.* (1973) and its atomic structure was determined by Moore and Araki (1974). The bjarebyite-group currently consists of seven International Mineralogical Association (IMA) approved minerals (Table 1). The minerals have the general formula  $XM_1M_2(PO_4)_3(OH)_3$  where  $X = \text{Ba, Sr, Pb}$ ;  $M_1 = \text{Fe}^{2+}, \text{Mn}^{2+}, \text{Mg}$  and  $M_2 = \text{Al, Fe}^{3+}$ . Until the discovery of strontioferroperloffite (Elliott, 2019), all group members contained Ba as the dominant cation at the  $X$  site. The new mineral species, plumboperloffite, characterized and described in this paper, is the first member of the group with Pb dominant at the  $X$  site. The name is for the relationship to perloffite with Pb dominating the  $X$  site. The mineral and mineral name have been approved by the International Mineralogical Association, Commission on New Minerals, Nomenclature and Classification (IMA 2020–007). The type specimen is deposited in the South Australian Museum, Adelaide, South Australia, (Registration number G34868).

## OCCURRENCE

Plumboperloffite was found on a specimen collected in 2018 from the dumps of the Wiperaminga Hill West Quarry, Boolcoomatta Reserve, Olary Province, South Australia, Australia (140°27'34"E, 31°57'42"S). Pegmatites at Wiperaminga Hill were exploited for feldspar, beryl and muscovite between 1957 and 1980 (Olliver and Steveson, 1984). Four pegmatites were worked in three workings on Wiperaminga Hill West and Wiperaminga Hill East. Recorded production is 31.7 tonnes of beryl and 2030 tonnes of feldspar. The

pegmatites occur near the base of a prominent range of hills which comprise banded quartz-mica gneiss and schist of the Early Middle Proterozoic age Willyama Complex. The metasediments have been granitized and in many places are intruded by granite and associated pegmatite and aplite bodies. The pegmatites are usually small lenticular bodies varying from 6 m to 30 m in length and from 1 m to 3 m in width (Hiern, 1966). The pegmatites are mineralogically zoned and comprise an outer border zone of fine- to medium-grained microcline, quartz, plagioclase and muscovite, an intermediate zone of coarse-grained muscovite, quartz, microcline, plagioclase, beryl and apatite (and triplite), and an inner quartz core or cores (Lottermoser and Lu, 1997). The Wiperaminga Hill pegmatites belong to the beryl-columbite phosphate-rare element type in the classification of Černý (1991). Phosphate minerals, dufrénite and floreneite, were first reported from Wiperaminga Hill by King (1954), and Hiern (1966) reported apatite, triplite, dickinsonite and barbosalite. During a visit to the locality in 2018 by one of the authors (PE), masses of orange to dark brown triplite-zwieselite were found on the dumps of the eastern quarry. Secondary phosphate minerals have formed in seams and cavities in the triplite-zwieselite by hydrothermal alteration and weathering, in an oxidizing, low-temperature, low-pH environment (Lottermoser and Lu, 1997). Plumboperloffite occurs in a single cavity in a matrix comprising triplite, barbosalite and minor quartz. Associated minerals are fluorapatite, phosphosiderite, natrodufrénite and fluorite.

## PHYSICAL AND OPTICAL PROPERTIES

Plumboperloffite occurs as thin tabular to bladed crystals, with a brownish-orange colour. Individual crystals are up to 40  $\mu\text{m}$  in length (Fig.1). The streak is pale orange and the mineral has a vitreous lustre. The hardness is estimated to be 4 based on the hardness of other bjarebyite-group minerals. The fracture is uneven, the tenacity is brittle and the mineral has an excellent cleavage on  $\{100\}$ . Density was not determined owing to the small amount of material available. The calculated density is 4.416  $\text{g/cm}^3$  from the empirical formula. Plumboperloffite is optically biaxial (+) with indices of refraction  $\alpha = 1.87(1)$ ,  $\beta = 1.88(1)$ ,  $\gamma = 1.89(1)$  measured in white light. The  $2V$  measured directly by conoscopic observation is  $88(1)^\circ$ . The calculated  $2V$  is  $90.5^\circ$ . Dispersion is apparently strong, based on extinction colours, but the sense could not be determined because of the indistinct conoscopic figure. The pleochroism in shades of yellow brown is  $X < Z < Y$ . The optical

orientation is  $Y = \mathbf{b}$ . The optical orientation could not be completely determined because crystals are subparallel intergrowths and the refractive indices are high. The compatibility index  $1 - (K_P/K_C)$  is 0.03 (excellent) based upon the empirical formula, density calculated using the single-crystal cell and the measured indices of refraction (Mandarino, 1981).

## CHEMICAL COMPOSITION

Analyses of plumboperloffite (12 spots) were performed with a Cameca SXFive electron microprobe (wavelength-dispersive spectroscopy mode, 20 kV, 20 nA and 5  $\mu\text{m}$  beam diameter). Data were processed using the  $\phi(\rho Z)$  correction procedure of Pouchou and Pichoir (1985). There was insufficient material for direct determination of  $\text{H}_2\text{O}$  content so it was based upon the crystal structure analysis. The results of the analyses are reported in Table 2. The empirical formula unit, based on 15 O atoms per formula unit (apfu) is  $(\text{Pb}_{0.92}\text{Ca}_{0.04}\text{Ba}_{0.01}\text{K}_{0.01})_{0.98}(\text{Mn}^{2+}_{1.84}\text{Fe}^{2+}_{0.13})_{1.97}(\text{Fe}^{3+}_{1.97}\text{Al}_{0.03})_{2.00}(\text{P}_{3.01}\text{O}_{11.94})(\text{OH})_{3.06}$ . The ideal formula is  $\text{PbMn}^{2+}_2\text{Fe}^{3+}_2(\text{PO}_4)_3(\text{OH})_3$  which requires PbO 29.19, MnO 18.55,  $\text{Fe}_2\text{O}_3$  20.87,  $\text{P}_2\text{O}_5$  27.86,  $\text{H}_2\text{O}$  3.53, total 100.00 wt%.

## INFRARED SPECTROSCOPY

The infrared spectrum of plumboperloffite (Fig. 2) was recorded using a Nicolet 5700 FTIR spectrometer equipped with a Nicolet Continuum IR microscope and a diamond-anvil cell. Data were acquired from a powdered sample during 50 scans in the wavenumber range 4000 to 700  $\text{cm}^{-1}$ . The spectrum shows bands at 3471  $\text{cm}^{-1}$  and 3209  $\text{cm}^{-1}$  that correspond to O–H-stretching vibrations of OH groups forming weak hydrogen bonds (Libowitzky and Beran, 2004). The bands at 1157, 1067, 989 and 898  $\text{cm}^{-1}$  are assigned to  $\nu_3$  vibrations of the  $\text{PO}_4$  tetrahedra and the band at 827  $\text{cm}^{-1}$  is assigned to the  $\nu_1$  vibration of the  $\text{PO}_4$  tetrahedra. No absorption attributed to H–O–H bending vibrations was observed in the spectrum.

## CRYSTALLOGRAPHY

### *Powder X-ray diffraction*

Powder X-ray diffraction data for plumboperloffite were obtained using a Rigaku R-Axis Rapid II curved-imaging-plate microdiffractometer, with monochromatised  $\text{MoK}\alpha$

radiation (50 kV, 40 mA). A Gandolfi-like motion on the  $\varphi$  and  $\omega$  axes was used to randomize the sample. Observed  $d$  values and intensities were derived by profile fitting using JADE 2010 software (Materials Data, Inc.). Data (in Å for MoK $\alpha$ ) are given in Table 3. Unit-cell parameters refined from the powder data using JADE 2010 with whole-pattern fitting are  $a = 9.17(2)$ ,  $b = 12.41(2)$ ,  $c = 5.02(2)$  Å,  $\beta = 100.84(5)^\circ$  and  $V = 561(3)$  Å<sup>3</sup>.

#### *Single-crystal X-ray diffraction*

The single-crystal X-ray diffraction experiment was carried out on the micro-focus macromolecular MX2 beamline at the Australian Synchrotron, part of ANSTO (Aragao *et al.*, 2018). A crystal was attached to a MiTeGen polymer loop and X-ray diffraction data was collected at 100 K using a Dectris EigerX 16M detector and monochromatic radiation with a wavelength of 0.710754 Å. The intensity data sets were processed using XDS (Kabsch, 2010) without scaling and with absorption correction and scaling performed using SADABS (Bruker, 2001). Crystal data and experimental details are reported in Table 4.

#### *Structure refinement*

The structure was solved in space group  $P2_1/m$  using SHELXT (Sheldrick, 2015a) as implemented in the WinGX suite (Farrugia, 1999). SHELXL-2018 (Sheldrick, 2015b) was used for the refinement of the structure. Following the solution of the structure, the atom coordinates were transformed to correspond to those in the structure of perloffite (Elliott and Willis, 2011). Difference Fourier syntheses then located all H atom positions. The H sites were refined with soft restraints on the O–H distances of 0.85(3) Å for the O–H groups and 0.90(3) Å. The final cycle of the refinement converged to  $R_1 = 0.0207$ . Final atom coordinates and anisotropic-displacement parameters are listed in Table 5, selected interatomic distances and framework angles are given in Table 6, and bond-valence values, calculated using the parameters of Gagné and Hawthorne (2015) are given in Table 7.

#### *Structure description*

Plumboperloffite<sub>2</sub> is isostructural with minerals of the bjarebyite group. The structure contains one  $X$  site (occupied by Pb and minor Ca, Ba and K), one  $M1$  site (occupied by Mn<sup>2+</sup> and minor Fe<sup>2+</sup>), one  $M2$  site, (occupied by Fe<sup>3+</sup> and minor Al) and two P sites. The  $M1$  and  $M2$  sites were refined with full occupancy of Mn and Fe respectively.

$[M2O_3(OH)_3]$  octahedra share edges to form  $[M_2O_6(OH)_4]$  dimers which corner-link to form  $(M2_2\phi_9)$  chains along the **b**-direction (Fig. 3). Octahedra that share corners are also linked by  $P1O_4$  tetrahedra. The chains are decorated by  $P2O_4$  tetrahedra, which link between chains in the **c**-direction via the O5 and O6 anions. Chains are linked in the **a**-direction via  $[M1O_5(OH)]$  octahedra, which share an edge with  $[M2^{3+}O_3(OH)_3]$  octahedra and corners with  $P1O_4$  and  $P2O_4$  tetrahedra (Fig. 4). The *X* site occupies channels parallel to  $[001]$ , also provides linkage between chains in the **a**-direction. The *X* site exhibits a lopsided coordination (Fig. 5) with three very short bonds to O1 and O7, four medium length bonds to O2, O4 and OH8 and six very long bonds to O5, O6 and O7. This coordination is distinct from the more regular coordination of the *X* site in the structures of other bjarebyite-group minerals and is attributable to the accommodation of the stereoactive  $6s^2$  electron lone pair on  $Pb^{2+}$ . Coordination of the *M1* and *M2* sites is similar to the sites in the structures of other bjarebyite-group minerals. The  $M1O_6$  octahedron is strongly distorted with bond angles in the range  $86.35$  to  $109.53^\circ$ . and The  $M1O_6$  octahedron is relatively undistorted with bond angles in the range  $86.77$  to  $96.22^\circ$ .

The hydrogen bonding scheme for plumboperloffite given in Table 6, is the same as reported in previous studies on bjarebyite-group minerals (e.g. Cooper and Hawthorne, 1994; Elliott, 2019). The H9 atom associated with OH9 hydrogen bonds to O3 with a donor-acceptor distance of  $H9 \cdots O3 = 1.89(3)$  Å and an  $OH9-H2 \cdots O3$  angle of  $168(8)^\circ$ , an arrangement typical of strong hydrogen bonding (Libowitzky and Beran, 2004). The H8 atom provides a weak hydrogen bond to O6 with a donor-acceptor distance of  $H8 \cdots O6 = 2.73(2)$  Å and an  $OH8-H8 \cdots O6$  angle of  $124(6)^\circ$ .

## ACKNOWLEDGMENTS

The authors thank Ben Wade of Adelaide Microscopy, The University of Adelaide for assistance with the microprobe analysis. The infrared spectrum was acquired with the assistance of the Forensic Science Centre, Adelaide. This research was undertaken in part using the MX2 beamline at the Australian Synchrotron, part of ANSTO, and made use of the Australian Cancer Research Foundation (ACRF) detector.

## REFERENCES

- Aragao, D., Aishima, J., Cherukuvada, H., Clarken, R., Clift, M., Cowieson, N.P., Ericsson, D.J., Gee, C.L., Macedo, S., Mudie, N., Panjekar, S., Price, J.R., Riboldi-Tunnicliffe, A., Rostan, R., Williamson, R. and Caradoc-Davies, T.T. (2018) MX2: a high-flux undulator microfocus beamline serving both the chemical and macromolecular crystallography communities at the Australian Synchrotron. *Journal of Synchrotron Radiation*, **25**, 885–891.
- Bowman, M.G., Downs, R.T, and Yang, H. (2013) Penikisite,  $\text{BaMg}_2\text{Al}_2(\text{PO}_4)_3(\text{OH})_3$ , isostructural with bjarebyite, *Acta Crystallographica*, **E69**, i4-i5.
- Bruker (2001) SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Černý, P. (1991) Rare element granitic pegmatites, part I. Anatomy and internal evolution of pegmatite deposits. *Geoscience Canada*, **18**, 49–67.
- Cooper, M. and Hawthorne, F.C. (1994) Refinement of the crystal structure of kulanite. *The Canadian Mineralogist*, **32**, 15–19.
- Elliott, P. and Willis, A.C. (2011) The crystal structure of perloffite. *Mineralogical Magazine*, **75**, 317–325.
- Elliott, P. (2019) Strontio-perloffite,  $\text{SrMn}^{2+}_2\text{Fe}^{3+}_2(\text{PO}_4)_3(\text{OH})_3$ , a new bjarebyite-group mineral from the Spring Creek mine, South Australia, Australia. *European Journal of Mineralogy*, **31**, 549–555.
- Farrugia, L.J. (2012) WinGX and ORTEP for Windows: an update. *Journal of Applied Crystallography*, **45**, 849–854.
- Gagné, O.C. and Hawthorne, F.C. (2015) Comprehensive derivation of bond-valence parameters for ion pairs involving oxygen. *Acta Crystallographica*, **B71**, 562–578.
- Hiern, M.N. (1966) Feldspar and beryl deposits - Wiperaminga Hill. South Australia. Department of Mines and Energy, *Mining Review*, 124, 44–51.
- Kabsch, W. (2010) XDS. *Acta Crystallographica*, **D66**, 125–132.
- Kampf, A.R. (1977) A new mineral: perloffite, the  $\text{Fe}^{3+}$  analogue of bjarebyite. *Mineralogical Record*, **8**, 112–114.
- King, D. (1954) Examination of Rare-Earth Pegmatite near Wiperaminga Hill, Old Boolcoomata Station. Department of Mines and Energy, *Mining Review* **97**, 93–94.
- Kolitsch, U., Pring, A. and Tiekink, E.R.T. (2000) Johntomaite, a new member of the bjarebyite group of barium phosphates: description and structure refinement. *Mineralogy and Petrology*, **70**, 1–14.

- Libowitzky, E. and Beran, A. (2004) IR spectroscopic characterisation of hydrous species in minerals. Pp. 227–279 in: *Spectroscopic Methods in Mineralogy*. (A. Beran and E. Libowitzky, editors). EMU Notes in Mineralogy, **6**. Eötvös University Press, Budapest.
- Lottermoser, B.G. and Lu, J. (1997) Petrogenesis of rare-element pegmatites in the Olary Block, South Australia, part 1. Mineralogy and chemical evolution. *Mineralogy and Petrology*, **59**, 1–19.
- Mandarino, J.A. (1981) The Gladstone-Dale relationship: Part IV: The compatibility concept and its application. *The Canadian Mineralogist*, **19**, 441–450.
- Moore, P.B., Lund, D.H., and Keester, K.L. (1973) Bjarebyite,  $(\text{Ba,Sr})(\text{Mn,Fe,Mg})_2\text{Al}_2(\text{OH})_3(\text{PO}_4)_3$ , a new species. *Mineralogical Record*, **4**, 282–285.
- Moore, P.B. and Araki, T. (1974) Bjarebyite,  $\text{Ba}(\text{Mn,Fe})^{2+}_2\text{Al}_2(\text{OH})_3[\text{PO}_4]_3$ : Its atomic arrangement. *American Mineralogist*, **59**, 567–572.
- Olliver, J.G. and Steveson, B.G. (1982) Pegmatites in the Olary Province. A review of feldspar and beryl mining north of Olary and the results of reconnaissance sampling of feldspar. *South Australian Department of Mines and Energy*, report **81/74**.
- Pouchou, J.-L., and Pichoir, F. (1991) Quantitative analysis of homogeneous or stratified microvolumes applying the model “PAP”. In K.F.J. Heinrich and D.E. Newbury, Eds., *Electron Probe Quantitation*, p. 31–75. Plenum Press, New York.
- Sheldrick, G.M. (2015a) SHELXT - Integrated space-group and crystal-structure determination. *Acta Crystallographica*, **A71**, 3–8.
- Sheldrick, G.M. (2015b) Crystal structure refinement with SHELXL. *Acta Crystallographica*, **C71**, 3–8.

## FIGURE CAPTIONS





Figure 1. Orange crystals of plumboperloffite associated with natrodufrénite (black) and fluorite (white) The scale bare is 200  $\mu\text{m}$ .

Prepublished




Figure 2. The FT-IR spectrum of powdered plumboperloffite.

Prepublished




Figure 3. The crystal structure of plumboperloffite projected on (100). Hydrogen atoms are grey spheres. Hydrogen bonds are indicated by dotted lines.




Figure 4. The crystal structure of plumboperloffite projected onto (001). The X positions are shown as purple spheres. The unit cell is outlined.




Figure 5. The Pb coordination in plumboperloffite showing Pb-O bond lengths in angstroms, Å.

Table 1. Comparison of related minerals.

Mineral name	plumboperloffite	strontioferroperloffite	bjarebyite	johntomaite	kulanite	perloffite	penikisite
Formula	$\text{PbMn}^{2+}_2\text{Fe}^{3+}_2(\text{PO}_4)_3(\text{OH})_3$	$\text{SrMn}^{2+}_2\text{Fe}^{3+}_2(\text{PO}_4)_3(\text{OH})_3$	$\text{BaMn}^{2+}_2\text{Al}_2(\text{PO}_4)_3(\text{OH})_3$	$\text{BaFe}^{2+}_2\text{Fe}^{3+}_2(\text{PO}_4)_3(\text{OH})_3$	$\text{BaFe}^{2+}_2\text{Al}_2(\text{PO}_4)_3(\text{OH})_3$	$\text{BaMn}^{2+}_2\text{Fe}_2(\text{PO}_4)_3(\text{OH})_3$	$\text{BaMg}_2\text{Al}_2(\text{PO}_4)_3(\text{OH})_3$
Crystal system	monoclinic	monoclinic	monoclinic	monoclinic	monoclinic	monoclinic	monoclinic
Space group	$P2_1/m$	$P2_1/m$	$P2_1/m$	$P2_1/m$	$P2_1/m$	$P2_1/m$	$P2_1/m$
$a$ (Å)	9.1765(18)	9.2195(18)	8.930(14)	9.199(9)	9.014(1)	9.223(5)	8.9577(4)
$b$ (Å)	12.340(3)	12.410(3)	12.073(24)	12.359(8)	12.074(1)	12.422(8)	12.0150(5)
$c$ (Å)	5.0092(10)	4.9990(10)	4.917(9)	5.004(2)	4.926(1)	4.995(2)	4.9079(2)
$\beta$ (°)	101.01(3)	100.09(3)	100.15(13)	100.19(6)	100.48(1)	100.39(4)	100.505(2)
$V$ (Å <sup>3</sup> )	556.8(2)	563.12(19)	521.8	559.9(5)	527.1(1)	562.9(5)	519.37
$Z$	2	2	2	2	2	2	2
Reference	This work	Elliott, (2019)	Moore and Araki (1974)	Kolitsch <i>et al.</i> (2000)	Cooper and Hawthorne (1994)	Kampf (1977)	Bowman <i>et al.</i> (2013)

Prepublished Article

Table 2. Analytical data for plumboperloffite.

Constituent	Wt. %	Range	Stand. Dev.	Probe Standard
PbO	26.60	25.82–27.44	0.48	galena
BaO	0.25	0.10–0.61	0.15	barite
CaO	0.28	0.08–0.86	0.25	plagioclase
K <sub>2</sub> O	0.23	0.11–0.39	0.09	sanidine
MnO	17.06	16.51–17.62	0.29	rhodonite
FeO*	1.23	0.78–2.09	0.41	almandine-pyrope
Fe <sub>2</sub> O <sub>3</sub> *	20.53	20.13–20.78	0.18	almandine-pyrope
Al <sub>2</sub> O <sub>3</sub>	0.18	0.01–0.50	0.14	almandine-pyrope
P <sub>2</sub> O <sub>5</sub>	27.98	27.79–28.58	0.23	apatite
H <sub>2</sub> O (calc)**	3.60			
Total	97.94			

\*FeO and Fe<sub>2</sub>O<sub>3</sub> calculated to give Fe<sup>3+</sup>+Al = 2.00\*\*H<sub>2</sub>O calculated from the refined formula.Table 3. Powder X-ray data (*d* in Å) for plumboperloffite. Only calculated lines with *I* ≥ 3 are listed. *I*<sub>calc</sub> values have been scaled so that the three lines contributing to the highest observed peak have a total intensity of 100.

<i>I</i> <sub>obs</sub>	<i>d</i> <sub>obs</sub>	<i>d</i> <sub>calc</sub>	<i>I</i> <sub>calc</sub>	<i>hkl</i>	<i>I</i> <sub>obs</sub>	<i>d</i> <sub>obs</sub>	<i>d</i> <sub>calc</sub>	<i>I</i> <sub>calc</sub>	<i>hkl</i>	<i>I</i> <sub>obs</sub>	<i>d</i> <sub>obs</sub>	<i>d</i> <sub>calc</sub>	<i>I</i> <sub>calc</sub>	<i>hkl</i>
25	9.029	9.0076	39	1 0 0	9	2.1673	2.1643	3	2 5 0	7	1.6255	1.6262	12	4 4 1
12	6.196	6.1700	10	0 2 0			2.1517	8	3 4 0			1.6117	6	-1 2 3
40	5.105	5.0903	38	1 2 0			2.1338	4	-1 3 2	14	1.5937	1.6068	4	-5 0 2
8	4.848	4.9170	19	0 0 1	16	2.1233	2.1258	4	1 2 2			1.5960	9	5 0 1
		4.7108	3	-1 0 1			2.1154	5	4 2 0			1.5872	6	-1 6 2
30	4.583	4.5677	32	0 1 1			2.1099	8	-3 0 2			1.5837	9	-4 4 2
10	4.349	4.4010	11	-1 1 1			2.1013	5	1 5 1			1.5775	3	0 6 2
5	4.027	4.0061	5	1 0 1			2.0836	3	-4 2 1			1.5703	3	-3 0 3
9	3.770	3.7442	11	-1 2 1	25	2.0669	2.0567	20	0 6 0	15	1.5595	1.5590	4	-5 4 1
		3.6907	7	-2 0 1			2.0516	5	-2 5 1			1.5577	7	-3 1 3
20	3.657	3.6377	25	2 2 0			2.0051	3	1 6 0			1.5553	7	2 5 2
6	3.553	3.5360	21	-2 1 1			2.0030	3	2 0 2			1.5531	3	3 6 1
18	3.367	3.3600	12	1 2 1	29	1.9919	1.9964	8	-3 2 2			1.5425	4	0 8 0
		3.1673	5	-2 2 1			1.9837	21	1 3 2			1.5269	3	-2 3 3
100	3.158	3.1549	50	0 3 1	57	1.9384	1.9772	8	2 1 2	23	1.5273	1.5255	8	2 7 1
		3.0984	44	-1 3 1			1.9493	16	-4 3 1			1.5226	6	0 3 3
18	3.064	3.0372	16	2 3 0			1.8974	3	0 6 1			1.5202	10	3 7 0
42	2.950	2.9556	36	2 1 1	14	1.8806	1.8790	4	3 4 1			1.5191	6	-6 0 1
23	2.898	2.9186	39	1 4 0			1.8721	11	-2 4 2			1.5132	5	1 2 3
		2.8699	28	1 3 1	6	1.8342	1.8551	6	-3 5 1			1.5106	3	4 1 2
		2.7470	3	-2 3 1			1.8278	3	4 2 1	9	1.4978	1.4967	3	-5 3 2
		2.7422	52	-3 1 1			1.8255	4	1 4 2			1.4937	11	-3 7 1
58	2.738	2.7300	28	2 2 1	9	1.8070	1.8015	12	5 0 0			1.4728	3	-3 6 2
		2.6998	32	3 2 0	4	1.7736	1.7826	3	5 1 0	6	1.4639	1.4684	3	-1 4 3
13	2.584	2.5808	16	-1 4 1			1.7680	10	-4 2 2			1.4547	4	-4 1 3
		2.5452	6	2 4 0			1.7458	4	3 0 2			1.4536	3	2 0 3
12	2.498	2.4959	16	-1 0 2	7	1.7424	1.7416	8	-3 4 2			1.4511	5	-2 4 3
11	2.452	2.4585	12	0 0 2			1.7286	5	3 1 2	5	1.4451	1.4395	4	1 8 1
		2.4464	7	-1 1 2	4	1.7079	1.7091	6	3 5 1			1.4326	3	0 7 2
		2.4111	5	0 1 2	7	1.6746	1.6837	5	-4 3 2	6	1.4251	1.4275	3	4 3 2
4	2.364	2.3692	3	3 0 1			1.6685	6	1 5 2			1.4251	7	-6 3 1
		2.2153	8	4 1 0			1.6594	4	0 7 1			1.4113	3	-2 7 2
30	2.205	2.2057	5	0 5 1	26	1.6519	1.6542	5	-5 3 1			1.4063	3	-6 0 2
		2.2005	5	-2 2 2			1.6479	6	-4 5 1			1.3759	4	2 8 1
		2.1861	22	-1 5 1			1.6390	3	0 0 3	14	1.3740	1.3720	4	3 8 0
												1.3650	3	4 4 2

Table 4. Crystal data, data collection and refinement details.

## Crystal data

Formula	PbMn <sup>2+</sup> <sub>2</sub> Fe <sup>3+</sup> <sub>2</sub> (PO <sub>4</sub> ) <sub>3</sub> (OH) <sub>3</sub>
Space group	<i>P</i> 2 <sub>1</sub> / <i>m</i>
<i>a, b, c</i> (Å)	9.1765(18), 12.340(3), 5.0092(10)
β (°)	101.01(3)
<i>V</i> (Å <sup>3</sup> ), <i>Z</i>	556.8(2), 2
<i>F</i> (000)	702.0
μ (mm <sup>-1</sup> )	20.344
Absorption correction	multi-scan, <i>T</i> <sub>min</sub> , <i>T</i> <sub>max</sub> = 0.34, 0.43
Crystal dimensions (μm)	0.25 x 0.20 x 0.05

#### Data collection

Diffractometer	Dectris EigerX 16M
Temperature (K)	100
Radiation	Synchrotron, λ = 0.710754 Å
Crystal detector distance (mm)	108.033
θ range (°)	2.80 - 28.70
<i>h, k, l</i> ranges	-12 → 12, -16 → 16, -6 → 6
Total reflections measured	9136
Unique reflections	1417 ( <i>R</i> <sub>int</sub> = 0.0355)

#### Refinement

Refinement on	<i>F</i> <sup>2</sup>
<i>R</i> 1 for <i>F</i> <sub>0</sub> > 4σ( <i>F</i> <sub>0</sub> ).	2.07%
w <i>R</i> 2 <sup>†</sup> for all <i>F</i> <sub>0</sub> <sup>2</sup>	7.23%
Reflections used <i>F</i> <sub>0</sub> > 4σ( <i>F</i> <sub>0</sub> )	1417
Number of parameters refined	119
Goof	0.927
(Δ/σ) <sub>max</sub>	0.000
Δρ <sub>max</sub> , Δρ <sub>min</sub> (e/Å)	1.119, -2.248

<sup>†</sup>w*R*2 = Σw(|*F*<sub>o</sub>|<sup>2</sup> - |*F*<sub>c</sub>|<sup>2</sup>)<sup>2</sup> / Σw|*F*<sub>o</sub>|<sup>2</sup>;<sup>1/2</sup>; w = 1/[σ<sup>2</sup>(*F*<sub>o</sub><sup>2</sup>) + (0.0646 *P*)<sup>2</sup> + 2.08 *P*];  
*P* = ([max of (0 or *F*<sub>o</sub><sup>2</sup>)] + 2*F*<sub>c</sub><sup>2</sup>) / 3



Table 5. Fractional atomic coordinates and displacement parameters for plumboperloffite.

ATOM	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> <sub>eq</sub>	<i>U</i> <sup>11</sup>	<i>U</i> <sup>22</sup>	<i>U</i> <sup>33</sup>	<i>U</i> <sup>12</sup>	<i>U</i> <sup>13</sup>	<i>U</i> <sup>23</sup>
<i>X</i>	0.53089(2)	0.75	0.81612(4)	0.01373(13)	0.01212(16)	0.02002(17)	0.00940(17)	0	0.00298(10)	0
<i>M1</i>	0.29861(6)	-0.10841(4)	0.21722(12)	0.00569(16)	0.0056(3)	0.0053(3)	0.0063(3)	0.00021(18)	0.0016(2)	0.00053(18)
<i>M2</i>	0.09489(6)	0.39784(4)	0.13454(11)	0.00540(15)	0.0048(3)	0.0068(3)	0.0049(3)	0.00017(16)	0.00149(19)	-0.00041(16)
<i>P1</i>	0.16275(14)	0.75	0.6835(3)	0.0038(2)	0.0041(5)	0.0049(5)	0.0027(6)	0	0.0010(4)	0
<i>P2</i>	0.33514(10)	0.44237(7)	0.71734(19)	0.0052(2)	0.0050(4)	0.0066(4)	0.0042(4)	0.0007(3)	0.0015(3)	0.0010(3)
<i>O1</i>	0.2878(4)	0.75	0.9401(8)	0.0064(7)	0.0072(16)	0.0076(16)	0.0040(17)	0	0.0004(13)	0
<i>O2</i>	0.2374(4)	0.75	0.4344(8)	0.0089(12)	0.0119(19)	0.0091(18)	0.008(2)	0	0.0084(14)	0
<i>O3</i>	0.0698(3)	0.6478(2)	0.6877(6)	0.0069(5)	0.0077(11)	0.0061(10)	0.0083(12)	-0.0016(9)	0.0053(10)	-0.0012(9)
<i>O4</i>	0.3775(3)	0.5521(2)	0.6183(6)	0.0095(5)	0.0103(12)	0.0096(12)	0.0089(13)	0.0041(10)	0.0025(10)	0.0004(10)
<i>O5</i>	0.2655(3)	0.4584(2)	0.9764(6)	0.0079(5)	0.0083(12)	0.0120(12)	0.0040(12)	-0.0014(9)	0.0022(10)	-0.0003(9)
<i>O6</i>	0.2256(3)	0.3832(2)	0.4964(6)	0.0103(5)	0.0101(12)	0.0134(12)	0.0063(13)	-0.0012(10)	-0.0010(10)	-0.0006(10)
<i>O7</i>	0.4718(3)	0.3689(2)	0.8075(6)	0.0086(5)	0.0051(12)	0.0102(12)	0.0099(13)	0.0031(10)	0.0000(10)	0.0019(9)
<i>OH8</i>	0.1341(5)	0.25	0.0026(9)	0.0096(8)	0.0123(18)	0.0087(17)	0.0075(18)	0	0.0015(15)	0
<i>OH9</i>	0.0617(3)	0.5601(2)	0.1952(6)	0.0066(5)	0.0070(11)	0.0086(12)	0.0039(12)	-0.0021(9)	0.0007(9)	0.0007(9)
<i>H8</i>	0.204(9)	0.25	-0.08(2)	0.032						
<i>H9</i>	0.053(8)	0.590(6)	0.334(10)	0.032						

Prepublished Article

Table 6. Selected interatomic distances (Å), angles (°) and hydrogen bonds for plumboperloffite.

<i>X</i>	O7	2.392(3) (x2)	<i>M2</i>	O3	1.977(3)
	O1	2.427(4)		O6	1.982(3)
	O4	2.897(3) (x2)		OH8	1.996(2)
	O2	2.992(3)		O5	2.028(3)
	O8	3.034(4)		OH9	2.039(3)
	O5	3.232(3) (x2)		OH9	2.057(3)
	O6	3.388(3) (x2)		< <i>M-O</i> >	2.013
	O7	3.447(3) (x2)			
	< <i>X-O</i> >	3.012			
<i>M1</i>			<i>P1</i>	O3	1.525(3) (x2)
	O4	2.119(3)		O2	1.534(4)
	O7	2.153(3)		O1	1.551(4)
	O2	2.187(2)	< <i>P-O</i> >	1.534	
	O5	2.198(3)			
	O1	2.222(2)	<i>P2</i>	O4	1.518(3)
	OH9	2.236(3)		O6	1.531(3)
	< <i>M-O</i> >	2.186		O7	1.544(3)
				O5	1.565(3)
				< <i>P-O</i> >	1.540

## Hydrogen bonds

<i>D-H...A</i>	<i>D-H</i>	<i>H...A</i>	<i>D...A</i>	$\angle D-H...A$
OH8-H8...O6	0.81(3)	2.73(6)	3.263(5)	124(6)
OH9-H9...O3	0.81(3)	1.89(3)	2.682(4)	168(8)

Table 7. Bond-valence\* analysis for plumboperloffite.

	<i>X</i>	<i>M1</i>	<i>M2</i>	<i>P1</i>	<i>P2</i>	H1	H2	Sum
O1	0.41	0.31		1.20				1.93
O2	0.11	0.34		1.26				1.71
O3			0.56	1.28 x2↓			0.21	2.05
O4	0.14 x2↓	0.40			1.30			2.00
O5	0.07	0.33	0.48		1.17			2.05
O6	0.05		0.55		1.26	0.03		1.89
O7	0.44 x2↓	0.41			1.22			2.03
		0.04 x2↓						
OH8	0.10		0.53 x2→			0.97		2.14
OH9		0.30	0.47				0.79	2.01
			0.45					
Sum	2.10	2.05	3.04	5.02	4.95	1.00	1.00	

\*Bond-valence parameters used are from Gagne and Hawthorne (2015). Bond-valences for the *X*, *M1* and *M2* sites are based on the refined occupancy.