

subspecialty choice. Responses were recorded and transcribed verbatim prior to thematic analysis, including triangulation via a co-coding process to check areas were not over-represented and/or subject to researcher thoughts and possible unconscious biases. **Result.** Semi-structured interviews included all North-East C&A trainees and 3 recently qualified Consultants.

Demographics results included 5 male and 6 female participants and 6 non-UK graduates. Emergent themes were analysed into key findings including:

- interest in children/specialty
- specialty experience
- supportive supervisors/team
- work-life balance
- opportunity to impact/intervene early.

Limitations include study of a single geographical area and possible researcher bias.

Conclusion. This Research provides novel findings re factors influencing career choice of C&A Psychiatry to inform future recruitment and retention. Clear themes have emerged re important recruitment/retention factors and the study highlights need for more research to investigate reasons why C&A is not chosen.

The feasibility and acceptability of a peer-support group for mental health in Filipino prisons

Bethany Platford

Nuffield Centre, Faculty of International Health and Development, School of Medicine, University of Leeds

doi: 10.1192/bjo.2021.747

Aims. To assess the feasibility and acceptability of implementing and facilitating a peer-support group for mental health in Filipino prisons.

Objectives. To identify the logistical issues faced in implementing and facilitating healthcare in Filipino Prisons

To explore attitudes of potential participants towards the implementation of a peer-support group for mental health in the prisons

To provide logical recommendations from my findings to inform future mental health support for prisoners in the Philippines

Method. Ethical approval was granted by the University of Leeds prior to data collection. In-country ethical approval was granted through my host Dr Rachael Pickering. Data collection occurred through observations and semi-structured interviews. Participants recruited included six secure-environment healthcare workers, eight prisoners and six ex-prisoners. Both prisoners and ex-prisoners were identified through gatekeepers and informed consent was gained. Interviews were transcribed before coding and themes identified.

Result. Feasibility: Bureaucracy and corruption were the main barriers to the potential successful implementation of a peer-support group, which were emergent themes. Space, time and staff were all themes identified that may help facilitate the group.

Attitudes: It was noted that there is a significant lack of knowledge surrounding what mental health is and its causes. Stigma and discriminatory actions were also noted by many participants as barrier to the group. However, seven prisoners and five ex-prisoners said they would join.

Conclusion. The findings highlighted many barriers but with persistence and local cultural competence this peer-support group could be feasible and be accepted in Filipino prisons. It will be reducing a high unmet need for mental health services in these prisons and if ran successfully with positive effects, will be an example for other prisons across the Philippines and other low-middle income countries.

Towards peak experience - exploring the influence of wilderness on conscious awareness

Oli Purnell

Swansea University Medical School

doi: 10.1192/bjo.2021.748

Aims. This study examines what effect wilderness has on our conscious awareness, and by extension of that meta-cognition; our physical, mental and spiritual health. It reviews available scientific and artistic literature and integrates this with interviews in order to generate original grounded theory. It was found that intensity of wilderness experience varied proportionally with four mediators; Challenge, Immersion, Beauty and Time. With these maximised, experiences broadly within four themes occurred; Increased Awareness, Confidence, Perspective and Connectedness. When sufficiently intense, these four experiences amalgamated to elicit what Maslow described as 'Peak Experience'. As such, this thesis unexpectedly provides a pragmatic recipe towards peak experience, and a map of one's potential psychological journey, in the context of wilderness.

Exploring the influence of wilderness on conscious awareness

Oli Purnell

Swansea University Medical School

doi: 10.1192/bjo.2021.749

Aims. This study examined what effect wilderness has on our conscious awareness, and by extension of that meta-cognition; our physical, mental and spiritual health.

Background.

Method. It reviews available scientific and artistic literature and integrates this with interviews in order to generate original grounded theory.

Result. It was found that intensity of wilderness experience varied proportionally with four mediators; Challenge, Immersion, Beauty and Time. With these maximised, experiences broadly within four themes occurred; Increased Awareness, Confidence, Perspective and Connectedness. When sufficiently intense, these four experiences amalgamated to elicit what Maslow described as 'Peak Experience'.

Conclusion. As such, this thesis unexpectedly provides a pragmatic recipe towards peak experience, and a map of one's potential psychological journey, in the context of wilderness.

Admission patterns in a psychiatric intensive care unit in Ireland: a longitudinal follow-up

Shaeraine Raaj^{1*}, Sujesha Navanathan², Basil Matti³, Anisha Bhagawan⁴, Pauline Twomey⁵, John Lally⁶ and Roy Browne⁵

¹Mater Misericordiae University Hospital, Phoenix Care Centre;

²National Forensic Mental Health Services; ³St. Patrick's University Hospital; ⁴national drug treatment centre; ⁵Phoenix Care Centre and

⁶Mater Misericordiae University Hospital

*Corresponding author.

doi: 10.1192/bjo.2021.750

Aims. This observational study aims to describe the course of the admission and clinical characteristics of admissions to the PICUs in the Phoenix Care Centre, Dublin, Ireland. The authors hypothesised that the length of stay (LOS) would be shorter in male patients as compared to females.

Method. This retrospective cohort study was carried out at the Phoenix Care Centre Dublin, Ireland. Informed consent was not sought as this was a retrospective chart study involving anonymised clinical data which was collected as part of routine clinical care and no items of information were reported that would enable the identification of any subject. We described primary outcomes using frequencies, percentages, mean and standard deviations, median and interquartile ranges (IQR). Between groups comparisons were made using χ^2 tests for categorical variables; t-tests, ANOVA tests, or Kruskal-Wallis tests, for continuous variables; All analyses were two-tailed, and a P-value ≤ 0.05 was considered statistically significant

Result. Over the study period from Jan 2014 to Jan 2017 inclusive, there were 96 admission episodes to the PICU. The mean age of admitted cases was 37.1 (SD = 11.3) years (range 18–63 years). The mean length of stay (LOS) was 59.3 (SD = 61.0) days (median 39.5 days). All patients were admitted under the Mental Health Act legislation. We identified assault as the primary risk factor for pre-admission 62% (n = 62) to the PICU. Antipsychotic polypharmacy was used in 61% (n = 55) of the admission. The mean daily antipsychotic dosage was 139.4 % (SD = 65.1) of BNF maximum daily dose. A diagnosis of acute psychotic disorder (B = -1.027, p = 0.003, 95% CI: -1.691 to -0.363) was associated with reduced LOS in PICU. Majority of admissions 43% (n = 39) had a diagnosis of schizophrenia, followed by Bipolar affective disorder BPAD 21% (n = 21), schizoaffective disorder 18% (n = 18), and acute psychotic disorder 9% (n = 9).

Conclusion. Psychiatric Intensive Care Unit is an essential service for the severely ill psychiatric patients and is a progressively developing sub-speciality. An important finding from our study describes the cohort of patients admitted being predominantly male, younger-aged, single, with a diagnosis of schizophrenia, legally detained, and from an Irish background. The primary indication for a referral is the risk of assault, showing the need for the intensive and secure treatment model that a PICU can provide.

Prevalence and correlates of depressive symptoms among professional drivers in Saudi Arabia: a cross-sectional study

Adnan Raed Alnaser*, Osama A. Zitoun, Tawfik Rajab, Abdullah Khojah, Juliann Saquib and Nazmus Saquib

College of Medicine, Sulaiman Al Rajhi University

*Corresponding author.

doi: 10.1192/bjo.2021.751

Aims. Due to the nature of their work, professional drivers face a considerable risk of developing depression and other mental illnesses. We sought to assess the prevalence and the factors influencing depressive symptoms among professional drivers in Saudi Arabia.

Method. Using convenience sampling, we have conducted an interviewer-administered survey on 324 professional drivers in Qassim Region in Saudi Arabia using Depression subscale from the Depression, Anxiety and Stress Scale 21 (DASS-21). Participants were interviewed in their native language, and responses were outlined directly into an online form in English. Data were then extracted and analyzed using SPSS software.

Result. Participants' mean age was 38.6 years, and mean driving hours per day were 9.86 hours/day. The mean DASS-21 depression score among the professional drivers was 2.88. Overall, 21.9% of the included drivers had variable degrees of depressive symptoms, with 7.4% suffered from extremely severe symptoms. Depressive symptoms were influenced by the driver's nationality, educational level, vehicle type, driving years, BMI, presence of chronic medical conditions, physical activity, and sexual activity. Moreover, poor sleep quality increased the risk of developing

depressive symptoms among the drivers by 31.9 times (OR: 31.9, CI: 9.03–112.63, P < 0.001).

Conclusion. Nearly one-fifth of professional drivers in Saudi Arabia (Qassim region) suffer from depressive symptoms. Unhealthy life-style practices (i.e. being obese and physically inactive) have been closely related to depressive symptoms. Education, sexual activity, type of driven vehicle, and the number of chronic conditions were also associated with depressive symptoms. Also, poor and fair sleep quality was strongly associated with the development of depressive symptoms as compared with excellent sleep quality. As drivers are always on the move and hardly reachable, we would propose psychological support and counseling to be administered via telemedicine services. Future research is needed to better comprehend the needs of this vulnerable population.

Can probiotics benefit young people with autism spectrum disorders?

Pooja Ramani^{1*} and Regina Sala²

¹Great Ormond Street Hospital and ²East London NHS Foundation Trust

*Corresponding author.

doi: 10.1192/bjo.2021.752

Aims. The aims are to evaluate the effectiveness of Probiotics on young people with Autism Spectrum Disorder.

We hypothesized that there will be an improvement of the comorbid gastrointestinal symptoms that can accompany Autism Spectrum Disorder.

We believe that the use of probiotics can exert bidirectional effects on the gut-brain axis which may result in improvements in core Autism symptoms.

Method. A literature search was performed in accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. We used databases including OVID MEDLINE, Pubmed, EMBASE, AMED and the Cochrane register of controlled trials. Studies using Probiotics as a treatment for children with ASD were identified by key search terms; Child*, young person*, adoles*, teenagers, ASD, Autism Spectrum Disorder, Autism, Pervasive developmental disorder, PDD, Probiotics, Supplements, Lactobacillus, and Bifidobacterium. Inclusion criteria: Children of age range 2-18 with a diagnosis of ASD and having at least one gastrointestinal symptom were included. Exclusion criteria: The following were excluded: studies looking at Autism with interventions aside from Probiotics; studies where Probiotics were tested in conjunction with other interventions; studies where there were additional neurodevelopmental disorders.

Result. Twelve studies identified all utilized probiotics. This included 7 Randomised Control Trials, 2 Open-Label studies, 1 pre and post-intervention design and 1 Case study. All RCTs gave probiotics or placebo to children.

Ten studies showed an improvement in gastrointestinal symptoms. Six studies showed improvements in various behavioral measures. Four studies showed improvements in core autism symptoms. However, the sample sizes in these studies were not large enough to prove statistical significance.

Conclusion. No studies showed an adverse reaction which indicates probiotics can be considered a safe treatment.

The improvements in a variety of parameters imply probiotics a suitable adjunctive intervention that may help improve ASD core symptoms in young people as well as improving physical and behavioural comorbidities which in some cases was noted by parents.

However, due to high dropout rates and generally small sample sizes, larger-scale trials are needed to critically confirm the efficacy of probiotics for children with ASD.