

CORRIGENDUM


Psychophysical analysis of contrast processing segregated into magnocellular and parvocellular systems in asymptomatic carriers of 11778 Leber's hereditary optic neuropathy

M. GUALTIERI,^{1,2} M. BANDEIRA,^{1,2} R.D. HAMER,^{2,8} M.E. COSTA,^{1,2} A.G.F. OLIVEIRA,^{1,2}
A.L.A. MOURA,^{1,2} F. SADUN,³ A.M. DE NEGRI,⁴ A. BEREZOVSKY,⁵ S.R. SALOMÃO,⁵ V. CARELLI,⁶
A.A. SADUN,⁷ AND D.F. VENTURA^{1,2}

¹Departamento de Psicologia Experimental, Universidade de São Paulo, São Paulo, Brasil
²Centro de Neurociências e Comportamento, Universidade de São Paulo, São Paulo, Brasil
³Ospedale S. Giovanni Evangelista, Tivoli, Italy
⁴Azienda Ospedaliera S. Camillo-Forlanini, Roma, Italy
⁵Departamento de Oftalmologia, Universidade Federal de São Paulo, São Paulo, Brasil
⁶Department of Neurology, University of Bologna, Bologna, Italy
⁷Doheny Eye Institute, Keck-USC School of Medicine, Los Angeles, California
⁸Smith-Kettlewell Eye Research Institute, San Francisco, California

(This article appeared in Volume 25, Number 3, 2008, pp. 469–474).

The abscissa in Fig. 1 (p. 471) was labeled in units of cd/m^2 . The correct units are trolands (Td), and a new Fig. 1, so labeled, is shown below.



On p. 473, column 1, the third sentence in the second paragraph under the heading *Effects of stimulus duration* should read as follows:

“The mean Tc-estimates for their two observers were 33.5 ms for SPP and 49 ms for PPP, compared with 26.3 (SPP) versus 38.4 (PPP) for our controls.”